

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

A9

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification : <b>C12Q</b>	<b>A2</b>	(11) International Publication Number: <b>WO 00/08202</b> (43) International Publication Date: 17 February 2000 (17.02.00)
(21) International Application Number: PCT/US99/17845 (22) International Filing Date: 4 August 1999 (04.08.99) (30) Priority Data: 09/129,256      4 August 1998 (04.08.98)      US 60/095,470      5 August 1998 (05.08.98)      US 60/102,178      28 September 1998 (28.09.98)      US 60/116,107      15 January 1999 (15.01.99)      US (71) Applicant (for all designated States except US): SUGEN, INC. [US/US]; 230 East Grand Avenue, South San Francisco, CA 94080-4811 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): TANG, Peng, Cho [-/US]; 827 Camino Ricardo, Moraga, CA 94556 (US). SUN, Li [-/US]; 64 Rockharbor Lane, Foster City, CA 94556 (US). MILLER, Todd, Anthony [-/US]; 4928 Colusa Street, Union City, CA 94587 (US). LIANG, Congxin [-/US]; 729 West Remington Drive, Sunnyvale, CA 94087 (US). TRAN, Ngoc, My [-/US]; 550 Shannon Way, Spt. #6306, Redwood City, CA 94065 (US). NGUYEN, Anh, Thi [-/US]; 503 Becado Place, Fremont, CA 94539 (US). NEMATALLA, Asaad [-/US]; 1324 Bodega Place, Walnut Creek, CA 94596 (US).		(74) Agents: ROSE, Bernard, F. et al.; Lyon & Lyon LLP, Suite 4700, 633 West Fifth Street, Los Angeles, CA 90071-2066 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report.
(54) Title: 3-METHYLIDENYL-2-INDOLINONE MODULATORS OF PROTEIN KINASE (57) Abstract <p>The present invention relates to novel 3-methylidenyl-2-indolinone compounds and physiologically acceptable salts and prodrugs thereof which modulate the activity of protein kinases and therefore are expected to be useful in the prevention and treatment of protein kinase related cellular disorders such as cancer.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

WO 00/08202

1

PCT/US99/17845 ,

## DESCRIPTION

### 5 3-METHYLIDENYL-2-INDOLINONE MODLATORS OF PROTEIN KINASE

#### INTRODUCTION

The present invention relates generally to organic chemistry, biochemistry, pharmacology and medicine. More particularly, it relates to 3-methylidenyl-2-indolinone derivatives, and their physiologically acceptable salts and prodrugs, which modulate the activity of protein kinases ("PKs") and are expected to exhibit a salutary effect against disorders related to abnormal PK activity.

15

#### BACKGROUND OF THE INVENTION

The following is offered as background information only and is not admitted to be prior art to the present invention.

PKs are enzymes that catalyze the phosphorylation of hydroxy groups on tyrosine, serine and threonine residues of proteins. The consequences of this seemingly simple activity are staggering; cell growth, differentiation and proliferation, i.e., virtually all aspects of cell life in one way or another depend on PK activity. Furthermore, abnormal PK activity has been related to a host of disorders, ranging from relatively non-life threatening diseases such as psoriasis to extremely virulent diseases such as glioblastoma (brain cancer).

The PKs can be conveniently be broken down into two classes, the protein tyrosine kinases (PTKs) and the serine-threonine kinases (STKs).

One of the prime aspects of PTK activity is their involvement with growth factor receptors. Growth factor receptors are cell-surface proteins. When bound by a growth



WO 00/08202

2

PCT/US99/17845,

factor ligand, growth factor receptors are converted to an active form which interacts with proteins on the inner surface of a cell membrane. This leads to phosphorylation on tyrosine residues of the receptor and other proteins and to the

5 formation inside the cell of complexes with a variety of cytoplasmic signaling molecules that, in turn, effect numerous cellular responses such as cell division (proliferation), cell differentiation, cell growth, expression of cellular metabolic processes to the extracellular microenvironment, etc. For a

10 more complete discussion, see Schlessinger and Ullrich, Neuron, 9:303-391 (1992) which is incorporated by reference, including any drawings, as if fully set forth herein.

Growth factor receptors with PTK activity are known as receptor tyrosine kinases ("RTKs"). They comprise a large

15 family of transmembrane receptors with diverse biological activity. At present, at least nineteen (19) distinct subfamilies of RTKs have been identified. An example of these is the subfamily designated the "HER" RTKs, which include EGFR (epithelial growth factor receptor), HER2, HER3 and HER4.

20 These RTKs consist of an extracellular glycosylated ligand binding domain, a transmembrane domain and an intracellular cytoplasmic catalytic domain that can phosphorylate tyrosine residues on proteins.

Another RTK subfamily consists of insulin receptor (IR),

25 insulin-like growth factor I receptor (IGF-1R) and insulin receptor related receptor (IRR). IR and IGF-1R interact with insulin, IGF-I and IGF-II to form a heterotetramer of two entirely extracellular glycosylated  $\alpha$  subunits and two  $\beta$  subunits which cross the cell membrane and which contain the

30 tyrosine kinase domain.

A third RTK subfamily is referred to as the platelet derived growth factor receptor ("PDGFR") group, which includes PDGFR $\alpha$ , PDGFR $\beta$ , CSFIR, c-kit and c-fms. These receptors

WO 00/08202

3

PCT/US99/17845

consist of glycosylated extracellular domains composed of variable numbers of immunoglobulin-like loops and an intracellular domain wherein the tyrosine kinase domain is interrupted by unrelated amino acid sequences.

5 Another group which, because of its similarity to the PDGFR subfamily, is sometimes subsumed into the later group is the fetus liver kinase ("flk") receptor subfamily. This group is believed to be made of up of kinase insert domain-receptor fetal liver kinase-1 (KDR/FLK-1), flk-1R, flk-4 and fms-like  
10 tyrosine kinase 1 (flt-1).

A further member of the tyrosine kinase growth factor receptor family is the fibroblast growth factor ("FGF") receptor subgroup. This group consists of four receptors, FGFR1-4, and seven ligands, FGF1-7. While not yet well  
15 defined, it appears that, like the PDGF receptors, the FGF receptors consist of a glycosylated extracellular domain containing a variable number of immunoglobulin-like loops and an intracellular domain in which the tyrosine kinase sequence is interrupted by regions of unrelated amino acid sequences.

20 Still another member of the tyrosine kinase growth factor receptor family is the vascular endothelial growth factor (VEGF) receptor subgroup. VEGF is a dimeric glycoprotein similar to PDGF but has different biological functions and target cell specificity in vivo. In particular, VEGF is  
25 presently thought to play an essential role in vasculogenesis and angiogenesis.

A more complete listing of the known RTK subfamilies is described in Plowman et al., DN&P, 7(6):334-339 (1994) which is incorporated by reference, including any drawings, as if  
30 fully set forth herein.

In addition to the RTKs, there also exists a family of entirely intracellular PTKs called "non-receptor tyrosine kinases" or "cellular tyrosine kinases." This latter designation, abbreviated "CTK," will be used herein. CTKs do

WO 00/08202

4

PCT/US99/17845 ,

not contain extracellular and transmembrane domains. At present, over 24 CTKs in 11 subfamilies (Src, Frk, Btk, Csk, Abl, Zap70, Fes, Fps, Fak, Jak and Ack) have been identified. The Src subfamily appear so far to be the largest group of CTKs and includes Src, Yes, Fyn, Lyn, Lck, Blk, Hck, Fgr and Yrk. For a more detailed discussion of CTKs, see Bolen, Oncogene, 8:2025-2031 (1993), which is incorporated by reference, including any drawings, as if fully set forth herein.

10       The serine/threonine kinases, STKs, like the CTKs, are predominantly intracellular although there are a few receptor kinases of the STK type. STKs are the most common of the cytosolic kinases; i.e., kinases that perform their function in that part of the cytoplasm other than the cytoplasmic  
15 organelles and cytoskelton. The cytosol is the region within the cell where much of the cell's intermediary metabolic and biosynthetic activity occurs; e.g., it is in the cytosol that proteins are synthesized on ribosomes.

RTKs, CTKs and STKs have all been implicated in a host of  
20 pathogenic conditions including, significantly, cancer. Other pathogenic conditions which have been associated with PTKs include, without limitation, psoriasis, hepatic cirrhosis, diabetes, angiogenesis, restenosis, ocular diseases, rheumatoid arthritis and other inflammatory disorders,  
25 immunological disorders such as autoimmune disease, cardiovascular disease such as atherosclerosis and a variety of renal disorders.

With regard to cancer, two of the major hypotheses advanced to explain the excessive cellular proliferation that  
30 drives tumor development relate to functions known to be PK regulated. That is, it has been suggested that malignant cell growth results from a breakdown in the mechanisms that control cell division and/or differentiation. It has been shown that the protein products of a number of proto-oncogenes are

WO 00/08202

5

PCT/US99/17845 ,

involved in the signal transduction pathways that regulate cell growth and differentiation. These protein products of proto-oncogenes include the extracellular growth factors, transmembrane growth factor PTK receptors (RTKs), cytoplasmic

5 PTKs (CTKs) and cytosolic STKs, discussed above.

In view of the apparent link between PK-related cellular activities and wide variety of human disorders, it is no surprise that a great deal of effort is being expended in an attempt to identify ways to modulate PK activity. Some of

10 these have involved biomimetic approaches using large molecules patterned on those involved in the actual cellular processes (e.g., mutant ligands (U.S. App. No. 4,966,849); soluble receptors and antibodies (App. No. WO 94/10202, Kendall and Thomas, Proc. Nat'l Acad. Sci., 90:10705-09

15 (1994), Kim, et al., Nature, 362:841-844 (1993)); RNA ligands (Jelinek, et al., Biochemistry, 33:10450-56); Takano, et al., Mol. Bio. Cell 4:358A (1993); Kinsella, et al., Exp. Cell Res. 199:56-62 (1992); Wright, et al., J. Cellular Phys., 152:448-57) and tyrosine kinase inhibitors (WO 94/03427; WO 92/21660; WO 91/15495; WO 94/14808; U.S. Pat. No. 5,330,992; Mariani, et

20 al., Proc. Am. Assoc. Cancer Res., 35:2268 (1994)).

In addition to the above, attempts have been made to identify small molecules which act as PK inhibitors. For example, bis- monocyclic, bicyclic and heterocyclic aryl

25 compounds (PCT WO 92/20642), vinylene-azaindole derivatives (PCT WO 94/14808) and 1-cyclopropyl-4-pyridylquinolones (U.S. Pat. No. 5,330,992) have been described as tyrosine kinase inhibitors. Styryl compounds (U.S. Pat. No. 5,217,999), styryl-substituted pyridyl compounds (U.S. Pat. No.

30 5,302,606), quinazoline derivatives (EP App. No. 0 566 266 A1), selenaindoles and selenides (PCT WO 94/03427), tricyclic polyhydroxylic compounds (PCT WO 92/21660) and benzylphosphonic acid compounds (PCT WO 91/15495) have all

WO 00/08202

6

PCT/US99/17845 ,


been described as PTK inhibitors useful in the treatment of cancer.

#### SUMMARY OF THE INVENTION

Our own efforts to identify small organic molecules which  
5 modulate PK activity and which, therefore, are expected to be  
useful in the treatment and prevention of disorders involving  
abnormal PK activity, has led us to the discovery of a family  
of 3-methylidenyl-2-indolinone derivatives and their prodrugs  
and physiologically acceptable salts, which exhibit PK  
10 modulating ability and are thereby expected to have a salutary  
effect against disorders related to abnormal PK activity; it  
is these compounds which is the subject of this invention.

Thus, the present invention relates generally to 3-  
methylidenyl-2-indolinones and their prodrugs and  
15 physiologically acceptable salts, which modulate the activity  
of receptor tyrosine kinases (RTKs), non-receptor protein  
tyrosine kinases (CTKs) and serine/threonine protein kinases  
(STKs). In addition, the present invention relates to the  
preparation and use of pharmaceutical compositions of the  
20 disclosed compounds and their physiologically acceptable salts  
and prodrugs in the treatment or prevention of PK driven  
disorders such as, by way of example and not limitation,  
cancer, diabetes, hepatic cirrhosis, cardiovasacular disease  
such as atherosclerosis, angiogenesis, immunological disease  
25 such as autoimmune disease and renal disease.

The terms "2-indolinone," indolin-2-one and "2-oxindole"  
are used interchangeably herein to refer to a molecule having  
the chemical structure:




WO 00/08202

7

PCT/US99/17845 ,

where A is carbon or nitrogen.

A "3-methylidenyl-2-indolinone" refers to a molecule having the chemical structure:



5

where, again, A can be carbon or nitrogen.

A "pharmaceutical composition" refers to a mixture of one or more of the compounds described herein, or physiologically acceptable salts or prodrugs thereof, with other chemical components, such as physiologically acceptable carriers and excipients. The purpose of a pharmaceutical composition is to facilitate administration of a compound to an organism.

A "prodrug" refers to an agent which is converted into the parent drug in vivo. Prodrugs are often useful because, in some situations, they may be easier to administer than the parent drug. They may, for instance, be bioavailable by oral administration whereas the parent drug is not. The prodrug may also have improved solubility in pharmaceutical compositions over the parent drug. An example, without limitation, of a prodrug would be a compound of the present invention which is administered as an ester (the "prodrug") to facilitate transmittal across a cell membrane where water solubility is detrimental to mobility but then is metabolically hydrolyzed to the carboxylic acid, the active entity, once inside the cell where water solubility is beneficial.

A further example of a prodrug might be a short polypeptide, for example, without limitation, a 2 - 10 amino acid polypeptide, bonded through a terminal amino group to a

WO 00/08202

8

PCT/US99/17845

carboxy group of a compound of this invention wherein the polypeptide is hydrolyzed or metabolized in vivo to release the active molecule.

As used herein, a "physiologically acceptable carrier" or  
5 a "pharmaceutically acceptable carrier" refers to a carrier or diluent that does not cause significant irritation to an organism and does not abrogate the biological activity and properties of the administered compound.

A "carrier" refers to a chemical compound which  
10 facilitates the incorporation of a compound of interest into cells and tissues. An example, without limitation, of a carrier is dimethyl sulfoxide (DMSO).


A "diluent" refers to a chemical compound, usually a liquid, which dissolves or disperses a compound of interest  
15 thereby reducing the concentration of the compound to less than that of the compound alone.

An "excipient" refers to an inert substance added to a pharmaceutical composition to further facilitate  
administration of a compound. Examples, without limitation,  
20 of excipients include calcium carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils and polyethylene glycols.

## 1. CHEMISTRY

### A. General Structural Features.

25 In one aspect, the the present invention relate to 2-indolinones having chemical structure 1:



WO 00/08202


9

PCT/US99/17845 ,

wherein:

A is selected from the group consisting of carbon and  
5 nitrogen;

Q is selected from the group consisting of



10 bonds a and b may be either single or double bonds as indicated by the dotted lines with the proviso that a and b are both single bonds or both double bonds in any one compound of this invention;

15 r is 1 when a and b are double bonds;

r is 2 when a and b are single bonds;

R<sup>0</sup> is selected from the group consisting of hydrogen, alkyl,  
20 -C(O)OR<sup>19</sup> and -C(O)R<sup>19</sup>;

R<sup>1</sup> is selected from the group consisting of hydrogen, alkyl, aryl, alkoxy, halo, -C(O)OR<sup>19</sup>, -(CH<sub>2</sub>)<sub>n</sub>OC(O)R<sup>19</sup>, -C(O)NR<sup>19</sup> and (CH<sub>2</sub>)<sub>n</sub>R<sup>20</sup>, wherein:

25

R<sup>19</sup> is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl and aryl;

n is 1, 2, 3 or 4;



WO 00/08202

10

PCT/US99/17845 ,

$R^{20}$  is selected from the group consisting of hydroxy, halo,  $-\text{OC}(\text{O})\text{NR}^{21}\text{R}^{22}$ ;  $-\text{OC}(\text{S})\text{NR}^{21}\text{R}^{22}$ ;  $-\text{OC}(\text{O})\text{NHSO}_2\text{R}^{21}$ ,  $-\text{C}(\text{O})\text{OR}^{19}$ ,  $-\text{NR}^{21}\text{R}^{22}$  and a heteroalicyclic group containing at least one  
5 nitrogen atom in the ring, the ring being bonded to the adjacent  $\text{CH}_2$  group through the nitrogen atom;

$R^2$  is selected from the group consisting of hydrogen, alkyl, trihalomethyl, aryl, heteroaryl, heteroalicyclic, alkoxy,  
10 halo,  $-(\text{CH}_2)_n\text{R}^{20}$ ,  $-\text{SO}_2\text{NR}^{21}\text{R}^{22}$ ,  $-\text{C}(\text{O})\text{OR}^{19}$ ,  $-\text{C}(\text{O})\text{R}^{19}$ ,  $-\text{NHC}(\text{O})\text{OR}^{19}$ ,  $\text{NHC}(\text{O})\text{R}^{19}$ ,  $-\text{C}(\text{O})(\text{CH}_2)_n\text{R}^{20}$ ,  $-\text{NR}^{21}\text{R}^{22}$  and  $-\text{N}=\text{CR}^{23}$  wherein

$R^{21}$  and  $R^{22}$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl,  
15 cycloalkyl, aryl, heteroaryl and heteroalicyclic;

$R^{23}$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, aryl and heteroaryl;

20  $R^3$  is selected from the group consisting of hydrogen, alkyl, trihalomethyl, alkoxy, aryl, aryloxy, heteroaryl, heteroalicyclic, hydroxy, halo,  $-\text{SO}_2\text{NR}^{21}\text{R}^{22}$ ,  $-\text{NHSO}_2\text{R}^{19}$ ,  $-\text{C}(\text{O})\text{OR}^{19}$ ,  $-\text{NR}^{21}\text{R}^{22}$  and  $-\text{NHC}(\text{O})\text{R}^{24}$ , wherein

25  $R^{24}$  is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl and heteroalicyclic;

$R^4$  is selected from the group consisting of hydrogen, alkyl, alkoxy and halo;

30


$R^5$  is selected from the group consisting of hydrogen, alkyl, cycloalkyl, halo, aryl and heteroaryl;

WO 00/08202

11

PCT/US99/17845

R<sup>6</sup> is selected from the group consisting of hydrogen, alkyl, alkoxy, halo, cycloalkyl, aryl and, combined with R<sup>18</sup>, a heteroalicyclic group having the structure




wherein


5

y and y' are independently selected from the group consisting of hydrogen, alkyl and aryl;

R<sup>7</sup> is OR<sup>18</sup>, wherein R<sup>18</sup> is selected from the group consisting of  
10 alkyl, -(CH)<sub>n</sub>R<sup>20</sup> and, combined with R<sup>6</sup> or R<sup>8</sup>, a heteroalicyclic group having the structure



R<sup>8</sup> is selected from the group consisting of hydrogen, alkyl,  
15 cycloalkyl, alkoxy, halo, aryl, heteroaryl and, combined with R<sup>18</sup>, a heteroalicyclic ring having the structure



WO 00/08202

12

PCT/US99/17845,

R<sup>9</sup> is selected from the group consisting of hydrogen, alkyl, alkoxy, halo and -NR<sup>21</sup>R<sup>22</sup>;

5 R<sup>10</sup> is selected from the group consisting of alkyl and -C(O)OR<sup>19</sup>;

R<sup>11</sup> is selected from the group consisting of hydrogen, alkyl and -C(O)OR<sup>19</sup>;

10 R<sup>12</sup> is -(CH<sub>2</sub>)<sub>n</sub>R<sup>20</sup>;

R<sup>13</sup> is either a bond through which Q is bonded to the rest of the molecule or, when it is not, a group selected from the group consisting of hydrogen and alkyl;

15

R<sup>14</sup> is either a bond through which Q is bonded to the rest of the molecule or, when it is not, a group selected from the group consisting of hydrogen, alkyl and -(CH<sub>2</sub>)<sub>n</sub>R<sup>20</sup>;

20 R<sup>15</sup> is either a bond through which Q is bonded to the rest of the molecule or, when it is not, hydrogen;

25 with the proviso that only one of R<sup>13</sup>, R<sup>14</sup> or R<sup>15</sup> is the bond linking Q to the rest of the molecule in any one compound herein;

R<sup>16</sup> and R<sup>17</sup>, when a and b are single bonds, are independently selected from the group consisting of hydrogen, alkyl and aryl and, when a and b are double bonds, R<sup>16</sup> is hydrogen and R<sup>17</sup> does  
30 not exist; and,

physiologically acceptable salts and prodrugs thereof.

As used herein, the term "alkyl" refers to a saturated aliphatic hydrocarbon including straight chain and branched

WO 00/08202

13

PCT/US99/17845 ,

chain groups. Preferably, the alkyl group has 1 to 20 carbon atoms (whenever a numerical range; e.g. "1-20", is stated herein, it means that the group, in this case the alkyl group, may contain 1 carbon atom, 2 carbon atoms, 3 carbon atoms, etc. up to and including 20 carbon atoms). More preferably, it is a medium size alkyl having 1 to 10 carbon atoms. Most preferably, it is a lower alkyl having 1 to 4 carbon atoms. The alkyl group may be substituted or unsubstituted. When substituted, the substituent group(s) is preferably one or more independently selected from the group consisting of halo hydroxy, unsubstituted lower alkoxy, aryl optionally substituted with one or more halo, hydroxy, unsubstituted lower alkyl or unsubstituted lower alkoxy groups, aryloxy optionally substituted with one or more halo, hydroxy, unsubstituted lower alkyl or unsubstituted lower alkoxy groups, 6-member heteroaryl having from 1 to 3 nitrogen atoms in the ring, the carbons in the ring being optionally substituted with one or more halo, hydroxy, unsubstituted lower alkyl or unsubstituted lower alkoxy groups, 5-member heteroaryl having from 1 to 3 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur, the carbon atoms of the group being optionally substituted with one or more halo, hydroxy, unsubstituted lower alkyl or unsubstituted lower alkoxy groups, 5- or 6-member heteroalicyclic group having from 1 to 3 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur, the carbon and nitrogen (if present) atoms in the group being optionally substituted with one or more halo, hydroxy, unsubstituted lower alkyl or unsubstituted lower alkoxy groups, mercapto, (unsubstituted lower alkyl)thio, arylthio optionally substituted with one or more halo, hydroxy, unsubstituted lower alkyl or unsubstituted lower alkoxy groups, cyano, acyl, thioacyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, nitro, N-sulfonamido, S-sulfonamido,  $R^{25}S(O)-$ ,  $R^{25}S(O)_2-$ ,  $-C(O)OR^{25}$ ,  $R^{25}C(O)O-$ , and

WO 00/08202

14

PCT/US99/17845 ,

-NR<sup>25</sup>R<sup>26</sup>, wherein R<sup>25</sup> and R<sup>26</sup> are independently selected from the group consisting of hydrogen, unsubstituted lower alkyl, trihalomethyl, unsubstituted (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, unsubstituted lower alkenyl, unsubstituted lower alkynyl and aryl optionally  
5 optionally substituted with one or more halo, hydroxy, unsubstituted lower alkyl or unsubstituted lower alkoxy groups.

A "cycloalkyl" group refers to a 3 to 8 member all-carbon monocyclic ring, an all-carbon 5-member/6-member or 6-  
10 member/6-member fused bicyclic ring or a multicyclic fused ring(a "fused" ring system means that each ring in the system shares an adjacent pair of carbon atoms with each other ring in the system)group wherein one or more of the rings may contain one or more double bonds but none of the rings has a  
15 completely conjugated pi-electron system. Examples, without limitation, of cycloalkyl groups are cyclopropane, cyclobutane, cyclopentane, cyclopentene, cyclohexane, cyclohexadiene, adamantane, cycloheptane and, cycloheptatriene. A cycloalkyl group may be substituted or  
20 unsubstituted. When substituted, the substituent group(s) is preferably one or more independently selected from the group consisting of unsubstituted lower alkyl, trihaloalkyl, halo, hydroxy, unsubstituted lower alkoxy, aryl optionally substituted with one or more halo, hydroxy, unsubstituted  
25 lower alkyl or unsubstituted lower alkoxy groups, aryloxy optionally substituted with one or more halo, hydroxy, unsubstituted lower alkyl or unsubstituted lower alkoxy groups, 6-member heteroaryl having from 1 to 3 nitrogen atoms in the ring, the carbons in the ring being optionally  
30 substituted with one or more halo, hydroxy, unsubstituted lower alkyl or unsubstituted lower alkoxy groups, 5-member heteroaryl having from 1 to 3 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur, the carbon atoms of the group being optionally substituted with one or

WO 00/08202

15

PCT/US99/17845 ,

more halo, hydroxy, unsubstituted lower alkyl or unsubstituted lower alkoxy groups, 5- or 6-member heteroalicyclic group having from 1 to 3 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur, the carbon and  
5 nitrogen (if present) atoms in the group being optionally substituted with one or more halo, hydroxy, unsubstituted lower alkyl or unsubstituted lower alkoxy groups, mercapto, (unsubstituted lower alkyl)thio, arylthio optionally substituted with one or more halo, hydroxy, unsubstituted  
10 lower alkyl or unsubstituted lower alkoxy groups, cyano, acyl, thioacyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, nitro, N-sulfonamido, S-sulfonamido,  $R^{25}S(O)-$ ,  $R^{25}S(O)_2-$ ,  $-C(O)OR^{25}$ ,  $R^{25}C(O)O-$ , and  $-NR^{25}R^{26}$  are as defined above.

15 An "alkenyl" group refers to an alkyl group, as defined herein, consisting of at least two carbon atoms and at least one carbon-carbon double bond.

An "alkynyl" group refers to an alkyl group, as defined herein, consisting of at least two carbon atoms and at least  
20 one carbon-carbon triple bond.

An "aryl" group refers to an all-carbon monocyclic or fused-ring polycyclic (i.e., rings which share adjacent pairs of carbon atoms) groups having a completely conjugated pi-electron system. Examples, without limitation, of aryl groups  
25 are phenyl, naphthalenyl and anthracenyl. The aryl group may be substituted or unsubstituted. When substituted, the substituted group(s) is preferably one or more independently selected from the group consisting of unsubstituted lower alkyl, trihaloalkyl, halo, hydroxy, unsubstituted lower  
30 alkoxy, mercapto, (unsubstituted lower alkyl)thio, cyano, acyl, thioacyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, nitro, N-sulfonamido, S-sulfonamido,  $R^{25}S(O)-$ ,  $R^{25}S(O)_2-$ ,  $-C(O)OR^{25}$ ,  $R^{25}C(O)O-$ , and  $-NR^{25}R^{26}$ , with  $R^{25}$  and  $R^{26}$  as defined above.

WO 00/08202

16

PCT/US99/17845.

As used herein, a "heteroaryl" group refers to a monocyclic or fused ring (i.e., rings which share an adjacent pair of atoms) group having in the ring(s) one or more atoms selected from the group consisting of nitrogen, oxygen and sulfur and, in addition, having a completely conjugated pi-electron system. Examples, without limitation, of heteroaryl groups are pyrrole, furan, thiophene, imidazole, oxazole, thiazole, pyrazole, pyridine, pyrimidine, quinoline, isoquinoline, purine and carbazole. The heteroaryl group may be substituted or unsubstituted. When substituted, the substituted group(s) is preferably one or more independently selected from the group consisting of unsubstituted lower alkyl, trihaloalkyl, halo, hydroxy, unsubstituted lower alkoxy, mercapto, (unsubstituted lower alkyl)thio, cyano, acyl, thioacyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, nitro, N-sulfonamido, S-sulfonamido,  $R^{25}S(O)-$ ,  $R^{25}S(O)_2-$ ,  $-C(O)OR^{25}$ ,  $R^{25}C(O)O-$ , and  $-NR^{25}R^{26}$ , with  $R^{25}$  and  $R^{26}$  as defined above.

A "heteroalicyclic" group refers to a monocyclic or fused ring group having in the ring(s) one or more atoms selected from the group consisting of nitrogen, oxygen and sulfur. The rings may also have one or more double bonds. However, the rings do not have a completely conjugated pi-electron system. The heteroalicyclic ring may be substituted or unsubstituted. When substituted, the substituted group(s) is preferably one or more independently selected from the group consisting of unsubstituted lower alkyl, trihaloalkyl, halo, hydroxy, unsubstituted lower alkoxy, mercapto, (unsubstituted lower alkyl)thio, cyano, acyl, thioacyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, nitro, N-sulfonamido, S-sulfonamido,  $R^{25}S(O)-$ ,  $R^{25}S(O)_2-$ ,  $-C(O)OR^{25}$ ,  $R^{25}C(O)O-$ , and  $-NR^{25}R^{26}$ , with  $R^{25}$  and  $R^{26}$  as defined above.

A "hydroxy" group refers to an -OH group.

An "alkoxy" group refers to both an -O-(unsubstituted alkyl) and an -O-(unsubstituted cycloalkyl) group.

WO 00/08202

17

PCT/US99/17845 ,

An "aryloxy" group refers to both an -O-aryl and an -O-heteroaryl group, as defined herein.

A "mercapto" group refers to an -SH group.

A "alkylthio" group refers to both an S-(unsubstituted alkyl) and an -S-(unsubstituted cycloalkyl) group.

A "arylthio" group refers to both an -S-aryl and an -S-heteroaryl group, as defined herein.

An "acyl" group refers to a -C(O)-R" group, where R" is selected from the group consisting of hydrogen, unsubstituted lower alkyl, trihalomethyl, unsubstituted cycloalkyl, aryl optionally substituted with one or more unsubstituted lower alkyl, trihalomethyl, unsubstituted lower alkoxy, halo and -NR<sup>25</sup>R<sup>26</sup> groups, heteroaryl (bonded through a ring carbon) optionally substituted with one or more unsubstituted lower alkyl, trihaloalkyl, unsubstituted lower alkoxy, halo and -NR<sup>25</sup>R<sup>26</sup> groups and heteroalicyclic (bonded through a ring carbon) optionally substituted with one or more unsubstituted lower alkyl, trihaloalkyl, unsubstituted lower alkoxy, halo and -NR<sup>25</sup>R<sup>26</sup> groups.

An "aldehyde" group refers to an acyl group in which R" is hydrogen.

A "thioacyl" group refers to a -C(S)-R" group, with R" as defined herein.

An "ester" group refers to a -C(O)O-R" group with R" as defined herein except that R" cannot be hydrogen.

An "acetyl" group refers to a -C(O)CH<sub>3</sub> group.

A "halo" group refers to fluorine, chlorine, bromine or iodine.

A "trihalomethyl" group refers to a -CX<sub>3</sub> group wherein X is a halo group as defined herein.

A "trihalomethanesulfonyl" group refers to a X<sub>3</sub>CS(=O)<sub>2</sub>- groups with X as defined above.

A "cyano" group refers to a -C≡N group.

A "methylenedioxy" group refers to a -OCH<sub>2</sub>O- group where



WO 00/08202

18

PCT/US99/17845 ,

the two oxygen atoms are bonded to adjacent carbon atoms.

An "ethylenedioxy" group refers to a  $-\text{OCH}_2\text{CH}_2\text{O}-$  where the two oxygen atoms are bonded to adjacent carbon atoms.

An "O-carbamyl" group refers to a  $-\text{OC}(\text{O})\text{NR}^{25}\text{R}^{26}$  group with  
5  $\text{R}^{25}$  and  $\text{R}^{26}$  as defined herein.

An "N-carbamyl" group refers to an  $\text{R}^{25}\text{OC}(\text{O})\text{NR}^{26}-$  group, with  $\text{R}^{25}$  and  $\text{R}^{26}$  as defined herein.

An "O-thiocarbamyl" group refers to a  $-\text{OC}(\text{S})\text{NR}^{25}\text{R}^{26}$  group with  $\text{R}^{25}$  and  $\text{R}^{26}$  as defined herein.

10 An "N-thiocarbamyl" group refers to a  $\text{R}^{25}\text{OC}(\text{S})\text{NR}^{26}-$  group, with  $\text{R}^{25}$  and  $\text{R}^{26}$  as defined herein.

An "amino" group refers to an  $-\text{NR}^{25}\text{R}^{26}$  group, wherein  $\text{R}^{25}$  and  $\text{R}^{26}$  are both hydrogen.


A "C-amido" group refers to a  $-\text{C}(\text{O})\text{NR}^{25}\text{R}^{26}$  group with  $\text{R}^{25}$   
15 and  $\text{R}^{26}$  as defined herein.

An "N-amido" group refers to a  $\text{R}^{25}\text{C}(\text{O})\text{NR}^{26}-$  group, with  $\text{R}^{25}$  and  $\text{R}^{26}$  as defined herein.

A "nitro" group refers to a  $-\text{NO}_2$  group.

#### B. Preferred Structural Features.

20 A presently preferred aspect of this invention is a compound in which Q is




Another presently preferred aspect of this invention is a compound in which Q is

WO 00/08202

19

PCT/US99/17845



and

R<sup>1</sup> is selected from the group consisting of hydrogen, lower alkyl and -C(O)OH;

- 5 R<sup>2</sup> is selected from the group consisting of hydrogen, lower alkyl, halo, -C(O)R<sup>19</sup>, -C(O)OR<sup>19</sup>, lower alkoxy, -NR<sup>21</sup>R<sup>22</sup>, -(CH<sub>2</sub>)<sub>n</sub>R<sup>20</sup> and -NHC(O)OR<sup>19</sup>;

R<sup>3</sup> is selected from the group consisting of hydrogen, hydroxy, halo, -NHC(O)O(lower alkyl), -NHSO<sub>2</sub>R<sup>19</sup>, -NHC(O)R<sup>24</sup>,

- 10 trihalomethyl, and aryl optionally substituted with one or more lower alkoxy groups;

R<sup>4</sup> is hydrogen;

R<sup>19</sup> is selected from the group consisting of hydrogen and lower alkyl;

- 15 n is 2 or 3;

R<sup>20</sup> is selected from the group consisting of hydroxy, -C(O)OH, morpholin-4-yl, piperidin-1-yl, piperazin-1-yl, pyrrolidin-1-yl and -NR<sup>21</sup>R<sup>22</sup>;

- 20 R<sup>21</sup> and R<sup>22</sup> are independently selected from the group consisting of hydrogen and lower alkyl; and,


R<sup>24</sup> is selected from the group consisting of hydrogen and lower alkyl.

It is likewise a presently preferred embodiment of this invention that Q is

WO 00/08202

20

PCT/US99/17845



R<sup>1</sup> is selected from the group consisting of hydrogen, lower alkyl and -C(O)OH;

R<sup>2</sup> is selected from the group consisting of hydrogen, lower  
5 alkyl, halo, -C(O)R<sup>19</sup>, -C(O)OR<sup>19</sup>, lower alkoxy, -NR<sup>21</sup>R<sup>22</sup>,  
- (CH<sub>2</sub>)<sub>n</sub>R<sup>20</sup> and -NHC(O)OR<sup>19</sup>;

R<sup>3</sup> is selected from the group consisting of hydrogen, hydroxy,  
halo, -NHC(O)O(lower alkyl), -NHSO<sub>2</sub>R<sup>19</sup>, -NHC(O)R<sup>24</sup>,  
trihalomethyl, and aryl optionally substituted with one or  
10 more lower alkoxy groups;

R<sup>4</sup> is hydrogen;

R<sup>19</sup> is selected from the group consisting of hydrogen and  
lower alkyl;

n is 2 or 3;

15 R<sup>20</sup> is selected from the group consisting of hydroxy,  
-C(O)OH, morpholin-4-yl, piperidin-1-yl, piperazin-1-yl,  
pyrrolidin-1-yl and -NR<sup>21</sup>R<sup>22</sup>;

R<sup>21</sup> and R<sup>22</sup> are independently selected from the group  
consisting of hydrogen and lower alkyl;

20 R<sup>24</sup> is selected from the group consisting of hydrogen and  
lower alkyl;

R<sup>5</sup> is selected from the group consisting of hydrogen, lower  
alkyl, halo, five-member ring heteroaryl and aryl optionally  
substituted with one or more groups selected from the group  
25 consisting of lower alkyl, halo, hydroxy, -NR<sup>21</sup>R<sup>22</sup> and lower  
alkoxy;


R<sup>6</sup> is selected from the group consisting of hydrogen, lower  
alkyl, 3 to 7-member cycloalkyl, lower alkoxy, halo, aryl

WO 00/08202

21

PCT/US99/17845

optionally substituted with one or more groups independently selected from the group consisting of lower alkyl, halo, hydroxy,  $-NR^{21}R^{22}$  and lower alkoxy and, combined with  $R^{18}$ , a group having the structure




5

wherein  $y$  and  $y'$  are either both hydrogen or both lower alkyl;

$R^7$  is  $-OR^{18}$  wherein  $R^{18}$  is selected from the group consisting of lower alkyl,  $-(CH_2)_nR^{20}$  and, combined with  $R^6$  or  $R^8$ , a group having the structure

10




$R^8$  is selected from the group consisting of hydrogen, lower alkyl, 3 to 6-member ring cycloalkyl, lower alkoxy, halo, aryl optionally substituted with one or more groups selected from the group consisting of lower alkyl, lower alkoxy, halo,  $-NR^{21}R^{22}$  and  $-NHC(O)(\text{lower alkyl})$ , five-member heteroaryl having from 1 to 3 heteroatoms in the ring and 6-member heteroaryl having from 1 to 3 heteroatoms in the ring and, combined with  $R^{18}$  a group having the structure

20

WO 00/08202


22

PCT/US99/17845



$R^9$  is selected from the group consisting of hydrogen, lower alkyl; hydroxy, lower alkoxy, halo and  $-NR^{21}R^{22}$ .

A further presently preferred aspect of this invention is a compound in which Q is



$R^1$  is selected from the group consisting of hydrogen, lower alkyl and  $-C(O)OH$ ;

$R^2$  is selected from the group consisting of hydrogen, lower alkyl, halo,  $-C(O)R^{19}$ ,  $-C(O)OR^{19}$ , lower alkoxy,  $-NR^{21}R^{22}$ ,  $-(CH_2)_nR^{20}$  and  $-NHC(O)OR^{19}$ ;

$R^3$  is selected from the group consisting of hydrogen, hydroxy, halo,  $-NHC(O)O(\text{lower alkyl})$ ,  $-NHSO_2R^{19}$ ,  $-NHC(O)R^{24}$ , trihalomethyl, and aryl optionally substituted with one or more lower alkoxy groups;

$R^4$  is hydrogen;

$R^{19}$  is selected from the group consisting of hydrogen and lower alkyl;

$n$  is 2 or 3;

$R^{20}$  is selected from the group consisting of hydroxy,  $-C(O)OH$ , morpholin-4-yl, piperidin-1-yl, piperazin-1-yl, pyrrolidin-1-yl and  $-NR^{21}R^{22}$ ;

WO 00/08202

23


PCT/US99/17845

$R^{21}$  and  $R^{22}$  are independently selected from the group consisting of hydrogen and lower alkyl;

$R^{24}$  is selected from the group consisting of hydrogen and lower alkyl;


- 5  $R^5$  is selected from the group consisting of hydrogen, lower alkyl, thien-2-yl, thien-3-yl and aryl optionally substituted with one or more lower alkoxy groups;

- $R^6$  is selected from the group consisting of hydrogen, lower alkyl, 5 or 6-member cycloalkyl, aryl optionally substituted  
10 with one or more lower alkoxy groups and, combined with  $R^{18}$ , a group having the structure



- wherein  $y$  and  $y'$  are either both hydrogen or both lower  
15 alkyl;

$R^7$  is  $-OR^{18}$ , wherein  $R^{18}$  is selected from the group consisting of lower alkyl,  $-(CH_2)_nR^{20}$  and, combined with  $R^6$  or  $R^8$ , a group having the structure



- 20  $n$  is 2 or 3;

$R^{20}$  is selected from the group consisting of hydroxy,  $-C(O)OH$ , morpholin-4-yl, piperidin-1-yl, piperazin-1-yl


WO 00/08202

24

PCT/US99/17845 ,


pyrrolidin-1-yl and  $-NR^{21}R^{22}$ ;

$R^8$  is selected from the group consisting of hydrogen, lower alkyl, 5 or 6-member ring cycloalkyl, lower alkoxy, aryl optionally substituted with a  $-NHC(O)$  (lower alkyl) group,  
 5 thien-2-yl, thien-3-yl, pyridin-2-yl, pyridin-3-yl and, combined with  $R^{18}$ , a group having the structure




$R^9$  is selected from the group consisting of hydrogen and lower  
 10 alkyl.

Still another presently preferred aspect of this invention is a compound in which Q is:



15

It is a presently preferred aspect of this compound that,  
 in a compound in which Q is



$R^1$  is selected from the group consisting of hydrogen, lower

WO 00/08202

25

PCT/US99/17845


alkyl and  $-(CH_2)_nR^{20}$ ;

n is 2 or 3;

$R^{20}$  is selected from the group consisting of hydroxy and  $-C(O)OH$ ;

- 5  $R^2$  is selected from the group consisting of hydrogen, halo, lower alkyl, lower alkoxy,  $-SO_2NR^{21}R^{22}$  and  $-C(O)OH$ ;
- $R^3$  is selected from the group consisting of hydrogen, halo, lower alkyl, lower alkoxy and aryl optionally substituted with one or more lower alkoxy groups; and,
- 10  $R^4$  is hydrogen.

Likewise, in a compound in which Q is



- it is a presently preferred aspect of this invention that
- 15  $R^1$  is selected from the group consisting of hydrogen, lower alkyl and  $-(CH_2)_nR^{20}$ ;
- n is 2 or 3;
- $R^{20}$  is selected from the group consisting of hydroxy and  $-C(O)OH$ ;
- 20  $R^2$  is selected from the group consisting of hydrogen, halo, lower alkyl, lower alkoxy,  $-SO_2NR^{21}R^{22}$  and  $-C(O)OH$ ;
- $R^3$  is selected from the group consisting of hydrogen, halo, lower alkyl, lower alkoxy and aryl optionally substituted with one or more lower alkoxy groups;
- 25  $R^4$  is hydrogen;
- $R^{10}$  is selected from the group consisting of lower alkyl and  $-C(O)OR^{19}$ ;
- $R^{11}$  is selected from the group consisting of hydrogen, lower alkyl and  $-C(O)OR^{19}$ ;



WO 00/08202

26

PCT/US99/17845

$R^{12}$  is  $-(CH_2)_nR^{20}$ ;


$R^{19}$  is selected from the group consisting of hydrogen and lower alkyl;

$n$  is 2 or 3; and,

5  $R^{20}$  is selected from the group consisting of hydroxy,  $-C(O)OH$ , morpholin-4-yl, piperidin-1-yl, piperazin-1-yl, pyrrolidin-1-yl and  $-NR^{21}R^{22}$ .

It is yet another presently preferred aspect of this invention that, in a compound in which Q is

10



$R^1$  is selected from the group consisting of hydrogen, lower alkyl and  $-(CH_2)_nR^{20}$ ;

15  $n$  is 2 or 3;

$R^{20}$  is selected from the group consisting of hydroxy and  $-C(O)OH$ ;

$R^2$  is selected from the group consisting of hydrogen, halo, lower alkyl, lower alkoxy,  $-SO_2NR^{21}R^{22}$  and  $-C(O)OH$ ;

20  $R^3$  is selected from the group consisting of hydrogen, halo, lower alkyl, lower alkoxy and aryl optionally substituted with one or more lower alkoxy groups;

$R^4$  is hydrogen;

$R^{10}$  is selected from the group consisting of lower alkyl and

25  $-C(O)O(\text{lower alkyl})$ ;

$R^{11}$  is selected from the group consisting of hydrogen, lower alkyl and  $-C(O)OR^{19}$ ;

$R^{12}$  is  $-(CH_2)_nR^{20}$ ;

$R^{19}$  is selected from the group consisting of hydrogen and

WO 00/08202

27

PCT/US99/17845 ,

lower alkyl;


n is 2; and,

R<sup>20</sup> is selected from the group consisting of -C(O)OH,

morpholin-1-yl, piperidin-1-yl, piperazin-1-yl,

5 pyrrolidin-1-yl and -N(lower alkyl)<sub>2</sub>.

It is also a presently preferred aspect of this invention that Q is




10 R<sup>1</sup> is selected from the group consisting of hydrogen and lower alkyl;

R<sup>2</sup> is selected from the group consisting of hydrogen, lower alkyl, halo, -C(O)OR<sup>19</sup>, -C(O)R<sup>19</sup>, -NR<sup>21</sup>R<sup>22</sup>, -SO<sub>2</sub>NR<sup>21</sup>R<sup>22</sup> and -N=C-R<sup>23</sup>;

15 R<sup>3</sup> is selected from the group consisting of hydrogen, halo, lower alkyl, lower alkoxy, morpholin-4-yl, piperidin-1-yl, piperazin-1-yl, pyrrolidin-1-yl and aryl optionally substituted with one or more lower alkoxy groups,; and, R<sup>4</sup> is hydrogen.

20 It is a presently preferred aspect of this invention that, in a compound in which Q is



a and b are both single bonds;

WO 00/08202

28

PCT/US99/17845

r is 2;

R<sup>1</sup> is selected from the group consisting of hydrogen and lower alkyl;

R<sup>2</sup> is selected from the group consisting of hydrogen, lower alkyl, halo, -C(O)OR<sup>19</sup>, -C(O)R<sup>19</sup>, -NR<sup>21</sup>R<sup>22</sup>, -SO<sub>2</sub>NR<sup>21</sup>R<sup>22</sup> and -N=C-R<sup>23</sup>;

R<sup>3</sup> is selected from the group consisting of hydrogen, halo, lower alkyl, lower alkoxy, morpholin-4-yl, piperidin-1-yl, piperazin-1-yl, pyrrolidin-1-yl and aryl optionally substituted with one or more lower alkoxy groups;

R<sup>4</sup> is hydrogen;

R<sup>13</sup> is a covalent bond through which Q is bonded to the rest of the molecule;


R<sup>14</sup> is selected from the group consisting of hydrogen, lower alkyl and -(CH<sub>2</sub>)<sub>n</sub>R<sup>20</sup>;

R<sup>15</sup> is hydrogen; and,

R<sup>16</sup> and R<sup>17</sup> are independently selected from the group consisting of hydrogen and lower alkyl.

It is also a presently preferred aspect of this invention that, in a compound having the substituents indicated in the paragraph immediately above, R<sup>20</sup> is -NR<sup>21</sup>R<sup>22</sup> wherein R<sup>21</sup> and R<sup>22</sup> are independently selected from the group consisting of hydrogen and lower alkyl.

It is yet another presently preferred aspect of this invention that, in a compound in which Q is



a and b are both double bonds;

WO 00/08202

29

PCT/US99/17845

$r$  is 1;

$R^1$  is selected from the group consisting of hydrogen and lower alkyl;

$R^2$  is selected from the group consisting of hydrogen, lower  
 5 alkyl, halo,  $-C(O)OR^{19}$ ,  $-C(O)R^{19}$ ,  $-NR^{21}R^{22}$ ,  $-SO_2NR^{21}R^{22}$  and  
 $-N=C-R^{23}$ ;

$R^3$  is selected from the group consisting of hydrogen, halo,  
 lower alkyl, lower alkoxy, morpholin-4-yl, piperidin-1-yl,  
 piperazin-1-yl, pyrrolidin-1-yl and aryl optionally  
 10 substituted with one or more lower alkoxy groups;

$R^4$  is hydrogen;


$R^{13}$  is a covalent bond through which  $Q$  is bonded to the rest of  
 the molecule;

$R^{14}$  is selected from the group consisting of hydrogen, lower  
 15 alkyl and  $-(CH_2)_nR^{20}$ ;

$R^{15}$  and  $R^{16}$  are hydrogen; and,

$R^{17}$  does not exist.

A compound in which  $Q$  is



20 a and b are both double bonds;

$r$  is 1;

$R^1$  is selected from the group consisting of hydrogen and lower  
 alkyl;

$R^2$  is selected from the group consisting of hydrogen, lower  
 25 alkyl, halo,  $-C(O)OR^{19}$ ,  $-C(O)R^{19}$ ,  $-NR^{21}R^{22}$ ,  $-SO_2NR^{21}R^{22}$  and  
 $-N=C-R^{23}$ ;

$R^3$  is selected from the group consisting of hydrogen, halo,  
 lower alkyl, lower alkoxy, morpholin-4-yl, piperidin-1-yl,

WO 00/08202

30

PCT/US99/17845

piperazin-1-yl, pyrrolidin-1-yl and aryl optionally substituted with one or more lower alkoxy groups;

R<sup>4</sup> is hydrogen;


R<sup>13</sup> is selected from the group consisting of hydrogen and lower alkyl;

R<sup>14</sup> is a covalent bond through which Q is bonded to the rest of the molecule;

R<sup>15</sup> and R<sup>16</sup> are hydrogen; and,

R<sup>17</sup> does not exist is another presently preferred aspect of this invention.

It is a presently preferred aspect of this invention that, in a compound where Q is



a and b are double bonds;

r is 1;

R<sup>1</sup> is selected from the group consisting of hydrogen and lower alkyl;

R<sup>2</sup> is selected from the group consisting of hydrogen, lower alkyl, halo, -C(O)OR<sup>19</sup>, -C(O)R<sup>19</sup>, -NR<sup>21</sup>R<sup>22</sup>, -SO<sub>2</sub>NR<sup>21</sup>R<sup>22</sup> and -N=C-R<sup>23</sup>;

R<sup>3</sup> is selected from the group consisting of hydrogen, halo, lower alkyl, lower alkoxy, morpholin-4-yl, piperidin-1-yl, piperazin-1-yl, pyrrolidin-1-yl and aryl optionally substituted with one or more lower alkoxy groups;

R<sup>4</sup> is hydrogen;

WO 00/08202

31

PCT/US99/17845

R<sup>13</sup> is selected from the group consisting of hydrogen and lower alkyl;

R<sup>14</sup> is selected from the group consisting of hydrogen, lower alkyl and -(CH<sub>2</sub>)<sub>n</sub>R<sup>20</sup>;

5 R<sup>15</sup> is a covalent bond through which Q is bonded to the rest of the molecule;

R<sup>16</sup> are hydrogen; and,

R<sup>17</sup> does not exist.

A compound selected from the group consisting of:

3-(3,5-Diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one  
5-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

N-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide

3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-hydroxy-1,3-dihydroindol-2-one

5-Acetyl-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid methyl ester

3-(3-Isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one

5-Chloro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one

3-(3-Cyclopentyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

3-(3-Cyclopentyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one

3-(3-Cyclohexyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

5-Bromo-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one

5-Chloro-3-(2,3-dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one

WO 00/08202

32

PCT/US99/17845

5-Chloro-3-(2,2-dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one

N-{3-[3-Cyclohexyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide

3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-methoxy-1,3-dihydroindol-2-one

N-[3-(4-Methoxy-3-thiophen-3-ylbenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide

3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-methyl-1,3-dihydroindol-2-one

5-Amino-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

5-Chloro-3-(4-methoxy-3,5-dimethylbenzylidene)-1,3-dihydroindol-2-one

3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-fluoro-1,3-dihydroindol-2-one

3-(2,2-Dimethylchroman-6-ylmethylene)-5-fluoro-1,3-dihydroindol-2-one

5-Chloro-3-[3,5-diisopropyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-1,3-dihydroindol-2-one

3-(3,5-Diisopropyl-4-methoxybenzylidene)-7-fluoro-1,3-dihydroindol-2-one

3-(4-Methoxy-3-thiophen-3-ylbenzylidene)-5-(2-morpholin-4-ylethyl)-1,3-dihydroindol-2-one

N-[3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide

3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-ethyl-1,3-dihydroindol-2-one

N-[2'-Methoxy-5'-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-biphenyl-3-yl]-acetamide

5-Fluoro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one

N-[3-(4-Methoxy-3-thiophen-2-ylbenzylidene)-2-oxo-2,3-dihydro-

WO 00/08202

33

PCT/US99/17845

1H-indol-6-yl]-acetamide  
6-Amino-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one  
N-[3-(2,2-Dimethylchroman-6-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide  
5-Bromo-3-(2,2-dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one  
3-(4-Methoxy-3-thiophen-3-ylbenzylidene)-1,3-dihydroindol-2-one  
5-Bromo-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one  
5-Fluoro-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one  
3-(3-Isopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one  
3-(4,5-Dimethoxy-2-thiophen-2-ylbenzylidene)-1,3-dihydroindol-2-one  
N-{3-[4-(2-Morpholin-4-ylethoxy)-3-thiophen-2-ylbenzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide  
3-(2,2-Dimethylchroman-6-ylmethylene)-4-methyl-1,3-dihydroindol-2-one  
3-(2,3-Dihydrobenzofuran-5-ylmethylene)-5-fluoro-1,3-dihydroindol-2-one  
3-(3-Cyclohexyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one  
5-Fluoro-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one  
3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one  
3-(3'-Ethoxy-6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one  
3-(3-Cyclopentyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one  
3-(3-Cyclopentyl-4-methoxybenzylidene)-4-methyl-1,3-



WO 00/08202

34

PCT/US99/17845

dihydroindol-2-one

3-(4,5,2'-Trimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one

N-{3-[4-(2-Morpholin-4-ylethoxy)-3-thiophen-3-ylbenzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide

5-Chloro-3-(3-cyclohexyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-carbamic acid tert-butyl ester

3-(3,5-Diisopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

5-Bromo-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

N-{3-[3-tert-Butyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide

3-(4-Methoxy-3,5-dimethylbenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one

5-Bromo-3-[3,5-diisopropyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-1,3-dihydroindol-2-one

3-(3'-Ethoxy-4,5-dimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one

5-Chloro-3-(4-methoxy-3-thiophen-2-ylbenzylidene)-1,3-dihydroindol-2-one

5-Chloro-3-(4-methoxy-3-pyridin-3-ylbenzylidene)-1,3-dihydroindol-2-one

5-Chloro-3-(4,5,3'-trimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one

3-(4,5-Dimethoxy-2-naphthalen-2-ylbenzylidene)-1,3-dihydroindol-2-one

N-[3-(3'-Acetylamino-6-methoxybiphenyl-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide

6-Methoxy-3-(4-methoxy-3-thiophen-3-ylbenzylidene)-1,3-dihydroindol-2-one

WO 00/08202

35

PCT/US99/17845

3-(6-Methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one  
3-(2,3-Dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one  
5-Chloro-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one  
3-(3-Cyclohexyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one  
3-(2,3-dihydrobenzofuran-5-ylmethylene)-4-methyl-1,3-dihydroindol-2-one  
3-(3-Isopropyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one  
3-(6-Methoxybiphenyl-3-ylmethylene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one  
3-(3-Cyclohexyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one  
3-(2,3-Dihydrobenzofuran-5-ylmethylene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one  
3-(3,5-Diisopropyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one  
5-Bromo-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one  
5-Bromo-3-(3-cyclopentyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one  
5-Chloro-3-(3-cyclopentyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one  
5-Chloro-3-(6-methoxybiphenyl-3-ylmethylene)-4-methyl-1,3-dihydroindol-2-one  
5-Chloro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-4-methyl-1,3-dihydroindol-2-one  
5-Chloro-3-(4-methoxy-3,5-dimethylbenzylidene)-4-methyl-1,3-dihydroindol-2-one  
3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-trifluoromethyl-1,3-dihydroindol-2-one  
6-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-

WO 00/08202

36

PCT/US99/17845

dihydroindol-2-one

3-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-5-yl]-propionic acid

3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-methoxy-1,3-dihydroindol-2-one

5-Butyl-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid

3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-(3-methoxyphenyl)-1,3-dihydroindol-2-one

7-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-5-methyl-1,3-dihydroindol-2-one

[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-5-yl]-carbamic acid tert-butyl ester

5-Chloro-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

5-Chloro-3-(3-cyclopentyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

3-(6-Methoxybiphenyl-3-ylmethylene)-4-methyl-1,3-dihydroindol-2-one

3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-4-methyl-1,3-dihydroindol-2-one

5-Bromo-3-(2,3-dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one

5-Chloro-3-(3-isopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

5-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

5-Chloro-3-(2,2-dimethylchroman-6-ylmethylene)-4-methyl-1,3-dihydroindol-2-one

3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid

WO 00/08202

37

PCT/US99/17845

3- (3,5-Diisopropyl-4-methoxybenzylidene)-5,6-dimethoxy-1,3-dihydroindol-2-one

N- [3- (3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-methanesulfonamide

N- [3- (3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-benzamide

3- (3,5-Diisopropyl-4-methoxybenzylidene)-6- (3-ethoxyphenyl)-1,3-dihydroindol-2-one

3- (3,5-Diisopropyl-4-methoxybenzylidene)-6-phenyl-1,3-dihydroindol-2-one

3- (3,5-Diisopropyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one

5-Fluoro-3- (4-methoxy-3,5-dimethylbenzylidene)-1,3-dihydroindol-2-one

3- (2,2-dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one

is also a preferred embodiment of this invention.

Likewise, a compound selected from the group consisting of:

5-methyl-3- (3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,

3- (3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-5-sulfonic acid amide,

3 (3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide,

3- (3-methyl-1H-indole-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide,

3- (3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid,

5-acetyl-3- (3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,

5-acetyl-3- (1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,

3- (1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-5-sulfonic acid amide,

5-amino-3- (1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,

WO 00/08202

38

PCT/US99/17845,

3-(1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid,  
6-chloro-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,  
3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,  
5-chloro-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,  
5-bromo-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,  
3-(1H-indol-2-ylmethylene)-4-methyl-1,3-dihydroindol-2-one,  
3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,  
5-chloro-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,  
5-bromo-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,  
4-methyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one, and  
3-(1H-indol-2-ylmethylene)-5[(1H-indol-2-ylmethylene)amino]-1,3-dihydroindol-2-one

is also a presently preferred embodiment of this invention.

Finally, with regard to chemical compounds of this invention, a presently preferred embodiment comprises compounds selected from the group consisting of:

3-[2-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
3-[2-(5-chloro-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
3-[2-(5-bromo-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
3-[2-(4-methyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
3-[2-(5-methyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
3-[2-(6-chloro-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
3-[2-(6-methoxy-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-

WO 00/08202

39

PCT/US99/17845

4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
N,N-dimethyl-3-[2-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-  
4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide,  
3-[3-(3-dimethylamino-propyl)-4,5,6,7-tetrahydro-1H-indol-2-  
ylmethylene]-1,3-dihydro-indol-2-one,  
3-[2-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-  
tetrahydro-1H-indol-3-yl]-propionamide,  
3-[3-(3-morpholin-4-yl-3-oxo-propyl)-4,5,6,7-tetrahydro-1H-  
indol-2-ylmethylene]-1,3-dihydro-indol-2-one,  
N-methyl-3-[2-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-  
4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide,  
N-(2-morpholin-4-yl-ethyl)-3-[2-(2-oxo-1,2-dihydro-indol-3-  
ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-  
propionamide, 3-[2-(2-oxo-1,2-dihydro-pyrrolo[2,3-b]pyridin-  
3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic  
acid,  
3-{2-[6-(3-methoxy-phenyl)-2-oxo-1,2-dihydro-indol-3-  
ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl}-propionic  
acid,  
3-{2-[6-(4-methoxy-phenyl)-2-oxo-1,2-dihydro-indol-3-  
ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl}-propionic  
acid,  
3-[2-(2-oxo-6-phenyl-1,2-dihydro-indol-3-ylidenemethyl)-  
4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
3-{2-[6-(2-methoxy-phenyl)-2-oxo-1,2-dihydro-indol-3-  
ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl}-propionic  
acid,  
3-[2-(5-isopropylsulfamoyl-2-oxo-1,2-dihydro-indol-3-  
ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic  
acid,  
3-[2-(6-morpholin-4-yl-2-oxo-1,2-dihydro-indol-3-  
ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic  
acid,

WO 00/08202

40

PCT/US99/17845

3-[2-(5-chloro-4-methyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,

3-[2-(5-bromo-4-methyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,




3-[2-(5-bromo-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-N-(2-morpholin-4-ylethyl)-propionamide,

3-[2-(5-chloro-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-N-(2-morpholin-4-ylethyl)-propionamide, and

3-[2-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-phenyl]-propionic acid.

## 2. SYNTHESIS/COMBINATORIAL LIBRARIES


An additional aspect of this invention is a combinatorial library of at least ten 3-methylidenyl-2-indolinone compounds that can be formed by reacting oxindoles of structure 5 with aldehydes of structure 6, 7 or 8:

567

WO 00/08202

41

PCT/US99/17845



8

wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{11}$ ,  $R^{12}$ ,  $R^{13}$ ,  $R^{14}$ ,  
 5  $R^{15}$ ,  $R^{16}$  and  $R^{17}$  have the meanings set forth above.

As used herein, a "combinatorial library" refers to all the compounds formed by the reaction of each compound of one dimension with a compound in each of the other dimensions in a multi-dimensional array of compounds. In the context of the  
 10 present invention, the array is two dimensional and one dimension represents all the oxindoles of the invention and the second dimension represents all the aldehydes of the invention. Each oxindole may be reacted with each and every aldehyde in order to form a substituted 3-methylidenyl-2-indolinone compound. All  
 15 substituted 3-methylidenyl-2-indolinone compounds formed in this way are within the scope of the present invention. Also within the scope of the present invention are smaller combinatorial libraries formed by the reaction of some of the oxindoles with all of the aldehydes, all of the oxindoles with some of the  
 20 aldehydes, or some of the oxindoles with some of the aldehydes.

The oxindole in the above combinatorial library is preferably selected from the group consisting of oxindole itself and substituted oxindoles such as, without limitation, 6-bromooxindole, 5-hydroxyoxindole, 5-methoxyoxindole, 6-methoxyoxindole, 5-phenylaminosulfonyloxindole, 4-[2-(2-isopropylphenoxy)ethyl]oxindole, 4-[2-(3-isopropylphenoxy)-ethyl]oxindole, 4-[2-(4-isopropylphenoxy)ethyl]oxindole, 5-fluorooxindole, 6-fluorooxindole, 7-fluorooxindole, 6-



WO 00/08202

42

PCT/US99/17845

trifluoromethyloxindole, 5-chlorooxindole, 6-chlorooxindole, 4-carboxyindole, 5-bromooxindole, 5-bromo-4-methyloxindole, 6-(N-acetamido)oxindole, 4-methyloxindole, 5-methyloxindole, 4-methyl-5-chlorooxindole, 5-ethyloxindole, 6-hydroxyoxindole, 5-acetyloxindole, 5-carboxyindole, 5-aminooxindole, 6-aminooxindole, 4-(2-N-morpholinoethyl)oxindole, 7-azaoxindole, oxindole-4-carabamic acid t-butyl ester, oxindole-6-carbamic acid t-butyl ester, 4-(2-carboxyethyl)oxindole, 5-n-butyloxindole, 5,6-dimethoxyoxindole, 6-(methanesulfonamido)-oxindole, 6-(benzamido)oxindole, 5-ethoxyoxindole, 6-phenyloxindole, 6-(2-methoxyphenyl)oxindole, 6-(3-methoxyphenyl)oxindole, 6-(4-methoxyphenyl)oxindole, 5-aminosulfonyloxindole, 5-isopropylaminosulfonyloxindole, 5-dimethylaminosulfonyloxindole, 5-(N-morpholinosulfonyl)oxindole, 4-(2-hydroxyethyl)oxindole, 6-(3-ethoxyphenyl)oxindole, 6-(morpholin-4-yl)oxindole, 5-(2-(N-morpholino)ethyl)oxindole, 5-(methanesulfonamido)oxindole, 5-methoxycarbonyloxindole and 5-carboxyethyloxindole.

The aldehyde in the above combinatorial library is preferably selected from the group consisting of 3-(1-Benzyl-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl)propionic acid, 3-(5-Formyl-1-methoxycarbonylmethyl-2,4-dimethyl-1H-pyrrol-3-yl)propionic acid, 3-(5-Formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)propionic acid, 3-[5-Formyl-1-(3-methoxybenzyl)-2,4-dimethyl-1H-pyrrol-3-yl]propionic acid methyl ester, 3-(1-Cyclohexylmethyl-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl)propionic acid methyl ester, 3-[1-(2,2-Dimethyl-propyl)-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl]propionic acid methyl ester, 1,3,5-Trimethyl-4-(3-morpholin-4-yl-3-oxo-propyl)1H-pyrrole-2-carbaldehyde, 3-(5-Formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-N-(2-morpholin-4-ylethyl)propionamide, 3-(5-Formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-N-phenylpropionamide, 1,3,5-trimethyl-4-(3-oxo-3-piperidin-1-ylpropyl)-1H-pyrrole-2-carbaldehyde, 1,3,5-Trimethyl-4-(3-oxo-3-pyrrolidin-1-ylpropyl)-1H-pyrrole-2-carbaldehyde, 3-(5-formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-N-(4-methoxyphenyl)propionamide, 3-(5-Formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-N-(4-

WO 00/08202

43

PCT/US99/17845 ,

methoxyphenyl)propionamide, N-(4-fluorophenyl)-3-(5-formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-propionamide, 3-(5-Formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-N-(4-trifluoromethylphenyl)-propionamide, 3-[5-Formyl-1-(3-methoxy-benzyl)-2,4-dimethyl-1H-pyrrol-3-yl]propionic acid, 3-(1-cyclohexylmethyl-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl)propionic acid, 3-[1-(3-fluorobenzyl)-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl]propionic acid methyl ester, 3-(1-Benzyl-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl)propionic acid, 3-[1-(4-fluorobenzyl)-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl]propionic acid methyl ester, 3-[1-(4-fluorobenzyl)-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl]propionic acid, 3-[1-(3-fluorobenzyl)-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl]propionic acid, 3,5-dimethyl-4-(3-morpholin-4-yl-propyl)-1H-pyrrole-2-carbaldehyde, 4-(3-dimethylaminopropyl)-3,5-dimethyl-1H-pyrrole-2-carbaldehyde, 5-formyl-2,4-dimethyl-1H-pyrrole-3-carboxylic acid, 3,5-dimethyl-4-(4-methylpiperazine-1-carbonyl)-1H-pyrrole-2-carbaldehyde, 5-formyl-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (2-dimethylaminoethyl)amide, 3-(2-formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)propionic acid, 3-(3-dimethylaminopropyl)-2-formyl-4,5,6,7-tetrahydro-1H-indole, indole-2-carbaldehyde and 3-methylindole-2-carbaldehyde.

Another aspect of this invention provides a method for the synthesis of a 3-methylidenyl-2-indolinone of formula 1 comprising reacting an oxindole of formula 5 with an aldehyde of formula 6, 7 or 8 in a solvent, preferably in the presence of a base.

Examples of the oxindoles of formula 5 which may be reacted with an aldehyde of formula 6, 7 or 8 to give the substituted 3-methylidenyl-2-indolinones of formula 1 are oxindole itself and substituted oxindoles such as, without limitation, 6-bromooxindole, 5-hydroxyoxindole, 5-methoxyoxindole, 6-methoxyoxindole, 5-phenylaminosulfonyloxindole, 4-[2-(2-isopropylphenoxy)ethyl]oxindole, 4-[2-(3-isopropylphenoxy)ethyl]oxindole, 4-[2-(4-isopropylphenoxy)ethyl]oxindole, 5-fluorooxindole, 6-fluorooxindole, 7-fluorooxindole, 6-

WO 00/08202

44

PCT/US99/17845

trifluoromethyloxindole, 5-chlorooxindole, 6-chlorooxindole, 4-carboxyindole, 5-bromooxindole, 5-bromo-4-methyloxindole, 6-(N-acetamido)oxindole, 4-methyloxindole, 5-methyloxindole, 4-methyl-5-chlorooxindole, 5-ethyloxindole, 6-hydroxyoxindole, 5-acetyloxindole, 5-carboxyindole, 5-aminooxindole, 6-aminooxindole, 4-(2-N-morpholinoethyl)oxindole, 7-azaoxindole, oxindole-4-carabamic acid t-butyl ester, oxindole-6-carbamic acid t-butyl ester, 4-(2-carboxyethyl)oxindole, 5-n-butyloxindole, 5,6-dimethoxyoxindole, 6-(methanesulfonamido)-oxindole, 6-(benzamido)oxindole, 5-ethoxyoxindole, 6-phenyloxindole, 6-(2-methoxyphenyl)oxindole, 6-(3-methoxyphenyl)oxindole, 6-(4-methoxyphenyl)oxindole, 5-aminosulfonyloxindole, 5-isopropylaminosulfonyloxindole, 5-dimethylaminosulfonyloxindole, 5-(N-morpholinosulfonyl)oxindole, 4-(2-hydroxyethyl)oxindole, 6-(3-ethoxyphenyl)oxindole, 6-(morpholin-4-yl)oxindole, 5-(2-(N-morpholino)ethyl)oxindole, 5-(methanesulfonamido)oxindole, 5-methoxycarbonyloxindole and 5-carboxyethyloxindole.

Examples of aldehydes of structure 6, 7 or 8 which may be reacted with oxindoles of structure 2 are, without limitation, 3-(1-benzyl-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl)propionic acid, 3-(5-formyl-1-methoxycarbonylmethyl-2,4-dimethyl-1H-pyrrol-3-yl)-propionic acid, 3-(5-formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-propionic acid, 3-[5-formyl-1-(3-methoxybenzyl)-2,4-dimethyl-1H-pyrrol-3-yl]propionic acid methyl ester, 3-(1-cyclohexylmethyl-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl)propionic acid methyl ester, 3-[1-(2,2-dimethylpropyl)-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl]-propionic acid methyl ester, 1,3,5-trimethyl-4-(3-morpholin-4-yl-3-oxopropyl)-1H-pyrrole-2-carbaldehyde, 3-(5-formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-N-(2-morpholin-4-ylethyl)propionamide, 3-(5-formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-N-phenylpropionamide, 1,3,5-trimethyl-4-(3-oxo-3-piperidin-1-ylpropyl)-1H-pyrrole-2-carbaldehyde, 1,3,5-trimethyl-4-(3-oxo-3-pyrrolidin-1-ylpropyl)-1H-pyrrole-2-carbaldehyde, 3-(5-formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-N-(4-methoxyphenyl)propionamide, 3-(5-formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-N-(4-methoxyphenyl)propionamide, N-(4-


WO 00/08202

45

PCT/US99/17845

fluorophenyl)-3-(5-formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-  
 propionamide, 3-(5-formyl-1,2,4-trimethyl-1H-pyrrol-3-yl)-N-(4-  
 trifluoromethylphenyl)propionamide, 3-[5-formyl-1-(3-methoxy-  
 benzyl)-2,4-dimethyl-1H-pyrrol-3-yl]propionic acid, 3-(1-  
 5 cyclohexylmethyl-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl)propionic  
 acid, 3-[1-(3-fluorobenzyl)-5-formyl-2,4-dimethyl-1H-pyrrol-3-  
 yl]propionic acid methyl ester, 3-(1-benzyl-5-formyl-2,4-  
 dimethyl-1H-pyrrol-3-yl)propionic acid, 3-[1-(4-fluorobenzyl)-5-  
 formyl-2,4-dimethyl-1H-pyrrol-3-yl]propionic acid methyl ester,  
 10 3-[1-(4-fluorobenzyl)-5-formyl-2,4-dimethyl-1H-pyrrol-3-yl]-  
 propionic acid, 3-[1-(3-fluorobenzyl)-5-formyl-2,4-dimethyl-1H-  
 pyrrol-3-yl]propionic acid, 3,5-dimethyl-4-(3-morpholin-4-yl-  
 propyl)-1H-pyrrole-2-carbaldehyde, 4-(3-dimethylaminopropyl)-3,5-  
 dimethyl-1H-pyrrole-2-carbaldehyde, 5-formyl-2,4-dimethyl-1H-  
 15 pyrrole-3-carboxylic acid, 3,5-dimethyl-4-(4-methylpiperazine-1-  
 carbonyl)-1H-pyrrole-2-carbaldehyde, 5-formyl-2,4-dimethyl-1H-  
 pyrrole-3-carboxylic acid (2-dimethylaminoethyl)amide, 3-(2-  
 formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)propionic acid, 3-(3-  
 dimethylaminopropyl)-2-formyl-4,5,6,7-tetrahydro-1H-indole,  
 20 indole-2-carbaldehyde and 3-methylindole-2-carbaldehyde.

The synthesis of an aminopropyl-4,5,6,7-tetrahydro-1H-indole, structure 9, below, which is an intermediate to some of the compounds of this invention, is another aspect of this



WO 00/08202

46

PCT/US99/17845

invention:

In Step 1, the base can be an inorganic or an organic base. Examples of inorganic and organic bases are presented elsewhere herein. Preferably the base is sodium acetate. The "solvent" may be any solvent in which base and the other reactants are sufficiently soluble for reaction to take place. Preferred solvents are polar protic solvents (defined elsewhere herein) such as water, methanol and ethanol. The presently preferred solvent is water. The reaction is carried out at temperatures of from about 60° C to about 180° C. Preferably the temperature is between about 60° C and 180° C, more preferably between 80° C and 150° C, most preferably between 100 and 120° C. The reaction is allowed to run for from 1 to 30 hours, preferably from 2 to 20 hours, most preferably from 4 to 15 hours.

In step 2, a carboxylic acid group is converted to an amide group. Procedures for accomplishing this conversion are well-known in the art. The reaction of an amine with the carboxylic acid in the presence of 1,1'-carbonyldiimidazole is the presently preferred method. Any aprotic solvent may be used; in a presently preferred embodiment the solvent is a non-polar aprotic solvent, in particular dichloromethane. The reaction may be carried out at room temperature or at an elevated temperature up to about 100° C; a presently preferred embodiment is to run the reaction at about room temperature.

Step 3 is the simultaneous reduction of the keto and the amido group using a reducing agent such as, for example and without limitation, lithium aluminum hydride. The reaction is carried out in an aprotic solvent (defined elsewhere herein).

A presently preferred solvent is tetrahydrofuran. The reaction is carried out at temperatures from about room temperature to about 80° C, preferably from about 50° C to

WO 00/08202

47

PCT/US99/17845

about 70° C, most preferably, the reaction is carried out in tetrahydrofuran at reflux.

Step 4 is the well-known formylation of the aromatic pyrrole ring using phosphorus oxychloride and N,N-dimethylformamide.

Step 5, the condensation of an aldehyde with an oxindole to form a 3-methylidenyl-2-indolinone of this invention is carried out in a solvent which may contain a base. The base may be an organic or an inorganic base. If an organic base is used, preferably it is a nitrogen base. Examples of organic nitrogen bases include, but are not limited to, diisopropylamine, trimethylamine, triethylamine, aniline, pyridine, 1,8-diazabicyclo-[5.4.1]undec-7-ene, pyrrolidine and piperidine.

Examples of inorganic bases are, without limitation, ammonia, alkali metal or alkaline earth hydroxides, phosphates, carbonates, bicarbonates, acetates, bisulfates and amides. The alkali metals include, lithium, sodium and potassium while the alkaline earths include calcium, magnesium and barium.

In a presently preferred embodiment of this invention, when the solvent is a protic solvent, such as water or alcohol, the base is an alkali metal or an alkaline earth inorganic base, preferably, a alkali metal or an alkaline earth hydroxide.

It will be clear to those skilled in the art, based both on known general principles of organic synthesis and on the disclosures herein which base would be most appropriate for the reaction contemplated.

The solvent in which the reaction is carried out may be a protic or an aprotic solvent, preferably it is a protic solvent. A "protic solvent" is a solvent which has hydrogen atom(s) covalently bonded to oxygen or nitrogen atoms which renders the hydrogen atoms appreciably acidic and thus capable of being "shared" with a solute through hydrogen bonding.

WO 00/08202

48

PCT/US99/17845

Examples of protic solvents include, without limitation, water and alcohols.

An "aprotic solvent" may be polar or non-polar but, in either case, does not contain acidic hydrogens and therefore is not capable of hydrogen bonding with solutes. Examples, without limitation, of non-polar aprotic solvents, are pentane, hexane, benzene, toluene, methylene chloride and carbon tetrachloride.

Examples of polar aprotic solvents are chloroform, tetrahydrofuran, dimethylsulfoxide and dimethylformamide.

In a presently preferred embodiment of this invention, the solvent is a protic solvent, preferably water or an alcohol such as ethanol.

The reaction is carried out at temperatures greater than room temperature. The temperature is generally from about 30° C to about 150° C, preferably about 80° C to about 100° C, most preferable about 75° C to about 85° C, which is about the boiling point of ethanol.

By "about" is meant that a temperature range described herein is preferably within 10 degrees Celcius of the indicated temperature, more preferably within 5 degrees Celcius of the indicated temperature and, most preferably, within 2 degrees Celcius of the indicated temperature. Thus, for example, by "about 75° C" is meant 75° C  $\pm$  10° C, preferably 75° C  $\pm$  5° C and most preferably, 75° C  $\pm$  2° C.










Some representative compounds of this invention are shown in Table 1. The compounds shown are examples only and are not to be construed as limiting the scope of this invention in any manner whatsoever.

WO 00/08202

49

PCT/US99/17845

TABLE 1











Compound	Structures	Names
1		3-(3,5-Diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
2		5-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
3		N-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide
4		3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-hydroxy-1,3-dihydroindol-2-one
5		5-Acetyl-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
6		3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid methyl ester
7		3-(3-Isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
8		3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one
9		5-Chloro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one



WO 00/08202

50











PCT/US99/17845

10		3-(3-Cyclopentyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
11		3-(3-Cyclopentyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one
12		3-(3-Cyclohexyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
13		5-Bromo-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one
14		5-Chloro-3-(2,3-dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one
15		5-Chloro-3-(2,2-dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one
16		N-[3-[3-Cyclohexyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide
17		3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-methoxy-1,3-dihydroindol-2-one
18		N-[3-(4-Methoxy-3-thiophen-3-ylbenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide
19		3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-methyl-1,3-dihydroindol-2-one

WO 00/08202

51

PCT/US99/17845

20		5-Amino-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
21		5-Chloro-3-(4-methoxy-3,5-dimethylbenzylidene)-1,3-dihydroindol-2-one
22		3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-fluoro-1,3-dihydroindol-2-one
23		3-(2,2-Dimethylchroman-6-ylmethylene)-5-fluoro-1,3-dihydroindol-2-one
24		5-Chloro-3-[3,5-diisopropyl-4-(2-morpholin-4-ylethoxy)benzylidene]-1,3-dihydroindol-2-one
25		3-(3,5-Diisopropyl-4-methoxybenzylidene)-7-fluoro-1,3-dihydroindol-2-one
26		3-(4-Methoxy-3-thiophen-3-ylbenzylidene)-5-(2-morpholin-4-ylethyl)-1,3-dihydroindol-2-one
27		N-[3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide
28		3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-ethyl-1,3-dihydroindol-2-one
29		N-[2'-Methoxy-5'-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-biphenyl-3-yl]-acetamide

WO 00/08202

52












PCT/US99/17845

30		5-Fluoro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one
31		N-[3-(4-Methoxy-3-thiophen-2-ylbenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide
32		6-Amino-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
33		N-[3-(2,2-Dimethylchroman-6-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide
34		5-Bromo-3-(2,2-dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one
35		3-(4-Methoxy-3-thiophen-3-ylbenzylidene)-1,3-dihydroindol-2-one
36		5-Bromo-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one
37		5-Fluoro-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one
38		3-(3-Isopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one
39		3-(4,5-Dimethoxy-2-thiophen-2-ylbenzylidene)-1,3-dihydroindol-2-one

WO 00/08202

53











PCT/US99/17845

40		N-[3-[4-(2-Morpholin-4-ylethoxy)-3-thiophen-2-ylbenzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide
41		3-(2,2-Dimethylchroman-6-ylmethylene)-4-methyl-1,3-dihydroindol-2-one
42		3-(2,3-Dihydrobenzofuran-5-ylmethylene)-5-fluoro-1,3-dihydroindol-2-one
43		3-(3-Cyclohexyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one
44		5-Fluoro-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
45		3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one
46		3-(3'-Ethoxy-6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one
47		3-(3-Cyclopentyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one
48		3-(3-Cyclopentyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one
49		3-(4,5,2'-Trimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one
50		N-[3-[4-(2-Morpholin-4-ylethoxy)-3-thiophen-3-ylbenzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide

WO 00/08202

54

PCT/US99/17845

51		5-Chloro-3-(3-cyclohexyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
52		[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-carbamic acid tert-butyl ester
53		3-(3,5-Diisopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one
54		5-Bromo-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
55		N-[3-[3-tert-Butyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide
56		3-(4-Methoxy-3,5-dimethylbenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one
57		5-Bromo-3-[3,5-diisopropyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-1,3-dihydroindol-2-one
58		3-(3'-Ethoxy-4,5-dimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one
59		5-Chloro-3-(4-methoxy-3-thiophen-2-ylbenzylidene)-1,3-dihydroindol-2-one
60		5-Chloro-3-(4-methoxy-3-pyridin-3-ylbenzylidene)-1,3-dihydroindol-2-one

WO 00/08202

55











PCT/US99/17845

61		5-Chloro-3-(4,5,3'-trimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one
62		3-(4,5-Dimethoxy-2-naphthalen-2-ylbenzylidene)-1,3-dihydroindol-2-one
63		N-[3-(3'-Acetylamino-6-methoxybiphenyl-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide
64		6-Methoxy-3-(4-methoxy-3-thiophen-3-ylbenzylidene)-1,3-dihydroindol-2-one
65		3-(6-Methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one
66		3-(2,3-Dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one
67		5-Chloro-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one
68		3-(3-Cyclohexyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one
69		3-(2,3-dihydrobenzofuran-5-ylmethylene)-4-methyl-1,3-dihydroindol-2-one
70		3-(3-Isopropyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one

WO 00/08202

56

PCT/US99/17845

71		3-(6-Methoxybiphenyl-3-ylmethylene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one
72		3-(3-Cyclohexyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one
73		3-(2,3-Dihydrobenzofuran-5-ylmethylene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one
74		3-(3,5-Diisopropyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one
75		5-Bromo-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
76		5-Bromo-3-(3-cyclopentyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
77		5-Chloro-3-(3-cyclopentyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one
78		5-Chloro-3-(6-methoxybiphenyl-3-ylmethylene)-4-methyl-1,3-dihydroindol-2-one
79		5-Chloro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-4-methyl-1,3-dihydroindol-2-one
80		5-Chloro-3-(4-methoxy-3,5-dimethylbenzylidene)-4-methyl-1,3-dihydroindol-2-one

WO 00/08202

57

PCT/US99/17845

81		3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-trifluoromethyl-1,3-dihydroindol-2-one
82		6-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
83		3-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-5-yl]-propionic acid
84		3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-methoxy-1,3-dihydroindol-2-one
85		5-Butyl-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
86		3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid
87		3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-(3-methoxyphenyl)-1,3-dihydroindol-2-one
88		7-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-5-methyl-1,3-dihydroindol-2-one
89		[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-5-yl]-carbamic acid tert-butyl ester
90		5-Chloro-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one



WO 00/08202

58










PCT/US99/17845

91		5-Chloro-3-(3-cyclopentyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one
92		3-(6-Methoxybiphenyl-3-ylmethylene)-4-methyl-1,3-dihydroindol-2-one
93		3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-4-methyl-1,3-dihydroindol-2-one
94		5-Bromo-3-(2,3-dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one
95		5-Chloro-3-(3-isopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one
96		5-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one
97		5-Chloro-3-(2,2-dimethylchroman-6-ylmethylene)-4-methyl-1,3-dihydroindol-2-one
98		3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid
99		3-(3,5-Diisopropyl-4-methoxybenzylidene)-5,6-dimethoxy-1,3-dihydroindol-2-one
100		N-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-methanesulfonamide
101		N-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-benzamide

WO 00/08202

59

PCT/US99/17845

102		3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-(3-ethoxyphenyl)-1,3-dihydroindol-2-one
103		3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-phenyl-1,3-dihydroindol-2-one
104		3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one
105		5-Fluoro-3-(4-methoxy-3,5-dimethylbenzylidene)-1,3-dihydroindol-2-one
106		3-(2,2-dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one
107		3-{2-[6-(4-Fluoro-phenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-5-methyl-1H-pyrrol-3-yl}-propionic acid
108		4-(2-Carboxy-ethyl)-5-[6-(4-fluoro-phenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester
109		3-{2-[6-(2-Methoxy-phenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-5-methyl-1H-pyrrol-3-yl}-propionic acid
110		4-(2-Carboxy-ethyl)-5-[6-(2-methoxy-phenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester

WO 00/08202

60









PCT/US99/17845

111		3-[2-(5-Chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-5-methyl-1H-pyrrol-3-yl]-propionic acid
112		4-(2-Carboxy-ethyl)-5-(5-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester
113		4-(2-Carboxy-ethyl)-2-methyl-5-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid ethyl ester
114		5-(5-Bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-(2-carboxyethyl)-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester
115		3-[5-Methyl-2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrol-3-yl]-propionic acid
116		3-[2-(5-Bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-5-methyl-1H-pyrrol-3-yl]-propionic acid
117		4-(2-Carboxy-ethyl)-3-methyl-5-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester
118		2-Methyl-5-(5-methylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-(3-morpholin-4-yl-propyl)-1H-pyrrole-3-carboxylic acid ethyl ester
119		2-Methyl-4-[3-(4-methyl-piperazin-1-yl)-propyl]-5-(5-methylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid ethyl ester

WO 00/08202

61

PCT/US99/17845

120		4-(3-Dimethylamino-propyl)-5-(5-methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester
121		5-(5-Methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-2-methyl-4-(3-morpholin-4-yl-propyl)-1H-pyrrole-3-carboxylic acid ethyl ester
122		5-(5-Methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-2-methyl-4-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-pyrrole-3-carboxylic acid ethyl ester
123		2-Methyl-4-(3-morpholin-4-yl-propyl)-5-(2-oxo-5-sulfamoyl-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid ethyl ester
124		2-Methyl-4-[3-(4-methyl-piperazin-1-yl)-propyl]-5-(2-oxo-5-sulfamoyl-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid ethyl ester
125		3-[4-Ethoxycarbonyl-5-methyl-3-(3-morpholin-4-yl-propyl)-1H-pyrrol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid
126		2-Methyl-5-(5-methylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-(3-pyrrolidin-1-yl-propyl)-1H-pyrrole-3-carboxylic acid ethyl ester
127		5-(5-Methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-2-methyl-4-(3-pyrrolidin-1-yl-propyl)-1H-pyrrole-3-carboxylic acid ethyl ester

WO 00/08202

62










PCT/US99/17845

128		3-[4-Ethoxycarbonyl-5-methyl-3-(3-pyrrolidin-1-yl-propyl)-1H-pyrrol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid
129		3-[4-Ethoxycarbonyl-5-methyl-3-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-pyrrol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid
130		2-Methyl-5-(2-oxo-5-sulfamoyl-1,2-dihydroindol-3-ylidenemethyl)-4-(3-pyrrolidin-1-yl-propyl)-1H-pyrrole-3-carboxylic acid ethyl ester
131		5-[4-(2-Hydroxy-ethyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-2-methyl-4-(3-morpholin-4-yl-propyl)-1H-pyrrole-3-carboxylic acid ethyl ester
132		5-[4-(2-Hydroxy-ethyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-2-methyl-4-(3-pyrrolidin-1-yl-propyl)-1H-pyrrole-3-carboxylic acid ethyl ester
133		4-(3-Dimethylamino-propyl)-2-methyl-5-(2-oxo-5-sulfamoyl-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid ethyl ester
134		3-[3-(3-Dimethylamino-propyl)-4-ethoxycarbonyl-5-methyl-1H-pyrrol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid
135		4-(3-Dimethylamino-propyl)-5-[4-(2-hydroxy-ethyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester

WO 00/08202

63

PCT/US99/17845

136		5-[4-(2-Hydroxy-ethyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-2-methyl-4-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-pyrrole-3-carboxylic acid ethyl ester
137		4-(3-Dimethylamino-propyl)-2-methyl-5-(5-methylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid ethyl ester
138		4-(3-Dimethylamino-propyl)-2-methyl-5-(5-methylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid
139		4-(2-Carboxy-ethyl)-3-methyl-5-(4-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester
140		4-(2-Carboxy-ethyl)-5-(5-methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester
141		4-(2-Carboxy-ethyl)-5-(6-methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester
142		5-(6-Bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-(2-carboxy ethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester
143		5-(5-Bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-(2-carboxy ethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester
144		4-(2-Carboxy-ethyl)-3-methyl-5-(2-oxo-6-phenyl-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester

WO 00/08202

64

PCT/US99/17845

145		4-(2-Carboxy-ethyl)-3-methyl-5-(2-oxo-5-sulfamoyl-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester
146		4-(2-Carboxy-ethyl)-3-methyl-5-(5-methylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester
147		4-(2-Carboxy-ethyl)-5-(5-dimethylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester
148		4-(2-Carboxy-ethyl)-5-(5-isopropylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester
149		3-[3-(3-Dimethylamino-propyl)-1H-indol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide
150		3-[3-(3-Dimethylamino-propyl)-1H-indol-2-ylmethylene]-4-(2-hydroxy-ethyl)-1,3-dihydroindol-2-one
151		3-[3-(3-Dimethylamino-propyl)-1H-indol-2-ylmethylene]-5-methoxy-1,3-dihydroindol-2-one
152		5-methyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one
153		3-(3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-5-sulfonic acid amide
154		3-(3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide

WO 00/08202

65

PCT/US99/17845












155		3-(3-methyl-1H-indole-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide
156		3-(3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid
157		5-acetyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one
158		5-acetyl-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one
159		3-(1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-5-sulfonic acid amide
160		5-amino-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one
161		3-(1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid
162		6-chloro-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one
163		3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one
164		5-chloro-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one



WO 00/08202

66

PCT/US99/17845

165		5-bromo-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one
166		3-(1H-indol-2-ylmethylene)-4-methyl-1,3-dihydroindol-2-one
167		3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one
168		5-chloro-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one
169		5-bromo-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one
170		4-methyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one
171		3-(1H-indol-2-ylmethylene)-5-[(1H-indol-2-ylmethylene)-amino]-1,3-dihydroindol-2-one
172		3-(1H-indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide
173		3-(1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide
174		3-(1H-indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid
175		3-(3-Methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide

WO 00/08202

67











PCT/US99/17845 ,

176		3-(1 <i>H</i> -Indol-5-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid methylamide
177		3-(1 <i>H</i> -Indol-5-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-carboxylic acid
178		3-(1 <i>H</i> -Indol-3-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid methylamide
179		5-Amino-3-(1 <i>H</i> -indol-5-ylmethylene)-1,3-dihydroindol-2-one
180		5-Amino-3-(3-methyl-1 <i>H</i> -indol-2-ylmethylene)-1,3-dihydroindol-2-one
181		3-(2-Methyl-1 <i>H</i> -indol-3-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid methylamide
182		3-(2-Methyl-1 <i>H</i> -indol-3-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-carboxylic acid
183		3-(1 <i>H</i> -Indol-5-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid dimethylamide
184		3-(2-Methyl-1 <i>H</i> -indol-3-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid dimethylamide
185		3-(1 <i>H</i> -Indol-5-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid amide
186		3-(1 <i>H</i> -Indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid dimethylamide

WO 00/08202

68

PCT/US99/17845

187		5-Amino-3-(2-methyl-1H-indol-3-ylmethylene)-1,3-dihydroindol-2-one
188		3-(1H-Indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide
189		3-(2-Methyl-1H-indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide
190		3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
191		3-[2-(5-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
192		3-[2-(5-bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
193		3-[2-(4-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
194		3-[2-(5-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
195		3-[2-(6-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
196		3-[2-(6-methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid

WO 00/08202

69

PCT/US99/17845

197		N,N-dimethyl-3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide
198		3-[3-(3-dimethylamino-propyl)-4,5,6,7-tetrahydro-1H-indol-2-ylmethylene]-1,3-dihydroindol-2-one
199		3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide
200		3-[3-(3-morpholin-4-yl-3-oxo-propyl)-4,5,6,7-tetrahydro-1H-indol-2-ylmethylene]-1,3-dihydroindol-2-one
201		N-methyl-3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide
202		N-(2-morpholin-4-yl-ethyl)-3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide
203		3-[2-(2-oxo-1,2-dihydro-pyrrolo[2,3-b]pyridin-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
204		3-[2-[6-(3-methoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
205		3-[2-[6-(4-methoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
206		3-[2-(2-oxo-6-phenyl-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid

WO 00/08202

70









PCT/US99/17845

207		3-[2-[6-(2-methoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
208		3-[2-(5-isopropylaminosulfonyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
209		3-[2-(6-morpholin-4-yl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
210		3-[2-(5-chloro-4-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
211		3-[2-(5-bromo-4-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid
212		3-[2-(5-bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-N-(2-morpholin-4-yl-ethyl)-propionamide
213		3-[2-(5-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-N-(2-morpholin-4-yl-ethyl)-propionamide
214		3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-phenyl]-propionic acid
215		3-[4-Methyl-2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrol-3-yl]-propionic acid
216		3-[2-(5-Chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid

WO 00/08202

71

PCT/US99/17845

217		3-[2-(6-Methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid
218		3-[2-(4-Methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid
219		3-[2-(6-Chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid
220		3-[2-(5-Bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid
221		3-[2-(5-Methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid
222		3-[2-(5-Methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid
223		3-[2-[6-(3-Methoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-4-methyl-1H-pyrrol-3-yl]-propionic acid
224		3-[2-[6-(3-Ethoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-4-methyl-1H-pyrrol-3-yl]-propionic acid

WO 00/08202

72

PCT/US99/17845 ,

### 3. BIOCHEMISTRY/PHARMACOTHERAPY

Another aspect of this invention relates to a method for the modulation of the catalytic activity of a PK by contacting a PK  
5 with a compound of this invention or a physiologically acceptable salt or prodrug thereof.

As used herein, "PK" refers to receptor protein tyrosine kinase (RTKs), non-receptor or "cellular" tyrosine kinase (CTKs) and serine-threonine kinases (STKs).

10 The term "method" refers to manners, means, techniques and procedures for accomplishing a given task including, but not limited to, those manners, means, techniques and procedures either known to, or readily developed from known manners, means, techniques and procedures by, practitioners of the chemical,  
15 pharmaceutical, biological, biochemical and medical arts.

As used herein, the term "modulation" or "modulating" refers to the alteration of the catalytic activity of RTKs, CTKs and STKs. In particular, modulating refers to the activation of the catalytic activity of RTKs, CTKs and STKs, preferably the  
20 activation or inhibition of the catalytic activity of RTKs, CTKs and STKs, depending on the concentration of the compound or salt to which the RTK, CTK or STK is exposed or, more preferably, the inhibition of the catalytic activity of RTKs, CTKs and STKs.

The term "catalytic activity" as used herein refers to the  
25 rate of phosphorylation of tyrosine under the influence, direct or indirect, of RTKs and/or CTKs or the phosphorylation of serine and threonine under the influence, direct or indirect, of STKs.

The term "contacting" as used herein refers to bringing a compound of this invention and a target PK together in such a  
30 manner that the compound can affect the catalytic activity of the PK, either directly, i.e., by interacting with the kinase itself, or indirectly, i.e., by interacting with another molecule on which the catalytic activity of the kinase is dependent. Such "contacting" can be accomplished "in vitro," i.e., in a test

WO 00/08202

73

PCT/US99/17845

tube, a petri dish or the like. In a test tube, contacting may involve only a compound and a PK of interest or it may involve whole cells. Cells may also be maintained or grown in cell culture dishes and contacted with a compound in that environment.

5 In this context, the ability of a particular compound to affect a PK related disorder, i.e., the  $IC_{50}$  of the compound, defined below, can be determined before use of the compounds in vivo with more complex living organisms is attempted. For cells outside the organism, multiple methods exist, and are well-known to those  
10 skilled in the art, to get the PKs in contact with the compounds including, but not limited to, direct cell microinjection and numerous transmembrane carrier techniques.

A further aspect of this invention is that the modulation of the catalytic activity of PKs using a compound of this  
15 invention may be carried out in vitro or in vivo.

"In vitro" refers to procedures performed in an artificial environment such as, e.g., without limitation, in a test tube or culture medium.

As used herein, "in vivo" refers to procedures performed  
20 within a living organism such as, without limitation, a mouse, rat, rabbit or human being.

A still further aspect of this invention is that the protein kinase whose catalytic activity is being modulated by a compound of this invention is selected from the group  
25 consisting of receptor protein tyrosine kinases, a cellular (or non-receptor) tyrosine kinases and serine-threonine kinases.

It is an aspect of this invention that the receptor protein kinase whose catalytic activity is modulated by a  
30 compound of this invention is selected from the group consisting of EGF, HER2, HER3, HER4, IR, IGF-1R, IRR, PDGFR $\alpha$ , PDGFR $\beta$ , CSF1R, C-Kit, C-fms, Flk-1R, Flk4, KDR/Flk-1, Flt-1, FGFR-1R, FGFR-2R, FGFR-3R and FGFR-4R.



WO 00/08202

74

PCT/US99/17845 ,

In addition, it is an aspect of this invention that the cellular tyrosine kinase whose catalytic activity is modulated by a compound of this invention is selected from the group consisting of Src, Frk, Btk, Csk, Abl, ZAP70, 5 Fes/Fps, Fak, Jak, Ack, Yes, Fyn, Lyn, Lck, Blk, Hck, Fgr and Yrk.

Another aspect of this invention is that the serine-threonine protein kinase whose catalytic activity is modulated by a compound of this invention is selected from the group 10 consisting of CDK2 and Raf.

A pharmaceutical composition of a compound of this invention with a pharmaceutically acceptable carrier or excipient is yet another aspect of this invention. Such pharmaceutical composition may contain both carriers and 15 excipients as well as other components generally known to those skilled in the formulation of pharmaceutical compositions.

A method for treating or preventing a protein kinase related disorder in an organism comprising administering a 20 therapeutically effective amount of a compound, salt or prodrug of this invention to an organism in need of such treatment is another aspect of this invention.

As used herein, "PK related disorder," "PK driven disorder," and "abnormal PK activity" all refer to a condition 25 characterized by inappropriate, i.e., under or, more commonly, over, PK catalytic activity, where the particular PK can be an RTK, a CTK or an STK. Inappropriate catalytic activity can arise as the result of either: (1) PK expression in cells which normally do not express PKs, (2) increased PK expression 30 leading to unwanted cell proliferation, differentiation and/or growth, or, (3) decreased PK expression leading to unwanted reductions in cell proliferation, differentiation and/or growth. Over-activity of a PK refers to either amplification of the gene encoding a particular PK or production of a level

WO 00/08202

75

PCT/US99/17845 ,

of PK activity which can correlate with a cell proliferation, differentiation and/or growth disorder (that is, as the level of the PK increases, the severity of one or more of the symptoms of the cellular disorder increases). Under-activity  
5 is, of course, the converse, wherein the severity of one or more symptoms of a cellular disorder increase as the level of the PK activity decreases.

As used herein, the terms "prevent", "preventing" and "prevention" refer to a method for barring an organism from  
10 acquiring a PK related disorder in the first place.

As used herein, the terms "treat", "treating" and "treatment" refer to a method of alleviating or abrogating a PK mediated cellular disorder and/or its attendant symptoms. With regard particularly to cancer, these terms simply mean  
15 that the life expectancy of an individual affected with a cancer will be increased or that one or more of the symptoms of the disease will be reduced.

The term "organism" refers to any living entity comprised of at least one cell. A living organism can be as simple as,  
20 for example, a single eukariotic cell or as complex as a mammal, including a human being.

The term "therapeutically effective amount" as used herein refers to that amount of the compound being administered which will relieve to some extent one or more of  
25 the symptoms of the disorder being treated. In reference to the treatment of cancer, a therapeutically effective amount refers to that amount which has the effect of (1) reducing the size of the tumor, (2) inhibiting (that is, slowing to some extent, preferably stopping) tumor metastasis, (3) inhibiting  
30 to some extent (that is, slowing to some extent, preferably stopping) tumor growth, and/or, (4) relieving to some extent (or, preferably, eliminating) one or more symptoms associated with the cancer.

WO 00/08202

76

PCT/US99/17845 ,

It is an aspect of this invention that the above-referenced protein kinase related disorder is selected from the group consisting of a receptor protein tyrosine kinase related disorder, a cellular tyrosine kinase disorder and a serine-threonine kinase related disorder.

In yet another aspect of this invention, the above referenced protein kinase related disorder is selected from the group consisting of an EGFR related disorder, a PDGFR related disorder, an IGFR related disorder and a flk related disorder.

The above referenced protein kinase related disorder is a cancer selected from the group consisting of squamous cell carcinoma, sarcomas such as Kaposi's sarcoma, astrocytoma, glioblastoma, lung cancer, bladder cancer, colorectal cancer, gastrointestinal cancer, head and neck cancer, melanoma, ovarian cancer, prostate cancer, breast cancer, small-cell lung cancer and glioma in a further aspect of this invention.

The above referenced protein kinase related disorder is selected from the group consisting of diabetes, an autoimmune disorder, a hyperproliferation disorder, von Hippel-Lindau disease, restenosis, fibrosis, psoriasis, osteoarthritis, rheumatoid arthritis, an inflammatory disorder and angiogenesis in yet another aspect of this invention.

Additional disorders which may be treated or prevented using the compounds of this invention are immunological disorders such as autoimmune disease (AIDS) and cardiovascular disorders such as atherosclerosis.

It is as aspect of this invention that a chemical compound that modulates the catalytic activity of a protein kinase may be identified by contacting cells expressing said protein kinase with a compound, salt or prodrug of the present invention and then monitoring said cells for an effect.

By "monitoring" is meant observing or detecting the effect of contacting a compound with a cell expressing a

WO 00/08202

77

PCT/US99/17845

particular PK. The observed or detected effect can be a change in cell phenotype, in the catalytic activity of a PK or a change in the interaction of a PK with a natural binding partner. Techniques for observing or detecting such effects are well-known in the art.

The above-referenced effect is selected from a change or an absence of change in a cell phenotype, a change or absence of change in the catalytic activity of said protein kinase or a change or absence of change in the interaction of said protein kinase with a natural binding partner in a final aspect of this invention.

"Cell phenotype" refers to the outward appearance of a cell or tissue or the biological function of the cell or tissue. Examples, without limitation, of a cell phenotype are cell size, cell growth, cell proliferation, cell differentiation, cell survival, apoptosis, and nutrient uptake and use. Such phenotypic characteristics are measurable by techniques well-known in the art.

A "natural binding partner" refers to a polypeptide that binds to a particular PK in a cell. Natural binding partners can play a role in propagating a signal in a PK-mediated signal transduction process. A change in the interaction of the natural binding partner with the PK can manifest itself as an increased or decreased concentration of the PK/natural binding partner complex and, as a result, in an observable change in the ability of the PK to mediate signal transduction.

It is also an aspect of this invention that a compound described herein, or its salt or prodrug, might be combined with other chemotherapeutic agents for the treatment of the diseases and disorders discussed above. For instance, a compound, salt or prodrug of this invention might be combined with alkylating agents such as fluorouracil (5-FU) alone or in further combination with leukovorin; or other alkylating

WO 00/08202

78

PCT/US99/17845

agents such as, without limitation, other pyrimidine analogs such as UFT, capecitabine, gemcitabine and cytarabine, the alkyl sulfonates, e.g., busulfan (used in the treatment of chronic granulocytic leukemia), improsulfan and piposulfan; 5 aziridines, e.g., benzodepa, carboquone, meturedepa and uredepa; ethyleneimines and methylmelamines, e.g., altretamine, triethylenemelamine, triethylenephosphoramidate, triethylenethiophosphoramidate and trimethylolmelamine; and the nitrogen mustards, e.g., chlorambucil (used in the treatment 10 of chronic lymphocytic leukemia, primary macroglobulinemia and non-Hodgkin's lymphoma), cyclophosphamide (used in the treatment of Hodgkin's disease, multiple myeloma, neuroblastoma, breast cancer, ovarian cancer, lung cancer, Wilm's tumor and rhabdomyosarcoma), estramustine, ifosfamide, 15 novembrichin, prednimustine and uracil mustard (used in the treatment of primary thrombocytosis, non-Hodgkin's lymphoma, Hodgkin's disease and ovarian cancer); and triazines, e.g., dacarbazine (used in the treatment of soft tissue sarcoma).

Likewise a compound, salt or prodrug of this invention 20 might be expected to have a beneficial effect in combination with other antimetabolite chemotherapeutic agents such as, without limitation, folic acid analogs, e.g. methotrexate (used in the treatment of acute lymphocytic leukemia, choriocarcinoma, mycosis fungoides breast cancer, head and 25 neck cancer and osteogenic sarcoma) and pteropterin; and the purine analogs such as mercaptopurine and thioguanine which find use in the treatment of acute granulocytic, acute lymphocytic and chronic granulocytic leukemias.

A compound, salt or prodrug of this invention might also 30 be expected to prove efficacious in combination with natural product based chemotherapeutic agents such as, without limitation, the vinca alkaloids, e.g., vinblastin (used in the treatment of breast and testicular cancer), vincristine and vindesine; the epipodophylotoxins, e.g., etoposide and

WO 00/08202

79

PCT/US99/17845

teniposide, both of which are useful in the treatment of testicular cancer and Kaposi's sarcoma; the antibiotic chemotherapeutic agents, e.g., daunorubicin, doxorubicin, epirubicin, mitomycin (used to treat stomach, cervix, colon, breast, bladder and pancreatic cancer), dactinomycin, temozolomide, plicamycin, bleomycin (used in the treatment of skin, esophagus and genitourinary tract cancer); and the enzymatic chemotherapeutic agents such as L-asparaginase.

In addition to the above, a compound, salt or prodrug of this invention might be expected to have a beneficial effect used in combination with the platinum coordination complexes (cisplatin, etc.); substituted ureas such as hydroxyurea; methylhydrazine derivatives, e.g., procarbazine; adrenocortical suppressants, e.g., mitotane, aminogluthethimide; and hormone and hormone antagonists such as the adrenocorticosteroids (e.g., prednisone), progestins (e.g., hydroxyprogesterone caproate); estrogens (e.g., diethylstilbesterol); antiestrogens such as tamoxifen; androgens, e.g., testosterone propionate; and aromatase inhibitors (such as anastrozole).

Finally, the combination of a compound of this invention might be expected to be particularly effective in combination with mitoxantrone or paclitaxel for the treatment of solid tumor cancers or leukemias such as, without limitation, acute myelogenous (non-lymphocytic) leukemia.

#### DETAILED DESCRIPTION OF THE INVENTION

##### 1. BRIEF DESCRIPTION OF THE TABLES

TABLE 1 shows the chemical structures of some exemplary compounds of this invention. The compound numbers correspond to the Example numbers in the Examples section. That is, the synthesis of Compound 1 in Table 1 is described in Example 1. The compounds presented in Table 1 are exemplary only and are not to be construed as limiting the scope of this invention in any manner.

WO 00/08202

80

PCT/US99/17845 ,

TABLE 2 shows the results of biological testing of some exemplary compounds of this invention. The results are reported in terms of  $IC_{50}$ , the micromolar ( $\mu M$ ) concentration of the compound being tested which causes a 50% change in the activity of the target PKT compared to the activity of the PTK in a control to which no test compound has been added. Specifically, the results shown indicate the concentration of a test compound needed to cause a 50% reduction of the activity of the target PTK. The bioassays used are described in detail below.

## 2. INDICATIONS/TARGET DISEASES

The PKs whose catalytic activity is modulated by the compounds of this invention include protein tyrosine kinases of which there are two types, receptor tyrosine kinases (RTKs) and cellular tyrosine kinases (CTKs), and serine-threonine kinases (STKs). RTK mediated signal transduction, is initiated by extracellular interaction with a specific growth factor (ligand), followed by receptor dimerization, transient stimulation of the intrinsic protein tyrosine kinase activity and phosphorylation. Binding sites are thereby created for intracellular signal transduction molecules and lead to the formation of complexes with a spectrum of cytoplasmic signaling molecules that facilitate the appropriate cellular response (e.g., cell division, metabolic effects on the extracellular microenvironment, etc.). See, Schlessinger and Ullrich, 1992, Neuron 9:303-391.

It has been shown that tyrosine phosphorylation sites on growth factor receptors function as high-affinity binding sites for SH2 (src homology) domains of signaling molecules. Fantl et al., 1992, Cell 69:413-423, Songyang et al., 1994, Mol. Cell. Biol. 14:2777-2785), Songyang et al., 1993, Cell 72:767-778, and Koch et al., 1991, Science 252:668-678. Several intracellular substrate proteins that associate with RTKs have been identified. They may be divided into two

WO 00/08202

81

PCT/US99/17845 ,

principal groups: (1) substrates that have a catalytic domain, and (2) substrates which lack such domain but which serve as adapters and associate with catalytically active molecules. Songyang et al., 1993, Cell 72:767-778. The specificity of the interactions between receptors and SH2 domains of their substrates is determined by the amino acid residues immediately surrounding the phosphorylated tyrosine residue. Differences in the binding affinities between SH2 domains and the amino acid sequences surrounding the phosphotyrosine residues on particular receptors are consistent with the observed differences in their substrate phosphorylation profiles. Songyang et al., 1993, Cell 72:767-778. These observations suggest that the function of each RTK is determined not only by its pattern of expression and ligand availability but also by the array of downstream signal transduction pathways that are activated by a particular receptor. Thus, phosphorylation provides an important regulatory step which determines the selectivity of signaling pathways recruited by specific growth factor receptors, as well as differentiation factor receptors.

STKs, being primarily cytosolic, affect the internal biochemistry of the cell, often as a down-line response to a PTK event. STKs have been implicated in the signaling process which initiates DNA synthesis and subsequent mitosis leading to cell proliferation.

Thus, PK signal transduction results in, among other responses, cell proliferation, differentiation, growth and metabolism. Abnormal cell proliferation may result in a wide array of disorders and diseases, including the development of neoplasia such as carcinoma, sarcoma, glioblastoma and hemangioma, disorders such as leukemia, psoriasis, arteriosclerosis, arthritis and diabetic retinopathy and other disorders related to uncontrolled angiogenesis and/or vasculogenesis.



WO 00/08202

82

PCT/US99/17845 ,

A precise understanding of the mechanism by which the compounds of this invention inhibit PKs is not required in order to practice the present invention. However, while not hereby being bound to any particular mechanism or theory, it is believed that the compounds interact with the amino acids in the catalytic region of PKs. PKs typically possess a bilobate structure wherein ATP appears to bind in the cleft between the two lobes in a region where the amino acids are conserved among PKs. Inhibitors of PKs are believed to bind by non-covalent interactions such as hydrogen bonding, van der Waals forces and ionic interactions in the same general region where the aforesaid ATP binds to the PKs. More specifically, it is thought that the 2-indolinone component of the compounds of this invention binds in the general space normally occupied by the adenine ring of ATP. Specificity of a particular molecule for a particular PK may then arise as the result of additional interactions between the various substituents on the 2-indolinone core and the amino acid domains specific to particular PKs. Thus, different indolinone substituents may contribute to preferential binding to particular PKs. The ability to select compounds active at different ATP (or other nucleotide) binding sites makes the compounds of this invention useful for targeting any protein with such a site. The compounds disclosed herein may thus have utility as in vitro assays for such proteins as well as exhibiting in vivo therapeutic effects through interaction with such proteins.

In another aspect, the protein kinase, the catalytic activity of which is modulated by contact with a compound of this invention, is a protein tyrosine kinase, more particularly, a receptor protein tyrosine kinase. Among the receptor protein tyrosine kinases whose catalytic activity can be modulated with a compound of this invention, or salt thereof, are, without limitation, EGF, HER2, HER3, HER4, IR,

WO 00/08202

83

PCT/US99/17845 ,

IGF-1R, IRR, PDGFR $\alpha$ , PDGFR $\beta$ , CSF1R, C-Kit, C-fms, Flk-1R, Flk4, KDR/Flk-1, Flt-1, FGFR-1R, FGFR-2R, FGFR-3R and FGFR-4R.

The protein tyrosine kinase whose catalytic activity is modulated by contact with a compound of this invention, or a salt or a prodrug thereof, can also be a non-receptor or cellular protein tyrosine kinase (CTK). Thus, the catalytic activity of CTKs such as, without limitation, Src, Frk, Btk, Csk, Abl, ZAP70, Fes, Fps, Fak, Jak, Ack, Yes, Fyn, Lyn, Lck, Blk, Hck, Fgr and Yrk, may be modulated by contact with a compound or salt of this invention.

Still another group of PKs which may have their catalytic activity modulated by contact with a compound of this invention are the serine-threonine protein kinases such as, without limitation, CDK2 and Raf.

In another aspect, this invention relates to a method for treating or preventing a PK related disorder by administering a therapeutically effective amount of a compound of this invention, or a salt or a prodrug thereof, to an organism.

It is also an aspect of this invention that a pharmaceutical composition containing a compound of this invention or a salt or prodrug thereof is administered to an organism for the purpose of preventing or treating a PK related disorder.

This invention is therefore directed to compounds that modulate PK signal transduction by affecting the enzymatic activity of RTKs, CTKs and/or STKs, thereby interfering with the signals transduced by such proteins. More particularly, the present invention is directed to compounds which modulate RTK, CTK and/or STK mediated signal transduction pathways as a therapeutic approach to the treatment of many kinds of solid tumors, including but not limited to carcinomas, sarcomas including Kaposi's sarcoma, erythroblastoma, glioblastoma, meningioma, astrocytoma, melanoma and myoblastoma. Treatment

WO 00/08202

84

PCT/US99/17845

or prevention of non-solid tumor cancers such as leukemia are also contemplated by this invention. Indications may include, but are not limited to brain cancers, bladder cancers, ovarian cancers, gastric cancers, pancreas cancers, colon cancers, blood cancers, lung cancers and bone cancers.

Further examples, without limitation, of the types of disorders related to inappropriate PK activity that the compounds described herein may be useful in preventing, treating and studying, are cell proliferative disorders, fibrotic disorders and metabolic disorders.

Cell proliferative disorders, which may be prevented, treated or further studied by the present invention include cancer, blood vessel proliferative disorders and mesangial cell proliferative disorders.

Blood vessel proliferative disorders refer to disorders related to abnormal vasculogenesis (blood vessel formation) and angiogenesis (spreading of blood vessels). While vasculogenesis and angiogenesis play important roles in a variety of normal physiological processes such as embryonic development, corpus luteum formation, wound healing and organ regeneration, they also play a pivotal role in cancer development where they result in the formation of new capillaries needed to keep a tumor alive. Other examples of blood vessel proliferation disorders include arthritis, where new capillary blood vessels invade the joint and destroy cartilage, and ocular diseases, like diabetic retinopathy, where new capillaries in the retina invade the vitreous, bleed and cause blindness.

Two structurally related RTKs have been identified to bind VEGF with high affinity: the fms-like tyrosine 1 (fit-1) receptor (Shibuya et al., 1990, Oncogene, 5:519-524; De Vries et al., 1992, Science, 255:989-991) and the KDR/FLK-1 receptor, also known as VEGF-R2. Vascular endothelial growth factor (VEGF) has been reported to be an endothelial cell

WO 00/08202

85

PCT/US99/17845 ,

specific mitogen with in vitro endothelial cell growth promoting activity. Ferrara & Henzel, 1989, Biochein. Biophys. Res. Comm., 161:851-858; Vaisman et al., 1990, J. Biol. Chem., 265:19461-19566. Information set forth in U.S. application

5 Ser. Nos. 08/193,829, 08/038,596 and 07/975,750, strongly suggest that VEGF is not only responsible for endothelial cell proliferation, but also is the prime regulator of normal and pathological angiogenesis. See generally, Klagsburn & Soker, 1993, Current Biology, 3(10):699-702; Houck, et al., 1992, J.

10 Biol. Chem., 267:26031-26037.

Normal vasculogenesis and angiogenesis play important roles in a variety of physiological processes such as embryonic development, wound healing, organ regeneration and female reproductive processes such as follicle development in

15 the corpus luteum during ovulation and placental growth after pregnancy. Folkman & Shing, 1992, J. Biological Chem., 267(16):10931-34. Uncontrolled vasculogenesis and/or angiogenesis has been associated with diseases such as diabetes as well as with malignant solid tumors that rely on

20 vascularization for growth. Klagsburn & Soker, 1993, Current Biology, 3(10):699-702; Folkham, 1991, J. Natl. Cancer Inst., 82:4-6; Weidner, et al., 1991, New Engl. J. Med., 324:1-5.

The surmised role of VEGF in endothelial cell proliferation and migration during angiogenesis and

25 vasculogenesis indicates an important role for the KDR/FLK-1 receptor in these processes. Diseases such as diabetes mellitus (Folkman, 198, in XIth Congress of Thrombosis and Haemostasis (Verstraeta, et al., eds.), pp. 583-596, Leuven University Press, Leuven) and arthritis, as well as malignant

30 tumor growth may result from uncontrolled angiogenesis. See e.g., Folkman, 1971, N. Engl. J. Med., 285:1182-1186. The receptors to which VEGF specifically binds are an important and powerful therapeutic target for the regulation and modulation of vasculogenesis and/or angiogenesis and a variety

WO 00/08202

86

PCT/US99/17845

of severe diseases which involve abnormal cellular growth caused by such processes. Plowman, et al., 1994, DN&P, 7(6):334-339. More particularly, the KDR/FLK-1 receptor's highly specific role in neovascularization make it a choice  
5 target for therapeutic approaches to the treatment of cancer and other diseases which involve the uncontrolled formation of blood vessels.

Thus, one aspect of the present invention relates to compounds capable of regulating and/or modulating tyrosine  
10 kinase signal transduction including KDR/FLK-1 receptor signal transduction in order to inhibit or promote angiogenesis and/or vasculogenesis, that is, compounds that inhibit, prevent, or interfere with the signal transduced by KDR/FLK-1 when activated by ligands such as VEGF. Although it is  
15 believed that the compounds of the present invention act on a receptor or other component along the tyrosine kinase signal transduction pathway, they may also act directly on the tumor cells that result from uncontrolled angiogenesis.

Although the nomenclature of the human and murine  
20 counterparts of the generic "flk-I" receptor differ, they are, in many respects, interchangeable. The murine receptor, Flk-1, and its human counterpart, KDR, share a sequence homology of 93.4% within the intracellular domain. Likewise, murine FLK-1 binds human VEGF with the same affinity as mouse VEGF, and  
25 accordingly, is activated by the ligand derived from either species. Millauer et al., 1993, Cell, 72:835-846; Quinn et al., 1993, Proc. Natl. Acad. Sci. USA, 90:7533-7537. FLK-1 also associates with and subsequently tyrosine phosphorylates human RTK substrates (e.g., PLC- $\gamma$  or p85) when co-expressed in  
30 293 cells (human embryonal kidney fibroblasts).

Models which rely upon the FLK-1 receptor therefore are directly applicable to understanding the KDR receptor. For example, use of the murine FLK-1 receptor in methods which

WO 00/08202

87

PCT/US99/17845

identify compounds that regulate the murine signal transduction pathway are directly applicable to the identification of compounds which may be used to regulate the human signal transduction pathway, that is, which regulate activity related to the KDR receptor. Thus, chemical compounds identified as inhibitors of KDR/FLK-1 in vitro, can be confirmed in suitable in vivo models. Both in vivo mouse and rat animal models have been demonstrated to be of excellent value for the examination of the clinical potential of agents acting on the KDR/FLK-1 induced signal transduction pathway.

Thus, in one aspect, this invention is directed to compounds that regulate, modulate and/or inhibit vasculogenesis and/or angiogenesis by affecting the enzymatic activity of the KDR/FLK-1 receptor and interfering with the signal transduced by KDR/FLK-1. In another aspect, the present invention is directed to compounds which regulate, modulate and/or inhibit the KDR/FLK-1 mediated signal transduction pathway as a therapeutic approach to the treatment of many kinds of solid tumors including, but not limited to, glioblastoma, melanoma and Kaposi's sarcoma, and ovarian, lung, mammary, prostate, pancreatic, colon and epidermoid carcinoma. In addition, data suggest the administration of compounds which inhibit the KDR/Flk-1 mediated signal transduction pathway may also be used in the treatment of hemangioma, restenosis and diabetic retinopathy.

A further aspect of this invention relates to the inhibition of vasculogenesis and angiogenesis by other receptor-mediated pathways, including the pathway comprising the flt-1 receptor.

Receptor tyrosine kinase mediated signal transduction is initiated by extracellular interaction with a specific growth factor (ligand), followed by receptor dimerization, transient stimulation of the intrinsic protein tyrosine kinase activity and autophosphorylation. Binding sites are thereby created for

WO 00/08202

88

PCT/US99/17845

intracellular signal transduction molecules which leads to the formation of complexes with a spectrum of cytoplasmic signalling molecules that facilitate the appropriate cellular response, e.g., cell division and metabolic effects to the extracellular microenvironment. See, Schlessinger and Ullrich, 1992, Neuron, 9:1-20.

The close homology of the intracellular regions of KDR/FLK-1 with that of the PDGF- $\beta$  receptor (50.3% homology) and/or the related flt-1 receptor indicates the induction of overlapping signal transduction pathways. For example, for the PDGF- $\beta$  receptor, members of the src family (Twamley et al., 1993, Proc. Natl. Acad. Sci. USA, 90:7696-7700), phosphatidylinositol-3'-kinase (Hu et al., 1992, Mol. Cell. Biol., 12:981-990), phospholipase  $\gamma$  (Kashishian & Cooper, 1993, Mol. Cell. Biol., 4:49-51), ras-GTPase-activating protein, (Kashishian et al., 1992, EMBO J., 11:1373-1382), PTP-ID/syp (Kazlauskas et al., 1993, Proc. Natl. Acad. Sci. USA, 10 90:6939-6943), Grb2 (Arvidsson et al., 1994, Mol. Cell. Biol., 14:6715-6726), and the adapter molecules Shc and Nck (Nishimura et al., 1993, Mol. Cell. Biol., 13:6889-6896), have been shown to bind to regions involving different autophosphorylation sites. See generally, Claesson-Welsh, 1994, Prog. Growth Factor Res., 5:37-54. Thus, it is likely that signal transduction pathways activated by KDR/FLK-1 include the ras pathway (Rozakis et al., 1992, Nature, 360:689-692), the PI-3'-kinase, the src-mediated and the plcy-mediated pathways. Each of these pathways may play a critical role in the angiogenic and/or vasculogenic effect of KDR/FLK-1 in endothelial cells. Consequently, a still further aspect of this invention relates to the use of the organic compounds described herein to modulate angiogenesis and vasculogenesis as such processes are controlled by these pathways.

WO 00/08202

89

PCT/US99/17845 ,

Conversely, disorders related to the shrinkage, contraction or closing of blood vessels, such as restenosis, are also implicated and may be treated or prevented by the methods of this invention.

5       Fibrotic disorders refer to the abnormal formation of extracellular matrices. Examples of fibrotic disorders include hepatic cirrhosis and mesangial cell proliferative disorders. Hepatic cirrhosis is characterized by the increase in extracellular matrix constituents resulting in the  
10       formation of a hepatic scar. An increased extracellular matrix resulting in a hepatic scar can also be caused by a viral infection such as hepatitis. Lipocytes appear to play a major role in hepatic cirrhosis. Other fibrotic disorders implicated include atherosclerosis.

15       Mesangial cell proliferative disorders refer to disorders brought about by abnormal proliferation of mesangial cells. Mesangial proliferative disorders include various human renal diseases such as glomerulonephritis, diabetic nephropathy and malignant nephrosclerosis as well as such disorders as  
20       thrombotic microangiopathy syndromes, transplant rejection, and glomerulopathies. The RTK PDGFR has been implicated in the maintenance of mesangial cell proliferation. Floege et al., 1993, Kidney International 43:47S-54S.

25       Many cancers are cell proliferative disorders and, as noted previously, PKs have been associated with cell proliferative disorders. Thus, it is not surprising that PKs such as, for example, members of the RTK family have been associated with the development of cancer. Some of these receptors, like EGFR (Tuzi et al., 1991, Br. J. Cancer 63:227-  
30       233, Torp et al., 1992, APMIS 100:713-719) HER2/neu (Slamon et al., 1989, Science 244:707-712) and PDGF-R (Kumabe et al., 1992, Oncogene, 7:627-633) are over-expressed in many tumors and/or persistently activated by autocrine loops. In fact, in the most common and severe cancers these receptor over-



WO 00/08202

90

PCT/US99/17845 ,

expressions (Akbasak and Suner-Akbasak et al., 1992, J. Neurol. Sci., 111:119-133, Dickson et al., 1992, Cancer Treatment Res. 61:249-273, Korc et al., 1992, J. Clin. Invest. 90:1352-1360) and autocrine loops (Lee and Donoghue, 1992, J. Cell. Biol., 118:1057-1070, Korc et al., supra, Akbasak and Suner-Akbasak et al., supra) have been demonstrated. For example, EGFR has been associated with squamous cell carcinoma, astrocytoma, glioblastoma, head and neck cancer, lung cancer and bladder cancer. HER2 has been associated with breast, ovarian, gastric, lung, pancreas and bladder cancer. PDGFR has been associated with glioblastoma and melanoma as well as lung, ovarian and prostate cancer. The RTK c-met has also been associated with malignant tumor formation. For example, c-met has been associated with, among other cancers, colorectal, thyroid, pancreatic, gastric and hepatocellular carcinomas and lymphomas. Additionally c-met has been linked to leukemia. Over-expression of the c-met gene has also been detected in patients with Hodgkins disease and Burkitts disease.

IGF-IR, in addition to being implicated in nutritional support and in type-II diabetes, has also been associated with several types of cancers. For example, IGF-I has been implicated as an autocrine growth stimulator for several tumor types, e.g. human breast cancer carcinoma cells (Arteaga et al., 1989, J. Clin. Invest. 84:1418-1423) and small lung tumor cells (Macauley et al., 1990, Cancer Res., 50:2511-2517). In addition, IGF-I, while integrally involved in the normal growth and differentiation of the nervous system, also appears to be an autocrine stimulator of human gliomas. Sandberg-Nordqvist et al., 1993, Cancer Res. 53:2475-2478. The importance of IGF-IR and its ligands in cell proliferation is further supported by the fact that many cell types in culture (fibroblasts, epithelial cells, smooth muscle cells, T-lymphocytes, myeloid cells, chondrocytes and osteoblasts (the stem cells of the bone marrow))

WO 00/08202

91

PCT/US99/17845

are stimulated to grow by IGF-I. Goldring and Goldring, 1991, Eukaryotic Gene Expression, 1:301-326. In a series of recent publications, Baserga suggests that IGF-IR plays a central role in the mechanism of transformation and, as such, could be a preferred target for therapeutic interventions for a broad spectrum of human malignancies. Baserga, 1995, Cancer Res., 55:249-252, Baserga, 1994, Cell 79:927-930, Coppola et al., 1994, Mol. Cell. Biol., 14:4588-4595.

STKs have been implicated in many types of cancer including, notably, breast cancer (Cance, et al., Int. J. Cancer, 54:571-77 (1993)).

The association between abnormal PK activity and disease is not restricted to cancer. For example, RTKs have been associated with diseases such as psoriasis, diabetes mellitus, endometriosis, angiogenesis, atheromatous plaque development, Alzheimer's disease, von Hippel-Lindau disease, epidermal hyperproliferation, neurodegenerative diseases, age-related macular degeneration and hemangiomas. For example, EGFR has been indicated in corneal and dermal wound healing. Defects in Insulin-R and IGF-1R are indicated in type-II diabetes mellitus. A more complete correlation between specific RTKs and their therapeutic indications is set forth in Plowman et al., 1994, DN&P 7:334-339.

As noted previously, not only RTKs but CTKs including, but not limited to, src, abl, fps, yes, fyn, lyn, lck, blk, hck, fgr and yrk (reviewed by Bolen et al., 1992, FASEB J., 6:3403-3409) are involved in the proliferative and metabolic signal transduction pathway and thus could be expected, and have been shown, to be involved in many PTK-mediated disorders to which the present invention is directed. For example, mutated src (v-src) has been shown to be an oncoprotein (pp60<sup>v-src</sup>) in chicken. Moreover, its cellular homolog, the proto-oncogene pp60<sup>c-src</sup> transmits oncogenic signals of many receptors. Over-expression of EGFR or HER2/neu in tumors leads to the constitutive

WO 00/08202

92

PCT/US99/17845

activation of pp60<sup>c src</sup>, which is characteristic of malignant cells but absent in normal cells. On the other hand, mice deficient in the expression of c-src exhibit an osteopetrotic phenotype, indicating a key participation of c-src in osteoclast function  
5 and a possible involvement in related disorders.

Similarly, Zap70 has been implicated in T-cell signaling which may relate to autoimmune disorders.

STKs have been associated with inflammation, autoimmune disease, immunoresponses, and hyperproliferation disorders such as  
10 restenosis, fibrosis, psoriasis, osteoarthritis and rheumatoid arthritis.

PKs have also been implicated in embryo implantation. Thus, the compounds of this invention may provide an effective method of preventing such embryo implantation and thereby be useful as  
15 birth control agents.

Finally, both RTKs and CTGs are currently suspected as being involved in hyperimmune disorders.

A method for identifying a chemical compound that modulates the catalytic activity of one or more of the above  
20 discussed protein kinases is another aspect of this invention.

The method involves contacting cells expressing the desired protein kinase with a compound of this invention (or its salt or prodrug) and monitoring the cells for any effect that the compound has on them. The effect may be any observable,  
25 either to the naked eye or through the use of instrumentation, change or absence of change in a cell phenotype. The change or absence of change in the cell phenotype monitored may be, for example, without limitation, a change or absence of change in the catalytic activity of the protein kinase in the cells  
30 or a change or absence of change in the interaction of the protein kinase with a natural binding partner.

Examples of the effect of a number of exemplary compounds of this invention on several PTKs are shown in Table 2. The

WO 00/08202

93

PCT/US99/17845

compounds and data presented are not to be construed as limiting the scope of this invention in any manner whatsoever.

### 3. PHARMACEUTICAL COMPOSITIONS AND USE

A compound of the present invention, a prodrug thereof or  
5 a physiologically acceptable salt of either the compound or  
its prodrug, can be administered as such to a human patient or  
can be administered in pharmaceutical compositions in which  
the foregoing materials are mixed with suitable carriers or  
excipient(s). Techniques for formulation and administration  
10 of drugs may be found in "Remington's Pharmacological  
Sciences," Mack Publishing Co., Easton, PA, latest edition.

#### Routes of Administration.

As used herein, "administer" or "administration" refers  
to the delivery of a compound, salt or prodrug of the present  
15 invention or of a pharmaceutical composition containing a  
compound, salt or prodrug of this invention to an organism for  
the purpose of prevention or treatment of a PK-related  
disorder.

Suitable routes of administration may include, without  
20 limitation, oral, rectal, transmucosal or intestinal  
administration or intramuscular, subcutaneous, intramedullary,  
intrathecal, direct intraventricular, intravenous,  
intravitreal, intraperitoneal, intranasal, or intraocular  
injections. The preferred routes of administration are oral  
25 and parenteral.

Alternatively, one may administer the compound in a local  
rather than systemic manner, for example, via injection of the  
compound directly into a solid tumor, often in a depot or  
sustained release formulation.

30 Furthermore, one may administer the drug in a targeted  
drug delivery system, for example, in a liposome coated with  
tumor-specific antibody. The liposomes will be targeted to  
and taken up selectively by the tumor.

WO 00/08202

94

PCT/US99/17845

Composition/Formulation.

Pharmaceutical compositions of the present invention may be manufactured by processes well known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Pharmaceutical compositions for use in accordance with the present invention may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

For injection, the compounds of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, lozenges, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient. Pharmaceutical preparations for oral use can be made using a solid excipient, optionally grinding the resulting mixture, and processing the mixture of granules, after adding other suitable auxiliaries if desired, to obtain tablets or dragee cores. Useful excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol, cellulose preparations such as, for example, maize starch, wheat starch, rice starch and

WO 00/08202

95

PCT/US99/17845 ,

potato starch and other materials such as gelatin, gum  
tragacanth, methyl cellulose, hydroxypropylmethyl- cellulose,  
sodium carboxymethylcellulose, and/or polyvinyl- pyrrolidone  
(PVP). If desired, disintegrating agents may be added, such  
5 as cross-linked polyvinyl pyrrolidone, agar, or alginic acid.  
A salt such as sodium alginate may also be used.

Dragee cores are provided with suitable coatings. For  
this purpose, concentrated sugar solutions may be used which  
may optionally contain gum arabic, talc, polyvinyl  
10 pyrrolidone, carbopol gel, polyethylene glycol, and/or  
titanium dioxide, lacquer solutions, and suitable organic  
solvents or solvent mixtures. Dyestuffs or pigments may be  
added to the tablets or dragee coatings for identification or  
to characterize different combinations of active compound  
15 doses.

Pharmaceutical compositions which can be used orally  
include push-fit capsules made of gelatin, as well as soft,  
sealed capsules made of gelatin and a plasticizer, such as  
glycerol or sorbitol. The push-fit capsules can contain the  
20 active ingredients in admixture with a filler such as lactose,  
a binder such as starch, and/or a lubricant such as talc or  
magnesium stearate and, optionally, stabilizers. In soft  
capsules, the active compounds may be dissolved or suspended  
in suitable liquids, such as fatty oils, liquid paraffin, or  
25 liquid polyethylene glycols. Stabilizers may be added in  
these formulations, also.

For administration by inhalation, the compounds for use  
according to the present invention are conveniently delivered  
in the form of an aerosol spray using a pressurized pack or a  
30 nebulizer and a suitable propellant, e.g., without limitation,  
dichlorodifluoromethane, trichlorofluoromethane,  
dichlorotetra- fluoroethane or carbon dioxide. In the case of  
a pressurized aerosol, the dosage unit may be controlled by  
providing a valve to deliver a metered amount. Capsules and

WO 00/08202

96

PCT/US99/17845 ,

cartridges of, for example, gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds may also be formulated for parenteral administration, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulating materials such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical compositions for parenteral administration include aqueous solutions of a water soluble form, such as, without limitation, a salt, of the active compound. Additionally, suspensions of the active compounds may be prepared in a lipophilic vehicle. Suitable lipophilic vehicles include fatty oils such as sesame oil, synthetic fatty acid esters such as ethyl oleate and triglycerides, or materials such as liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers and/or agents that increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile, pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, using, e.g., conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as depot preparations. Such

WO 00/08202

97

PCT/US99/17845 ,

long acting formulations may be administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection. A compound of this invention may be formulated for this route of administration with suitable  
5 polymeric or hydrophobic materials (for instance, in an emulsion with a pharmaceutically acceptable oil), with ion exchange resins, or as a sparingly soluble derivative such as, without limitation, a sparingly soluble salt.

A non-limiting example of a pharmaceutical carrier for  
10 the hydrophobic compounds of the invention is a cosolvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer and an aqueous phase such as the VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant Polysorbate 80™,  
15 and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:D5W) consists of VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic  
20 administration. Naturally, the proportions of such a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used  
25 instead of Polysorbate 80™, the fraction size of polyethylene glycol may be varied, other biocompatible polymers may replace polyethylene glycol, e.g., polyvinyl pyrrolidone, and other sugars or polysaccharides may substitute for dextrose.

Alternatively, other delivery systems for hydrophobic  
30 pharmaceutical compounds may be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. In addition, certain organic



WO 00/08202

98

PCT/US99/17845 ,

solvents such as dimethylsulfoxide also may be employed, although often at the cost of greater toxicity.

Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

The pharmaceutical compositions herein also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include, but are not limited to, calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

Many of the PK modulating compounds of the invention may be provided as physiologically acceptable salts wherein the claimed compound may form the negatively or the positively charged species. Examples of salts in which the compound forms the positively charged moiety include, without limitation, quaternary ammonium (defined elsewhere herein), salts such as the hydrochloride, sulfate, carbonate, lactate, tartrate, maleate, succinate wherein the nitrogen atom of the quaternary ammonium group is a nitrogen of the selected compound of this invention which has reacted with the appropriate acid. Salts in which a compound of this invention forms the negatively charged species include, without limitation, the sodium, potassium, calcium and magnesium salts formed by the reaction of a carboxylic acid group in the compound with an appropriate base (e.g. sodium hydroxide (NaOH), potassium hydroxide (KOH), Calcium hydroxide (Ca(OH)<sub>2</sub>), etc.).

WO 00/08202

99

PCT/US99/17845

Dosage.

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an amount sufficient to achieve  
5 the intended purpose, i.e., the modulation of PK activity or the treatment or prevention of a PK-related disorder.

More specifically, a therapeutically effective amount means an amount of compound effective to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the  
10 subject being treated.

Determination of a therapeutically effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

15 For any compound used in the methods of the invention, the therapeutically effective amount or dose can be estimated initially from cell culture assays. Then, the dosage can be formulated for use in animal models so as to achieve a circulating concentration range that includes the  $IC_{50}$  as  
20 determined in cell culture (i.e., the concentration of the test compound which achieves a half-maximal inhibition of the PK activity). Such information can then be used to more accurately determine useful doses in humans.

Toxicity and therapeutic efficacy of the compounds  
25 described herein can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., by determining the  $IC_{50}$  and the  $LD_{50}$  (both of which are discussed elsewhere herein) for a subject compound. The data obtained from these cell culture assays and animal studies can be used  
30 in formulating a range of dosage for use in humans. The dosage may vary depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See e.g.,

WO 00/08202

100

PCT/US99/17845

Fingl, et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1).

Dosage amount and interval may be adjusted individually to provide plasma levels of the active species which are  
5 sufficient to maintain the kinase modulating effects. These plasma levels are referred to as minimal effective concentrations (MECs). The MEC will vary for each compound but can be estimated from in vitro data, e.g., the  
10 concentration necessary to achieve 50-90% inhibition of a kinase may be ascertained using the assays described herein. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. HPLC assays or bioassays can be used to determine plasma concentrations.

Dosage intervals can also be determined using MEC value.  
15 Compounds should be administered using a regimen that maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%.

In cases of local administration or selective uptake, the effective local concentration of the drug may not be related  
20 to plasma concentration and other procedures known in the art may be employed to determine the correct dosage amount and interval.

The amount of a composition administered will, of course, be dependent on the subject being treated, the severity of the  
25 affliction, the manner of administration, the judgment of the prescribing physician, etc.

#### Packaging.

The compositions may, if desired, be presented in a pack or dispenser device, such as an FDA approved kit, which may  
30 contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser may also be accompanied by a notice

WO 00/08202

101

PCT/US99/17845

associated with the container in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the compositions or of human or veterinary administration. Such notice, for example, may be of the labeling approved by the U.S. Food and Drug Administration for prescription drugs or of an approved product insert. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition. Suitable conditions indicated on the label may include treatment of a tumor, inhibition of angiogenesis, treatment of fibrosis, diabetes, and the like.

15

#### 4. SYNTHESIS

The compounds of this invention, as well as the precursor oxindoles and aldehydes, may be readily synthesized using techniques well known in the chemical arts. It will be appreciated by those skilled in the art that other synthetic pathways for forming the compounds of the invention are available and that the following is offered by way of example and not limitation.

##### General synthetic procedures.

The following general methodologies may be used to prepare intermediates to and compounds of this invention.

##### Preparation of substituted oxindoles

##### 5-Amino-2-oxindole

5-Nitro-2-oxindole (6.3 g) was hydrogenated in methanol over 10% palladium on carbon to give 3.0 g (60% yield) of the title compound as a white solid.

##### 5-Bromo-2-oxindole

2-Oxindole (1.3 g) in 20 mL acetonitrile was cooled to -

WO 00/08202

102

PCT/US99/17845

10 °C and 2.0 g N-bromosuccinimide was slowly added with stirring. The reaction was stirred for 1 hour at -10 °C and 2 hours at 0 °C. The precipitate was collected, washed with water and dried to give 1.9 g (90 % yield) of the title compound.

5 **4-Methyl-2-oxindole**

Diethyl oxalate (30 mL) in 20 mL of dry ether was added with stirring to 19 g of potassium ethoxide suspended in 50 mL of dry ether. The mixture was cooled in an ice bath and 20 mL of 3-nitro-o-xylene in 20 mL of dry ether was slowly added.

10 The thick dark red mixture was heated to reflux for 0.5 hr, concentrated to a dark red solid, and treated with 10% sodium hydroxide until almost all of the solid dissolved. The dark red mixture was treated with 30% hydrogen peroxide until the red color changed to yellow. The mixture was treated  
15 alternately with 10% sodium hydroxide and 30% hydrogen peroxide until the dark red color was no longer present. The solid was filtered off and the filtrate acidified with 6N hydrochloric acid. The resulting precipitate was collected by vacuum filtration, washed with water, and dried under vacuum  
20 to give 9.8 g (45% yield) of 2-methyl-6-nitrophenylacetic acid as an off-white solid. The solid was hydrogenated in methanol over 10 % palladium on carbon to give 9.04 g of the title compound as a white solid.

**7-Bromo-5-chloro-2-oxindole**

25 5-Chloro-2-oxindole (16.8 g) and 19.6 g of N-bromosuccinimide were suspended in 140 mL of acetonitrile and refluxed for 3 hours. Thin layer chromatography (silica, ethyl acetate) at 2 hours of reflux showed 5-chloro-2-oxindole or N-bromosuccinimide (Rf 0.8), product (Rf 0.85) and a second  
30 product (Rf 0.9) whose proportions did not change after another hour of reflux. The mixture was cooled to 10 °C, the precipitate was collected by vacuum filtration, washed with 25 mL of ethanol and sucked dry for 20 minutes in the funnel to

WO 00/08202

103

PCT/US99/17845

give 14.1 g of wet product (56 % yield). The solid was suspended in 200 mL of denatured ethanol and slurry-washed by stirring and refluxing for 10 minutes. The mixture was cooled in an ice bath to 10 °C. The solid product was collected by vacuum filtration, washed with 25 mL of ethanol and dried under vacuum at 40 °C to give 12.7 g (51% yield) of 7-bromo-5-chloro-2-oxindole.

**5-Fluoro-2-oxindole**

5-Fluoroisatin (8.2 g) was dissolved in 50 mL of hydrazine hydrate and refluxed for 1.0 hr. The reaction mixtures were then poured in ice water. The precipitate was then filtered, washed with water and dried in a vacuum oven to afford the title compound.

**5-Nitro-2-oxindole**

2-Oxindole (6.5 g) was dissolved in 25 mL concentrated sulfuric acid and the mixture maintained at -10 to -15 °C while 2.1 mL of fuming nitric acid was added dropwise. After the addition of the nitric acid the reaction mixture was stirred at 0 °C for 0.5 hr and poured into ice-water. The precipitate was collected by filtration, washed with water and crystallized from 50% acetic acid. The crystalline product was then filtered, washed with water and dried under vacuum to give 6.3 g (70%) of 5-nitro-2-oxindole.

**5-Aminosulfonyl-2-oxindole**

To a 100 mL flask charged with 27 mL of chlorosulfonic acid was added slowly 13.3 g of 2-oxindole. The reaction temperature was maintained below 30 °C during the addition. After the addition, the reaction mixture was stirred at room temperature for 1.5 hr, heated to 68 °C for 1 hr, cooled, and poured into water. The precipitate was washed with water and dried in a vacuum oven to give 11.0 g of 5-chlorosulfonyl-2-oxindole (50% yield) which was used without further purification.

WO 00/08202

104

PCT/US99/17845

5-Chlorosulfonyl-2-oxindole (2.1 g) was added to 10 mL of ammonium hydroxide in 10 mL of ethanol and stirred at room temperature overnight. The mixture was concentrated and the solid collected by vacuum filtration to give 0.4 g (20% yield) of the title compound as an off-white solid.

**5-Methylaminosulfonyl-2-oxindole**

A suspension of 3.38 g of 5-chlorosulfonyl-2-oxindole in 10 mL 2M methylamine in tetrahydrofuran was stirred at room temperature for 4 hours during which time a white solid formed. The precipitate was collected by vacuum filtration, washed twice with 5 mL of water and dried under vacuum at 40 °C overnight to give 3.0 g (88 % yield) of 5-methylaminosulfonyl-2-oxindole.

**5-(4-Trifluoromethylphenylaminosulfonyl)-2-oxindole**

A suspension of 2.1 g of 5-chlorosulfonyl-2-oxindole, 1.6 g of 4-trifluoromethylaniline and 1.4 g of pyridine in 20 mL of dichloromethane was stirred at room temperature for 4 hours. The precipitate which formed was collected by vacuum filtration, washed twice with 5 mL of water and dried under vacuum at 40 °C overnight to give 2.4 g of crude product containing some impurities by thin layer chromatography. The crude product was chromatographed on silica gel eluting with ethyl acetate:hexane (1:2) to give 1.2 g (37 % yield) of 5-(4-trifluoromethylphenyl-aminosulfonyl)-2-oxindole.

**5-(Morpholinosulfonyl)-2-oxindole**

A suspension of 2.3 g of 5-chlorosulfonyl-2-oxindole and 2.2 g of morpholine in 50 mL of dichloromethane was stirred at room temperature for 3 hours. The white precipitate was collected by vacuum filtration, washed with ethyl acetate and hexane and dried under vacuum at 40 °C overnight to give 2.1 g (74 % yield) of 5-(morpholinosulfonyl)-2-oxindole.

**6-Trifluoromethyl-2-oxindole**

WO 00/08202

105

PCT/US99/17845

Dimethylsulfoxide (330 mL) was added to 7.9 g of sodium hydride followed by dropwise addition of 43.6 g diethyloxalate. The mixture was heated to 100 °C for 1.0 hour and cooled to room temperature. 2-Nitro-4-trifluoromethyltoluene (31.3 g) was added, the reaction stirred for 30 minutes at room temperature and then heated to 100 °C for 1 hour. The reaction was cooled and poured into a mixture of saturated aqueous ammonium chloride, ethyl acetate and hexane. The organic layer was washed with saturated ammonium chloride, water and brine, dried, and concentrated to give dimethyl 2-(2-nitro-4-trifluoromethylphenyl)malonate.

The diester was dissolved in a mixture of 6.4 g of lithium chloride and 2.7 mL of water in 100 mL of dimethylsulfoxide and heated to 100 °C for 3 hours. The reaction was cooled and poured into a mixture of ethyl acetate and brine. The organic phase was washed with brine, dried with sodium sulfate, concentrated and chromatographed on silica gel (10 % ethyl acetate in hexane). The fractions containing product were evaporated to give 25.7 g of methyl 2-nitro-4-trifluoromethylphenylacetate.

Methyl 2-nitro-4-trifluoromethylphenylacetate (26 mg) was hydrogenated over 10 % palladium on carbon and then heated at 100 °C for 3 hours. The catalyst was removed by filtration and the solvent evaporated to give the title compound.

#### 4-Carboxy-2-oxindole

A solution of trimethylsilyldiazomethane in hexane (2M) was added dropwise to a solution of 2.01 g of 2-chloro-3-carboxy-nitrobenzene in 20 mL methanol at room temperature until no further gas evolution occurred. The excess trimethylsilyldiazomethane was quenched with acetic acid. The reaction mixture was dried by rotary pump and the residue was further dried in a vacuum oven overnight. The product (2-chloro-3-methoxycarbonylnitrobenzene) was pure enough for the following reaction.



WO 00/08202

106

PCT/US99/17845 ,

Dimethyl malonate (6.0 mL) was added to an ice-cold suspension of 2.1 g of sodium hydride in 15 mL of DMSO. The reaction mixture was then stirred at 100 °C for 1.0 h and then cooled to room temperature. 2-Chloro-3-methoxycarbonyl-  
5 nitrobenzene (2.15 g) was added to the above mixture in one portion and the mixture was heated to 100 °C for 1.5 h. The reaction mixture was then cooled to room temperature and poured into ice water, acidified to pH 5, and extracted with ethyl acetate. The organic layer was washed with brine, dried  
10 over anhydrous sodium sulfate and concentrated to give 3.0 g of the dimethyl 2-methoxycarbonyl-6-nitrophenylmalonate.

Dimethyl 2-methoxycarbonyl-6-nitrophenylmalonate (3.0 g) was refluxed in 50 mL of 6 N hydrochloric acid overnight. The mixture was concentrated to dryness and refluxed for 2 hours  
15 with 1.1 g of tin(II) chloride in 20 mL of ethanol. The mixture was filtered through Celite, concentrated and chromatographed on silica gel (ethyl acetate:hexane:acetic acid) to give 0.65 g (37% yield) of 4-carboxy-2-oxindole as a white solid.

#### 20 5-Carboxyethyl-2-oxindole

5-Cyanoethyl-2-oxindole (4.02 g) in 10 mL of water containing 25 mL of concentrated hydrochloric acid was refluxed for 4 hours. The mixture was cooled, water added and the resulting solid collected by vacuum filtration, washed  
25 with water and dried to give 1.9 g (44% yield) of the title compound as a yellow solid.

#### 5-(Morpholin-4-ethyl)-2-oxindole

5-Chloroethyl-2-oxindole (2.3 g), 1.2 mL of morpholine and 1.2 mL of diisopropylethylamine were heated overnight at  
30 100 °C in 10 mL of dimethylsulfoxide. The mixture was cooled, poured into water and extracted with ethyl acetate. The organic layer was washed with brine, dried and evaporated. The residue

WO 00/08202

107

PCT/US99/17845

was chromatographed on silica gel (5 % methanol in chloroform) to give 0.9 g (31%) of the title compound as a white solid.

**5-(4-Methoxycarbonylbenzamido)-2-oxindole**

A mixture of 82.0 mg 5-amino-2-oxindole and 131.0 mg 4-methoxycarbonylbenzoyl chloride in pyridine was stirred at room temperature for 3 hr and poured into ice water. The precipitate was filtered, washed with water and dried in a vacuum oven to give 138.0 mg of 5-(4-methoxycarbonylbenzamido)-2-oxindole (81% yield).

**10 5-Methoxy-2-oxindole**

Chloral hydrate (9.6 g) was dissolved in 200 mL of water containing 83 g of sodium sulfate. The solution was warmed to 60 °C, a solution of 11.4 g of hydroxylamine hydrochloride in 50 mL of water was added and the mixture was held at 60 °C. In a separate flask, 6.4 g of 4-anisidine and 4.3 mL of concentrated hydrochloric acid in 80 mL of water was warmed to 80 °C. The first solution was added to the second and the mixture refluxed for 2 minutes after which it was cooled slowly to room temperature and then cooled in an ice bath. The tan precipitate was collected by vacuum filtration, washed with water and dried under vacuum to give 8.6 g (85% yield) of N-(2-hydroximino-acetyl)anisidine.

Concentrated sulfuric acid (45 mL) containing 5 mL of water was warmed to 60 °C and 8.6 g of N-(2-hydroximinoacetyl)anisidine was added in one portion. The stirred mixture was heated to 93 °C for 10 minutes and then allowed to cool to room temperature. The mixture was poured into 500 g of ice and extracted 3 times with ethyl acetate. The combined extracts were dried over anhydrous sodium sulfate and concentrated to give 5.1 g (65% yield) of 5-methoxyisatin as a dark red solid. 5-Methoxyisatin (5.0 g) and 30 mL of hydrazine hydrate were heated to reflux for 15 minutes. The reaction mixture was cooled to room temperature and 50 mL of

WO 00/08202

108

PCT/US99/17845

water was added. The mixture was extracted 3 times with 25 mL of ethyl acetate each time, the organic layers combined, dried over anhydrous sodium sulfate and concentrated to give a yellow solid. The solid was stirred in ethyl acetate and 1.1 g of insoluble material was removed by vacuum filtration and saved. This material proved to be 2-hydrazinocarbonylmethyl-4-anisidine. The filtrate was concentrated and chromatographed on silica gel eluting with ethyl acetate:hexane (1:1) to give 0.7 g of 5-methoxy-2-oxindole as a yellow solid. The 1.1 g of 2-hydrazino- carbonylmethyl-4-anisidine was refluxed for 1 hour in 20 mL of 1N sodium hydroxide. The mixture was cooled, acidified to pH 2 with concentrated hydrochloric acid and extracted 3 times with 25 mL of ethyl acetate each time. The organic extracts were combined, washed with brine, dried over anhydrous sodium sulfate and concentrated to give 0.8 g of 5-methoxy-2-oxindole as a yellow solid. The combined yield was 1.5 g or 33%.

#### 7-Azaoxindole

3,3-Dibromo-7-azaoxindole (2.9 g) was dissolved in a mixture of 20 mL of acetic acid and 30 mL of acetonitrile. To the solution was added 6.5 g of zinc dust. The mixture was stirred for 2 hrs at room temperature. The solid was filtered from the mixture and the solvent evaporated. The residue was slurried with ethyl acetate. The ethyl acetate solution containing insoluble solid was passed through a short column of silica gel. The collected ethyl acetate solution was evaporated and the residue dried under vacuum to give 1.8 g (yield 91%) of 7-azaoxindole acetic acid salt.

#### 5-Dimethylaminosulfonyl-2-oxindole

A suspension of 2.3 g 5-chlorosulfonyl-2-oxindole in 10 mL 2M dimethylamine in methanol was stirred at room temperature for 4 hours at which time a white solid formed. The precipitate was collected by vacuum filtration, washed with 5 mL of 1N sodium hydroxide and 5 mL of 1N hydrochloric

WO 00/08202

109

PCT/US99/17845 ,

acid and dried under vacuum at 40 °C overnight to give 1.9 g (79% yield) of 5-dimethylamino- sulfonyl-2-oxindole.

#### 6-Phenyl-2-oxindole

Dimethyl malonate (10 mL) in 25 mL of dimethylsulfoxide was added dropwise to 3.5 g sodium hydride suspended in 25 mL dimethylsulfoxide and the mixture heated at 100 °C for 10 minutes. The mixture was cooled to room temperature and 4.7 g of 4-fluoro-3-nitrobiphenyl in 25 mL dimethylsulfoxide was added. The mixture was heated at 100 °C for 2 hours, cooled and quenched with 300 mL of saturated ammonium chloride solution. The mixture was extracted three times with ethyl acetate and the combined organic layers washed with water and brine and evaporated to give, as a yellow oil, crude dimethyl-3-nitrobiphenyl- 4-malonate.

Crude dimethyl-3-nitrobiphenyl-4-malonate was refluxed in 30 mL of 6 N hydrochloric acid for 24 hours. The precipitate was collected by filtration, washed with water and dried to give 4.5 g of 3-nitrobiphenyl-4-acetic acid as a cream colored solid.

Iron powder (2.6 g) was added all at once to 4.5 g of 3-nitrobiphenyl-4-acetic acid in 40 mL of acetic acid. The mixture was refluxed for 2 hours, concentrated to dryness and taken up in ethyl acetate. The solids were removed by filtration and the filtrate washed twice with 1N hydrochloric acid and brine and dried over anhydrous sodium sulfate. The filtrate was concentrated to give 3.4 g (93% yield) of 6-phenyl-2-oxindole as a light brown solid.

#### 6-(3-Methoxyphenyl)-2-oxindole

Tetrakis(triphenylphosphine)palladium (0.8 g) was added to a mixture of 5 g 3-methoxyphenylboronic acid, 5 g 5-bromo-2-fluoro- nitrobenzene and 11 mL of 2 M sodium carbonate solution in 100 mL of toluene. The mixture was refluxed for 2 hours, diluted with water and extracted with ethyl acetate.

WO 00/08202

110

PCT/US99/17845

The ethyl acetate was washed with saturated sodium bicarbonate and brine and then dried and concentrated to give an oily solid. The solid was chromatographed on silica gel (ethyl acetate:hexane (1:6)) to give 4.3 g (77% yield) of 4-fluoro-3'-methoxy-3-nitrobiphenyl.

Dimethyl malonate (9.7 mL) was added dropwise to 2.0 g sodium hydride suspended in 50 mL dimethylsulfoxide. The mixture was heated to 100 °C for 35 minutes and cooled to room temperature. 4-Fluoro-2'-methoxy-3-nitrobiphenyl (4.2 g) in 50 mL dimethylsulfoxide was added and the mixture was heated at 100 °C for 1 hour. The reaction mixture was cooled and quenched with 300 mL of saturated ammonium chloride solution and extracted twice with ethyl acetate. The extracts were combined, washed with brine, dried over anhydrous sodium sulfate and concentrated to give crude dimethyl 3'-methoxy-3-nitrobiphenyl-4-malonate as a pale yellow solid.

Crude dimethyl 3'-methoxy-3-nitrobiphenyl-4-malonate was heated at 110 °C in 45 mL 6N hydrochloric acid for 4 days and then cooled. The precipitate was collected by filtration, washed with water and hexane, and dried to give 5.3 g of 3'-methoxy-2-nitrobiphenyl-4-acetic acid as a light tan solid.

3'-Methoxy-3-nitrobiphenyl-4-acetic acid (5.2 g) was dissolved in methanol and hydrogenated over 0.8 g of 10% palladium on carbon for 3 hours at room temperature. The catalyst was removed by filtration, washed with methanol and the filtrates combined and concentrated to give a brown solid. The solid was chromatographed on silica gel in ethyl acetate:hexane:acetic acid (33:66:1) to give 3.0 g of 6-(3-methoxyphenyl)-2-oxindole as a pink solid.

### 5-Acetyl-2-oxindole

2-Oxindole (3 g) was suspended in 1,2-dichloroethane and 3.2 mL acetyl chloride were slowly added. The resulting suspension was heated to 50 °C for 5 hours, cooled, and poured

WO 00/08202

111

PCT/US99/17845 ,

into water. The resulting precipitate was collected by vacuum filtration, washed copiously with water and dried under vacuum to give 2.9 g (73% yield) of the title compound as a brown solid.

5 **5-Cyanoethyl-2-oxindole**

Potassium cyanide (2.0 g) was added to 15 mL of dimethylsulfoxide and heated to 90 °C. 5-Chloroethyl-2-oxindole (3.0 g) dissolved in 5 mL dimethyl sulfoxide was added slowly with stirring, and the reaction heated to 150 °C for 2 hours. The mixture was cooled, poured into ice water and the precipitate collected by vacuum filtration, washed with water, dried and then chromatographed on silica gel (5% methanol in chloroform) to give 1.2 g (42% yield) of the title compound.

**6-(Morpholin-4-yl)-2-oxindole**

6-Amino-2-oxindole (2.2 g), 4.0 g 2, 2'-dibromoethyl ether and 7.9 g sodium carbonate were refluxed in 20 ml ethanol overnight, concentrated and diluted with 50 ml of water. The mixture was extracted three times with 50 ml of ethyl acetate and the organic extracts combined, washed with 20 ml of brine, dried over anhydrous sodium sulfate and concentrated to dryness. The solid was chromatographed on a column of silica gel (ethyl acetate:hexane (1:1) containing 0.7% acetic acid) to give 1.2 g (37% yield) of the title compound as a beige solid.

25 **Methylation of phenols**

An appropriate phenol (1 equivalent) is stirred overnight at room temperature with an equal volume of dimethylformamide containing 1.3 equivalents of methyl iodide and 1.3 equivalents of potassium carbonate. The mixture is diluted with water and extracted with ethyl acetate. The ethyl acetate layer is washed with water and brine. The organic layer is separated, dried over anhydrous sodium sulfate and evaporated to dryness to give the anisole derivative.

WO 00/08202

112

PCT/US99/17845

### Formylation of anisoles

The anisole (1 equivalent), dimethylformamide (1-3 equivalents) and phosphorus oxychloride (1-3 equivalents) are heated to 100° C for 1-3 hours. The reaction mixture is cooled to room temperature and dichloromethane is added. The mixture is cooled in ice and water is then added, followed by 4-12 equivalents of concentrated sodium hydroxide until a pH of 9-10 is reached. The organic layer is separated and washed with water and then with brine, dried over anhydrous sodium sulfate and evaporated to give the crude aldehyde. The crude aldehyde is dissolved in boiling hexane containing activated carbon and the hexane solution is decanted and filtered hot through a layer of silica gel. The filtrate is evaporated to dryness to give the benzaldehyde derivative.

### Formylation of pyrroles

POCl<sub>3</sub> (1.1 equiv.) is added dropwise to dimethylformamide (3 equiv.) in dichloromethane at -10° C followed by the appropriate pyrrole. After stirring for two hours, the reaction mixture is diluted with H<sub>2</sub>O and basified to pH 11 with 10 N KOH. The precipitate which forms is collected by filtration, washed with H<sub>2</sub>O and dried in a vacuum oven to give the desired aldehyde.

### Saponification of pyrrolecarboxylic acid esters

A mixture of a pyrrolecarboxylic acid ester and KOH (2 - 4 equiv.) in EtOH is refluxed until reaction completion is indicated by thin layer chromatography (TLC). The cooled reaction mixture is acidified to pH 3 with 1 N HCl. The precipitate which forms is collected by filtration, washed with H<sub>2</sub>O and dried in a vacuum oven to give the desired pyrrolecarboxylic acid.

### Amidation of pyrrolecarboxylic acids

To a stirred solution of a pyrrolecarboxylic acid dissolved in dimethylformamide (0.3M) is added 1-ethyl-3-(3-

WO 00/08202

113

PCT/US99/17845 ,

dimethylaminopropyl)carboiimide (1.2 equiv.), 1-hydroxybenzotriazole (1.2 equiv.), and triethylamine (2 equiv.). The appropriate amine is added (1 equiv.) and the reaction stirred until completion is indicated by TLC. The  
5 reaction is diluted with saturated NaHCO<sub>3</sub> and brine (with extra salt), dried over anhydrous MgSO<sub>4</sub> and concentrated to afford the desired amide.

Preparation of pyrrole and indole carbaldehydes

3-(2-Formyl-5-methyl-1H-pyrrol-3-yl)propionic acid

10 4-(2-Carboxyethyl)-3-ethoxycarbonyl-2-methylpyrrole (Bulter, A.R., and George, S.D. (1993) Tetrahedron 49: 7017-7026) was hydrolyzed using potassium hydroxide (KOH) in ethanol (EtOH) to give 4-(2-carboxyethyl)-2-methyl-1H-pyrrole-3-carboxylic acid (64%).

15 <sup>1</sup>HNMR (DMSO-d<sub>6</sub>) δ 11.63 (s, 1H, COOH), 10.81 (s, 1H, NH), 6.36 (s, 1H, CH), 2.78, 2.45 (2 x t, 4H, CH<sub>2</sub>CH<sub>2</sub>), 2.34 (s, 3H, CH<sub>3</sub>).

MS m/z 225 [M+2].

4-(2-Carboxyethyl)-2-methyl-1H-pyrrole-3-carboxylic acid  
20 was decarboxylated by heating for one hour at 200° C to give 3-(5-methyl-1H-pyrrol-3-yl)propionic acid.

<sup>1</sup>HNMR (DMSO-d<sub>6</sub>) δ 10.07 (s, 1H, NH), 9.60 (br s, 1H, COOH), 6.30 (s, 1H, CH), 5.56 (s, 1H, CH), 2.51, 2.36 (2 x t, 4H, CH<sub>2</sub>CH<sub>2</sub>), 2.10 (s, 3H, CH<sub>3</sub>).

25 MS m/z 154 [M+1].

3-(5-Methyl-1H-pyrrol-3-yl)propionic acid was formylated using phosphorus oxychloride (POCl<sub>3</sub>) and N,N-dimethylformamide (DMF) to give 3-(2-formyl-5-methyl-1H-pyrrol-3-yl)propionic acid.

30 <sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.09 (br s, 1H, COOH), 11.54 (br s, 1H, NH), 9.44 (s, 1H, CHO), 5.86 (d, J = 1.8 Hz, 1H), 2.87 (t, J = 7.5 Hz, 2H, CH<sub>2</sub>), 2.47 (t, J = 7.5 Hz, 2H, CH<sub>2</sub>), 2.16 (s, 3H, CH<sub>3</sub>).



WO 00/08202

114

PCT/US99/17845

MS-EI  $m/z$  181  $[M^+]$ .**4-(2-Carboxyethyl)-5-formyl-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester**

To a suspension of 10 g (38.9 mmol) ethyl 2,4-dimethyl-  
5 5-(ethoxycarbonyl)-3-pyrrolepropionate (commercially  
available) in a mixture of water (250 mL) and methanol  
(MeOH) (50 mL) was added 3.0 g of KOH in 50 mL of water. The  
mixture was stirred at 70°C for 4 hr. The reaction mixture  
was cooled and the methanol was stripped from the mixture  
10 under vacuum. The remaining solution was filtered and the  
filtrate was then acidified with 6N hydrochloric acid to pH  
3. The precipitate which formed was collected by filtration,  
washed with a 2:1 mixture of water and ethanol and dried  
under vacuum to give 7.5 g of 4-(2-carboxy-ethyl)-3,5-  
15 dimethyl-1H-pyrrole-2-carboxylic acid ethyl ester as a white  
solid.

$^1\text{H}$ NMR (360 MHz, DMSO- $d_6$ )  $\delta$  12.03 (br s, 1H, COOH), 11.02  
(br s, 1H, NH), 4.16 (q,  $J$  = 7.2 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 2.53 (t,  $J$  =  
7.8 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>), 2.25 (t,  $J$  = 7.8 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>), 2.16 (s,  
20 3H, CH<sub>3</sub>), 2.11 (s, 3H, CH<sub>3</sub>), 1.25 (t,  $J$  = 7.2 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI  $m/z$  239  $[M^+]$ .

To a mixture of 7 g 4-(2-carboxyethyl)-3,5-dimethyl-1H-  
pyrrole-2-carboxylic acid ethyl ester in 35 mL each of  
tetrahydrofuran (THF), acetic acid (AcOH) and water at -10°C  
25 was added 70 g of ceric ammonium nitrate in portion over 20  
minutes, the reaction temperature being maintained at about 5°  
C. The resulting mixture was then cooled to 0°C, stirred for  
another 2 hr and then diluted with 250 mL of brine. The  
mixture was then extracted with 2 x 300 mL 10% MeOH in  
30 dichloromethane (DCM). The organic extracts were combined and  
dried over anhydrous magnesium sulfate (MgSO<sub>4</sub>), filtered and  
concentrated. The residue was purified on silica gel column  
with hexanes(hex)/ethyl acetate (EtOAc)/AcOH (6:4:0.05)

WO 00/08202

115

PCT/US99/17845 ,

followed by crystallization in EtOAc to give 1.5 g of 4-(2-carboxyethyl)-5-formyl-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester as a white solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 12.36 (br s, 1H, COOH), 12.01  
5 (br s, 1H, NH), 9.77 (s, 1H, CHO), 4.27 (q, J = 7.1 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 2.89 (t, J = 7.6 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>), 2.37 (t, J = 7.6 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>), 2.21 (s, 3H, CH<sub>3</sub>), 1.30 (t, J = 7.1 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI m/z 253 [M<sup>+</sup>].

10 4-(3-Dimethylaminopropyl)-5-formyl-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester

To a suspension of 4-(2-carboxyethyl)-3-ethoxycarbonyl-2-methylpyrrole (2 g, 8.88 mmol) (Bulter, A.R., and George, S.D. (1993) Tetrahedron 49: 7017-7026) in 18 mL of DMF was added  
15 1.73 g (10.65 mmol) 1,1'-carbonyldiimidazole (CDI) followed by the dropwise addition of 8.9 mL (17.76 mmol) of 2M dimethylamine in THF. After stirring for 2 hr, the reaction was diluted with water (200 mL) and cooled. The precipitate which formed was collected by filtration, washed with water  
20 and dried to give 1.0 g of the product as a white crystalline solid. The filtrate was extracted with EtOAc, the organic layer was washed with brine, dried and concentrated to obtain an additional 0.9 g of the product for a total of 1.9 g of 4-(2-dimethylcarbamoyl-ethyl)-2-methyl-1H-pyrrole-3-carboxylic  
25 acid ethyl ester (85%).

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 10.89 (br s, 1H, NH), 6.40 (d, J = 2.5 Hz, 1H), 4.13 (q, J = 7.1 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 2.92 (s, 3H, NCH<sub>3</sub>), 2.79 (s, 3H, NCH<sub>3</sub>), 2.75 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 2.45 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>), 2.35 (s, 3H, CH<sub>3</sub>), 1.23 (t, J = 7.1 Hz, 3H,  
30 OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI m/z 252 [M<sup>+</sup>].

To a heterogeneous mixture of 4-(2-dimethylcarbamoyl-ethyl)-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester (1.01

WO 00/08202

116

PCT/US99/17845 ,

g, 4 mmol) in THF (9 mL) was added dropwise 8 mL of borane-tetrahydrofuran complex (1M in THF). The mixture was refluxed overnight. Then 9.0 mL of MeOH was added slowly to the reaction and refluxing was continued for another 2 hr. The  
5 reaction mixture was cooled, quenched with 1N HCl and extracted with EtOAc. The aqueous layer was basified with aqueous KOH and extracted with EtOAc. The EtOAc extract was washed with brine, dried and concentrated to give 616 mg (65%) of 4-(3-dimethylaminopropyl)-2-methyl-1H-pyrrole-3-carboxylic  
10 acid ethyl ester as a faint orange oil.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 10.84 (br s, 1H, NH), 6.36 (d, J = 2.5 Hz, 1H), 4.12 (q, J = 7.1 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 2.52 (m, 2H, CH<sub>2</sub>), 2.34 (s, 3H, CH<sub>3</sub>), 2.17 (m, 2H, CH<sub>2</sub>), 2.08 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.57 (m, 2H, CH<sub>2</sub>CH<sub>2</sub> CH<sub>2</sub>), 1.23 (t, J = 7.1 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).  
15

MS-EI m/z 238 [M<sup>+</sup>].

4-(3-Dimethylaminopropyl)-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester (600 mg, 2.5 mmol) was formylated using POCl<sub>3</sub> and DMF to give 645 mg (96%) of 4-(3-dimethylamino-  
20 propyl)-5-formyl-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 12.15 (br s, 1H, NH), 9.59 (s, 1H, CHO), 4.19 (q, J = 7.2 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 2.93 (m, 2H, CH<sub>2</sub>), 2.41 (s, 3H, CH<sub>3</sub>), 2.18 (m, 2H, CH<sub>2</sub>), 2.09 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>),  
25 1.63 (m, 2H, CH<sub>2</sub>CH<sub>2</sub> CH<sub>2</sub>), 1.27 (t, J = 7.2 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI m/z 266 [M<sup>+</sup>].

### 3-(3-Dimethylaminopropyl)-1H-indole-2-carbaldehyde

Ethyl indole-2-carboxylate (10 g, 52.8 mmol) was formylated using POCl<sub>3</sub> (1.3 equiv.) and DMF (1.3 equiv.) as  
30 above to give 3-formyl-1H-indole-2-carboxylic acid ethyl ester as a white solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 12.0 (br s, 1H, NH), 10.6 (s, 1H, CHO), 8.24 (dd, J = 0.7 & 8.0 Hz, 1H), 7.57 (d, 1H), 7.38

WO 00/08202

117

PCT/US99/17845

(m, 1H), 7.26 (m, 1H), 4.44 (q,  $J = 7.12$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 1.39 (t,  $J = 7.12$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  217 [ $\text{M}^+$ ].

Butyllithium (1.1 equiv.) was added to a suspension of  
5 (2-dimethylaminoethyl)triphenylphosphonium bromide in THF (0.2 M) at 0°C. After stirring for 30 minutes, lithium diethylamine (1.1 equiv.) was added dropwise followed by the cold suspension of 3-formyl-1H-indole-2-carboxylic acid ethyl ester (3.96 g, 18.2 mmol) in THF. The resulting orange suspension  
10 was stirred for 18 hours. The reaction was then quenched with saturated ammonium chloride and extracted with 10% MeOH in DCM. The organic layer was washed with saturated sodium bicarbonate, dried and concentrated to give 3-(3-dimethylaminopropenyl)-1H-indole-2-carboxylic acid ethyl ester  
15 as a brownish-red waxy solid.

$^1\text{H}$ NMR (360 MHz,  $\text{DMSO}-d_6$ )  $\delta$  11.8 (br s, 1H, NH), 7.93 (d, 1H), 7.49 (d, 1H), 7.41 (d, 1H), 7.3 (t, 1H), 7.14 (t, 1H), 6.33-6.42 (m, 1H), 4.36 (q, 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.38 (d, 2H,  $\text{CH}_2$ ), 2.44 (s, 6H,  $2 \times \text{CH}_3$ ), 1.36 (t, 3H,  $\text{OCH}_2\text{CH}_3$ ).

20 MS-EI  $m/z$  272 [ $\text{M}^+$ ].

3-(3-Dimethylaminopropenyl)-1H-indole-2-carboxylic acid ethyl ester was hydrogenated using 10% palladium over carbon to reduce the double bond followed by a lithium aluminum hydride (LAH) reduction of the ester group to an alcohol  
25 resulting in the formation of [3-(3-dimethylaminopropyl)-1H-indol-2-yl]methanol as an orange oil.

Manganese dioxide (10 equiv.) was added to a solution of [3-(3-dimethylaminopropyl)-1H-indol-2-yl]methanol (2.2 g) in DCM. The mixture was stirred at room temperature for 18 hr.  
30 Insoluble materials were removed by filtration and the filtrate was concentrated. The residue was column chromatographed to give 3-(3-dimethylaminopropyl)-1H-indole-2-carbaldehyde as a cream-colored solid.

WO 00/08202

118

PCT/US99/17845 ,

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 11.58 (br s, 1H, NH), 10.0 (s, 1H, CHO), 7.72 (d, 1H), 7.39 (d, 1H), 7.29 (t, 1H), 7.07 (t, 1H), 3.08 (t, 2H, CH<sub>2</sub>), 2.20 (t, 2H), 2.12 (s, 6H, 2xCH<sub>3</sub>), 1.73 (t, 2H, CH<sub>2</sub>).

5 MS-EI m/z 230 [M<sup>+</sup>].

**Preparation of aminotetrahydroindolecarbaldehydes**

A mixture of 5-aminolevulinic acid hydrochloride (1 equiv.), 1,3-cyclohexanedione (1 equiv.) and sodium acetate (2 equiv.) in water (1M) is stirred at 110° C for 4 - 12 hr and  
10 then cooled. The precipitate which forms is collected by vacuum filtration, washed with 30% ethanol in water and dried under vacuum to give the amido-keto-tetrahydroindole in 50-70% yield.

Lithium aluminum hydride (LAH, 4 equiv.) is added dropwise  
15 to a suspension of the appropriate amido-keto-tetrahydroindole (1 equiv.) in THF (0.5 M). The mixture is refluxed overnight. The mixture is then cooled and water is added carefully until no more gas is generated, then a few drops of 15% NaOH/water is added. The mixture is then stirred at room temperature for  
20 0.5 hr and filtered to remove insoluble materials. The filtrate is concentrated to give the amino-tetrahydroindole.

**Examples of syntheses of  
aminotetrahydroindolecarbaldehydes**

The following syntheses of aminotetrahydroindole-  
25 carbaldehydes are shown by way of example only and are not to be construed as limiting the scope of this invention in any manner whatsoever.

**3-[3-(4-methylpiperazin-1-yl)propyl]-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde**

30 **Step 1:** A mixture of 5-aminolevulinic acid hydrochloride (1.68 g, 10 mmol), 1,3-cyclohexanedione (1.12 g, 10 mmol) and sodium acetate (1.64 g, 20 mmol) in water (10 mL) was stirred at 110° C for 4 hr and then cooled. The precipitate which

WO 00/08202

119

PCT/US99/17845 ,

formed was collected by vacuum filtration, washed with 30% of ethanol in water and dried under vacuum to give 1.7 g (82%) of 3-(4-oxo-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 11.91 (br s, 1H, COOH), 10.99 (br s, 1H, NH), 6.45 (d, J = 1.4 Hz, 1H), 2.76 (t, 2H, CH<sub>2</sub>), 2.69 (t, 2H, CH<sub>2</sub>), 2.44 (t, 2H, CH<sub>2</sub>), 2.26 (t, 2H, CH<sub>2</sub>), 1.96 (m, 2H, CH<sub>2</sub>).

MS-EI m/z 207 [M<sup>+</sup>].

**Step 2:** To a suspension of 10 g of the product of step 1 (48 mmol) in dichloromethane (60 mL) was added 9.3 g (57.6 mmol) of 1,1'-carbonyldiimidazole. The mixture was stirred at room temperature for 2 hours and then 5.3 mL (48 mmol) 1-methylpiperazine and 8.4 mL (48 mmol) N,N-diisopropylethylamine was added. The dark red reaction mixture was then stirred at room temperature overnight. The reaction was then poured into water, the organic layer separated and washed with brine, dried and concentrated to give 8 g (57%) of 3-[3-(4-methylpiperazin-1-yl)-3-oxo-propyl]-1,5,6,7-tetrahydroindol-4-one.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 10.97 (br s, 1H, NH), 6.47 (d, J = 2.0 Hz, 1H), 3.43 (m, 4H, 2xCH<sub>2</sub>), 2.67-2.75 (m, 4H, 2xCH<sub>2</sub>), 2.51 (m, 2H, CH<sub>2</sub>), 2.27 (m, 2H, CH<sub>2</sub>), 2.20 (m, 4H, 2xCH<sub>2</sub>), 2.15 (s, 3H, CH<sub>3</sub>), 1.97 (m, 2H, CH<sub>2</sub>).

MS-EI m/z 289 [M<sup>+</sup>].

**Step 3:** LAH (2.6 g, 68 mmol) was added dropwise to a suspension of 3-[3-(4-methylpiperazin-1-yl)-3-oxo-propyl]-1,5,6,7-tetrahydroindol-4-one (5g, 17 mmol) in THF (300 mL). The mixture was then refluxed overnight. The mixture was then cooled and 2.6 mL each of water followed by a few drops of 15% NaOH was added. The reaction was stirred at room temperature for 30 min and then filtered to remove insolubles. The filtrate was concentrated to give 4.5 g (100%) of 3-[3-(4-methylpiperazin-1-yl)propyl]-4,5,6,7-tetrahydro-1H-indole.

WO 00/08202

120

PCT/US99/17845

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 9.79 (br s, 1H, NH), 6.22 (d, J = 2.0 Hz, 1H), 2.44 (m, 2H, CH<sub>2</sub>), 2.21-2.3 (m, 14H, 7xCH<sub>2</sub>), 2.12 (s, 3H, CH<sub>3</sub>), 1.65 (m, 4H, 2xCH<sub>2</sub>), 1.53 (m, 2H, CH<sub>2</sub>).

MS-EI m/z 261 [M<sup>+</sup>].

- 5       **Step 4:** POCl<sub>3</sub> (1.8 mL, 18.9 mmol) was added dropwise to N,N-dimethylformamide (DMF, 3.8 mL, 51.6 mmol) at -5° C. The mixture was then allowed to come to room temperature and then stirred for 30 minutes after which it was again cooled to -5°C.
- 10       A solution of 3-[3-(4-methylpiperazin-1-yl)-propyl]-4,5,6,7-tetrahydro-1H-indole (4.5 g, 17.2 mmol) in DMF (9 mL) was added dropwise. The mixture was again allowed to come to room temperature and then stirred at that temperature overnight. The reaction was then quenched with ice, followed by 10 N NaOH to adjust the pH to 10-11. After stirring at room temperature
- 15       for 1 hr, the reaction was extracted with ethyl acetate (EtOAc), the organic layer separated, washed with brine, dried and concentrated to give 3.1 g (62%) of 3-[3-(4-methylpiperazin-1-yl)-propyl]-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde.

- 20       <sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 11.24 (br, s, 1H, NH), 9.42 (s, 1H, CHO), 2.60 (t, 2H, CH<sub>2</sub>), 2.51 (m, 2H, CH<sub>2</sub>), 2.35 (m, 2H, CH<sub>2</sub>), 2.28 (m, 8H, 4xCH<sub>2</sub>), 2.21 (m, 2H, CH<sub>2</sub>), 2.12 (s, 3H, CH<sub>3</sub>), 1.57-1.68 (m, 6H, 3xCH<sub>2</sub>).

MS-EI m/z 289 [M<sup>+</sup>].

- 25       **3-(3-dimethylaminopropyl)-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde**

The procedure was the same as that employed in Example A except that the amine in Step 2 was dimethylamine (2.0M solution in tetrahydrofuran).

- 30       **Step 2: N,N-Dimethyl-3-(4-oxo-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionamide:**

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 11.0 (br s, 1H, NH), 6.48 (d, J = 1.6 Hz, 1H), 2.95 (s, 3H, CH<sub>3</sub>), 2.79 (s, 3H, CH<sub>3</sub>), 2.71 (m,

WO 00/08202

121

PCT/US99/17845 ,

4H, 2xCH<sub>2</sub>), 2.47 (m, 2H, CH<sub>2</sub>), 2.27 (t, 2H, CH<sub>2</sub>), 1.96 (m, 2H, CH<sub>2</sub>).

MS-EI m/z 234 [M<sup>+</sup>].

5 Step 3: Dimethyl-[3-(4,5,6,7-tetrahydro-1H-indol-3-yl)propyl]-amine:

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.83 (br s, 1H, NH), 6.22 (d, J = 2.3 Hz, 1H), 2.43 (m, 2H, CH<sub>2</sub>), 2.28 (m, 2H, CH<sub>2</sub>), 2.14-2.25 (m, 4H, CH<sub>2</sub>), 2.08 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.64 (m, 4H, 2xCH<sub>2</sub>), 1.53 (m, 2H, CH<sub>2</sub>).

10 MS-EI m/z 206 [M<sup>+</sup>].

Step 4: 3-(3-Dimethylaminopropyl)-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde:

15 <sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 11.29 (br s, 1H, NH), 9.40 (s, 1H, CHO), 2.59 (t, 2H, CH<sub>2</sub>), 2.5 (m, 2H, CH<sub>2</sub>), 2.34 (m, 2H, CH<sub>2</sub>), 2.16 (m, 2H, CH<sub>2</sub>), 2.08 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.67 (m, 4H, 2xCH<sub>2</sub>), 1.56 (m, 2H, CH<sub>2</sub>).

MS-EI m/z 234 [M<sup>+</sup>].

20 3-(3-pyrrolidin-1-ylpropyl)-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde

The procedure employed was that same as that in Example A except that the amine used in Step 2 was pyrrolidine.

Step 2: 3-(3-Oxo-3-pyrrolidin-1-ylpropyl)-1,5,6,7-tetrahydroindol-4-one:

25 <sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 11.05 (br s, 1H, NH), 6.46 (d, J = 1.5 Hz, 1H), 3.35 (m, 2H, CH<sub>2</sub>), 2.24 (m, 2H, CH<sub>2</sub>), 2.66-2.73 (m, 4H, 2xCH<sub>2</sub>), 2.44 (m, 2H, CH<sub>2</sub>), 2.26 (m, 2H, CH<sub>2</sub>), 1.96 (m, 2H, CH<sub>2</sub>), 1.82 (m, 2H, CH<sub>2</sub>), 1.73 (m, 2H, CH<sub>2</sub>).

MS-EI m/z 260 [M<sup>+</sup>].

30 Step 3: 3-(3-Pyrrolidin-1-ylpropyl)-4,5,6,7-tetrahydro-1H-indole:

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.82 (br s, 1H, NH), 6.22 (s, 1H), 2.2-2.5 (m, 12H, 6xCH<sub>2</sub>), 1.5-1.64 (m, 10H, 5xCH<sub>2</sub>).



WO 00/08202

122

PCT/US99/17845

MS-EI  $m/z$  232  $[M^+]$ .

Step 4: 3-(3-Pyrrolidin-1-ylpropyl)-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde:

$^1\text{H}$ NMR (300 MHz, DMSO- $d_6$ )  $\delta$  11.28 (br, s, 1H, NH), 9.38 (s, 1H, CHO), 2.59 (t, 2H,  $\text{CH}_2$ ), 2.46 (m, 2H,  $\text{CH}_2$ ), 2.3-2.44 (m, 8H,  $4\times\text{CH}_2$ ), 1.55-1.65 (m, 10H,  $5\times\text{CH}_2$ ).

MS-EI  $m/z$  260  $[M^+]$ .

3-(3-diethylamino-propyl)-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde

10 The procedure used was the same as that in Example A except that the amine used was diethylamine.

Step 2: N,N-Diethyl-3-(4-oxo-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionamide:

$^1\text{H}$ NMR (300 MHz, DMSO- $d_6$ )  $\delta$  11.0 (br s, 1H, NH), 6.47 (d,  $J$  = 2.0 Hz, 1H), 3.24 (m, 4H,  $\text{N}(\text{CH}_2\text{CH}_3)_2$ ), 2.69 (m, 4H,  $2\times\text{CH}_2$ ), 2.46 (m, 2H,  $\text{CH}_2$ ), 2.27 (m, 2H,  $\text{CH}_2$ ), 1.96 (m, 2H,  $\text{CH}_2$ ), 1.04 (t,  $J$  = 7.0 Hz, 3H,  $\text{NCH}_2\text{CH}_3$ ), 0.98 (t,  $J$  = 7.0 Hz, 3H,  $\text{NCH}_2\text{CH}_3$ ).

Step 3: Diethyl-[3-(4,5,6,7-tetrahydro-1H-indol-3-yl)-propyl]-amine:

20  $^1\text{H}$ NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.83 (br s, 1H, NH), 6.22 (d,  $J$  = 2.0 Hz, 1H), 2.19-2.44 (m, 12H,  $6\times\text{CH}_2$ ), 1.64 (m, 4H,  $2\times\text{CH}_2$ ), 1.51 (m, 2H,  $\text{CH}_2$ ), 0.90 (t,  $J$  = 7.0 Hz, 6H,  $\text{N}(\text{CH}_2\text{CH}_3)_2$ ).

MS-EI  $m/z$  234  $[M^+]$ .

25 Step 4: 3-(3-Diethylaminopropyl)-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde:

$^1\text{H}$ NMR (360 MHz, DMSO- $d_6$ )  $\delta$  11.24 (br, s, 1H, NH), 9.42 (s, 1H, CHO), 2.59 (t, 2H,  $\text{CH}_2$ ), 2.34-2.53 (m, 10H,  $5\times\text{CH}_2$ ), 1.67 (m, 4H,  $2\times\text{CH}_2$ ), 1.57 (m, 2H,  $\text{CH}_2$ ), 0.91 (t,  $J$  = 7.0 Hz, 6H,  $\text{N}(\text{CH}_2\text{CH}_3)_2$ ).

30 MS-EI  $m/z$  262  $[M^+]$ .

3-(3-Diethylaminopropyl)-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde

Step 1: A mixture of 5-aminolevulinic acid hydrochloride

WO 00/08202

123

PCT/US99/17845

(1.68 g, 10 mmol) , 5,5-dimethyl-1,3-cyclohexandione (1.4 g, 10 mmol) and sodium acetate (1.64 g, 20 mmol) in water (10 mL) was stirred at 110°C for 4 hr and then cooled. The precipitate  
5 which formed was collected by vacuum filtration, washed with 30% of ethanol (EtOH) in water and dried under vacuum to give 1.6 g (68%) of 3-(4-oxo-6-dimethyl-4,5,6,7-tetrahydro-1H-indol-3-yl)propionic acid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 11.89 (br s, 1H, COOH), 10.94  
10 (br s, 1H, NH), 6.45 (d, J = 1.4 Hz, 1H), 2.76 (t, 2H, CH<sub>2</sub>), 2.57 (s, 2H, CH<sub>2</sub>), 2.44 (t, 2H, CH<sub>2</sub>), 2.16 (s, 2H, CH<sub>2</sub>), 0.99 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 235 [M<sup>+</sup>].

**Step 2:** To a suspension of 1.18 g (5 mmol) of the product  
15 of step 1 in dichloromethane (25 mL) was added 0.97 g (6 mmol) of CDI. After stirring at room temperature for 2 hr, 2.1 mL (20 mmol) diethylamine was added. The mixture was stirred at room temperature overnight. The reaction was concentrate and the residue was dissolved in dichloromethane, washed with  
20 brine, dried and concentrated to give 1.2 g (83%) of 3-(6,6-dimethyl-4-oxo-4,5,6,7-tetrahydro-1H-indol-3-yl)-N,N-diethylpropionamide as a white solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 10.91 (br s, 1H, NH), 6.46 (s, 1H), 3.20-3.29 (m, 4H, 2xCH<sub>2</sub>), 2.72-2.76 (m, 2H, CH<sub>2</sub>), 2.57 (s, 2H, CH<sub>2</sub>), 2.45 (m, 2H, CH<sub>2</sub>), 2.17 (s, 2H, CH<sub>2</sub>), 0.96-1.06 (m, 12H, 4xCH<sub>3</sub>).

MS-EI m/z 290 [M<sup>+</sup>].

**Step 3:** LAH (0.57 g, 15.1 mmol) was added dropwise to a suspension of 3-(6,6-dimethyl-4-oxo-4,5,6,7-tetrahydro-1H-indol-3-yl)-N,N-diethylpropionamide (1.1 g, 3.8 mmol) in THF  
30 (80 mL). The mixture was refluxed overnight. The reaction was cooled and ice was added until no more gas was generated. A few drops of 15% NaOH in water was then added. The mixture was

WO 00/08202

124

PCT/US99/17845 ,

stirred at room temperature for 30 minutes and then filtered to remove insolubles. The filtrate was concentrated to give 0.9 g of [3-(6,6-dimethyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propyl]-diethyl-amine as a light yellow oil.

5       <sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 9.75 (br s, 1H, NH), 6.24 (s, 1H), 2.19-2.44 (m, 14H, 7xCH<sub>2</sub>), 1.53 (m, 2H, CH<sub>2</sub>), 1.40 (m, 2H, CH<sub>2</sub>), 0.88-0.92 (m, 12H, 4xCH<sub>3</sub>).

MS-EI m/z 262 [M<sup>+</sup>].

Step 4: POCl<sub>3</sub> (0.35 mL, 3.74 mmol) was added dropwise to  
10 DMF (0.8 mL, 10.3 mmol) at -5° C. After stirring at room temperature for 30 min, the mixture was cooled to -5°C. A solution of [3-(6,6-dimethyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propyl]-diethylamine (0.9 g, 3.4 mmol) in DMF (2 mL) was then added dropwise. The mixture was stirred at room  
15 temperature for 3 hours and then quenched with ice, followed by 10 N KOH to adjust pH to 10-11. After stirring at room temperature for 1 hour, the mixture was extracted with EtOAc, washed with brine, dried and concentrated to give 0.55 g 3-(3-diethylaminopropyl)-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-  
20 2-carbaldehyde.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 11.23 (br, s, 1H, NH), 9.41 (s, 1H, CHO), 2.61 (t, 2H, CH<sub>2</sub>), 2.30-2.43 (m, 10 H, 5x CH<sub>2</sub>), 1.58 (m, 2H, CH<sub>2</sub>), 1.45 (t, 2H, CH<sub>2</sub>), 0.93 (s, 6H, 2xCH<sub>3</sub>), 0.89 (t, 6H, N(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>).

25       MS-EI m/z 290 [M<sup>+</sup>].

**6,6-Dimethyl-3-(3-pyrrolidin-1-ylpropyl)-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde**

The procedure employed was the same as that in Example E except that the amine in Step 2 was pyrrolidine.

30       **Step 2: 6,6-Dimethyl-3-(3-oxo-3-pyrrolidin-1-ylpropyl)-1,5,6,7-tetrahydroindol-4-one**

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 10.90 (br s, 1H, NH), 6.46 (s, 1H), 3.34 (t, 2H, CH<sub>2</sub>), 3.25 (t, 2H, CH<sub>2</sub>), 2.76 (m, 2H, CH<sub>2</sub>),

WO 00/08202

125

PCT/US99/17845

2.58 (s, 2H, CH<sub>2</sub>), 2.43 (m, 2H, CH<sub>2</sub>), 2.17 (s, 2H, CH<sub>2</sub>), 1.74-1.74 (m, 4H, 2xCH<sub>2</sub>), 1.00 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 288 [M<sup>+</sup>].

5 Step 3: 6,6-Dimethyl-3-(3-pyrrolidin-1-ylpropyl)-4,5,6,7-tetrahydro-1H-indole

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 9.75 (br s, 1H, NH), 6.23 (d, 1H), 2.22-2.37 (m, 12H, 6xCH<sub>2</sub>), 1.55-1.66 (m, 6H, 3xCH<sub>2</sub>), 1.40 (m, 2H, CH<sub>2</sub>), 0.92 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 260 [M<sup>+</sup>].

10 Step 4: 6,6-Dimethyl-3-(3-pyrrolidin-1-ylpropyl)-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 11.25 (br, s, 1H, NH), 9.41 (s, 1H, CHO), 2.64 (t, 2H, CH<sub>2</sub>), 2.31-2.38 (m, 10 H, 5x CH<sub>2</sub>), 1.59-1.67 (m, 6H, 3xCH<sub>2</sub>), 1.46 (t, 2H, CH<sub>2</sub>), 0.93 (s, 6H, 2xCH<sub>3</sub>).

15 MS-EI m/z 288 [M<sup>+</sup>].

6,6-Dimethyl-3-[3-(4-methylpiperazin-1-yl)-propyl]-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde

The procedure used was that same as that in Example E except that the amine in Step 2 was 1-methylpiperazine.

20 Step 2: 6,6-Dimethyl-3-[3-(4-methylpiperazin-1-yl)-3-oxopropyl]-1,5,6,7-tetrahydroindol-4-one

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 10.93 (br s, 1H, NH), 6.48 (s, 1H), 3.42 (m, 4H, 2xCH<sub>2</sub>), 3.73 (m, 2H, CH<sub>2</sub>), 2.57 (s, 2H, CH<sub>2</sub>), 2.48 (m, 2H, CH<sub>2</sub>), 2.21 (m, 4H, 2xCH<sub>2</sub>), 2.17 (s, 2H, CH<sub>2</sub>), 2.15 (s, 3H, CH<sub>3</sub>), 1.00 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 317 [M<sup>+</sup>].

Step 3: 6,6-Dimethyl-3-[3-(4-methylpiperazin-1-yl)-propyl]-4,5,6,7-tetrahydro-1H-indole

30 <sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 9.74 (br s, 1H, NH), 6.24 (s, 1H), 2.21-2.30 (m, 16H, 8xCH<sub>2</sub>), 2.12 (s, 3H, CH<sub>3</sub>), 1.56 (m, 2H, CH<sub>2</sub>), 1.40 (m, 2H, CH<sub>2</sub>), 0.92 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 289 [M<sup>+</sup>].

Step 4: 6,6-Dimethyl-3-[3-(4-methylpiperazin-1-yl)-

WO 00/08202

126

PCT/US99/17845

propyl]-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 11.21 (br, s, 1H, NH), 9.42 (s, 1H, CHO), 2.62 (t, 2H, CH<sub>2</sub>), 2.18-2.37 (m, 14 H, 7x CH<sub>2</sub>), 2.12 (s, 3H, CH<sub>3</sub>), 1.61 (m, 2H, CH<sub>2</sub>), 1.45 (t, 2H, CH<sub>2</sub>), 0.93 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 317 [M<sup>+</sup>].

6,6-Dimethyl-3-(3-morpholin-4-ylpropyl)-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde

The procedure used was the same as that in Example E except that the amine used was morpholine.

Step 2: 6,6-Dimethyl-3-(3-morpholin-4-yl-3-oxopropyl)-1,5,6,7-tetrahydroindol-4-one

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 10.94 (br s, 1H, NH), 6.49 (s, 1H), 3.42-3.50 (m, 8H, 4xCH<sub>2</sub>), 2.74 (m, 2H, CH<sub>2</sub>), 2.57 (s, 2H, CH<sub>2</sub>), 2.48 (m, 2H, CH<sub>2</sub>), 2.17 (s, 2H, CH<sub>2</sub>), 1.00 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 304 [M<sup>+</sup>].

Step 3: 6,6-Dimethyl-3-(3-morpholin-4-ylpropyl)-4,5,6,7-tetrahydro-1H-indole

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 9.75 (br s, 1H, NH), 6.24 (s, 1H), 3.54 (m, 4H, 2xCH<sub>2</sub>), 2.23-2.31 (m, 12H, 6xCH<sub>2</sub>), 1.58 (m, 2H, CH<sub>2</sub>), 1.40 (m, 2H, CH<sub>2</sub>), 0.92 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 276 [M<sup>+</sup>].

Step 4: 6,6-Dimethyl-3-(3-morpholin-4-ylpropyl)-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 11.25 (br, s, 1H, NH), 9.43 (s, 1H, CHO), 3.54 (m, 4H, 2xCH<sub>2</sub>), 2.63 (t, 2H, CH<sub>2</sub>), 2.20-2.37 (m, 10 H, 5xCH<sub>2</sub>), 1.62 (m, 2H, CH<sub>2</sub>), 1.45 (t, 2H, CH<sub>2</sub>), 0.93 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 304 [M<sup>+</sup>].

3-(3-Dimethylaminopropyl)-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde

The procedure was the same as that in Example E except that the amine used was dimethylamine.

WO 00/08202

127

PCT/US99/17845

**Step 2: 3-(6,6-Dimethyl-4-oxo-4,5,6,7-tetrahydro-1H-indol-3-yl)-N,N-dimethylpropionamide**

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 10.93 (br s, 1H, NH), 6.48 (s, 1H), 2.94 (s, 3H, CH<sub>3</sub>), 2.79 (s, 3H, CH<sub>3</sub>), 2.72 (m, 2H, CH<sub>2</sub>), 2.57 (s, 2H, CH<sub>2</sub>), 2.48 (m, 2H, CH<sub>2</sub>), 2.17 (s, 2H, CH<sub>2</sub>), 1.00 (m, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 262 [M<sup>+</sup>].

**Step 3: [3-(6,6-Dimethyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propyl]dimethylamine**

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 9.75 (br s, 1H, NH), 6.24 (s, 1H), 3.36 (m, 2H, CH<sub>2</sub>), 2.07-2.3 (m, 14H, 7xCH<sub>2</sub>), 1.53 (m, 2H, CH<sub>2</sub>), 1.40 (m, 2H, CH<sub>2</sub>), 0.92 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 234 [M<sup>+</sup>].

**Step 4: 3-(3-Dimethylaminopropyl)-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-2-carbaldehyde**

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 11.25 (br s, 1H, NH), 9.41 (s, 1H, CHO), 2.62 (t, 2H, CH<sub>2</sub>), 2.36 (t, 2H, CH<sub>2</sub>), 2.30 (s, 2H, CH<sub>2</sub>), 2.21 (t, 2H, CH<sub>2</sub>), 2.12 (s, 6H, 2xCH<sub>3</sub>), 1.60 (m, 2H, CH<sub>2</sub>), 1.46 (t, 2H, CH<sub>2</sub>), 0.93 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 262 [M<sup>+</sup>].

**Condensation of aldehydes and oxindoles containing carboxylic acid substituents**

A mixture of the appropriate oxindole (1 equivalent), 1 equivalent of the appropriate aldehyde and 1 - 3 equivalents of piperidine (or pyrrolidine) in ethanol (0.4 M) is stirred at 90-100°C until the reaction complete as indicated by thin layer chromatography. The mixture is then concentrated and the residue is triturated with dilute hydrochloric acid. The resulting precipitate is collected by vacuum filtration, washed with water and ethanol and dried to give the product.

**Condensation of aldehyde and oxindole not containing carboxylic acid substituents**

WO 00/08202

128

PCT/US99/17845

A mixture of the appropriate oxindole (1 equivalent), 1 equivalent of the appropriate aldehyde and 1 - 3 equivalents of piperidine (or pyrrolidine) in ethanol (0.4 M) is stirred at 90-100° C until the reaction complete as indicated by thin layer chromatography. The mixture is cooled to room temperature and the precipitate which forms is collected by vacuum filtration, washed with ethanol and dried to give the product. Where a precipitate does not form, the mixture is concentrated and the product isolated by column chromatography.

**Condensation of aldehyde and oxindole containing amino substituents**

The aminooxindole is first protected with a BOC-group. After the condensation with the appropriate aldehyde, the solid which forms is deprotected using trifluoroacetic acid to yield the desired product.

**C. Examples - Synthesis of 3-methylidenyl-2-indolinones of this invention.**

The following syntheses of representative compounds of this invention are shown by way of example only and are not to be construed as limiting the scope of this invention in any manner whatsoever.

**Example 1: 3-(3,5-Diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one**

3,5-Diisopropyl-4-hydroxybenzaldehyde was methylated to give 3,5-diisopropyl-4-methoxybenzaldehyde.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.1 (s, 1H, CHO), 6.9 (s, 2H, aromatic), 3.9 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 2H, 2xCH), 1.2 (d, 12H, 4xCH<sub>3</sub>).

3,5-Diisopropyl-4-methoxybenzaldehyde was then condensed with 2-oxindole to give 0.25 g of 3-(3,5-Diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

WO 00/08202

129

PCT/US99/17845

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.54 (s, 1H, CONH), 7.6, 7.59, 7.49, 7.21, 6.81-6.88 (multiplets, 7H, =CH-, aromatic), 3.73 (s, 3H, OCH<sub>3</sub>), 3.26-3.33 (m, 2H, 2x-CH), 1.2 (d, J = 7.23Hz, 12H, 4xCH<sub>3</sub>).

5 MS m/z 336.

**Example 2: 5-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one**

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-chloro-2-oxindole to give 0.3 g of 5-chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.69 (s, 1H, CONH), 7.67 (s, 1H, =CH-), 7.6 (d, J = 1.85Hz, 1H, H-4), 7.51 (s, 2H, aromatic), 7.26 (dd, J = 1.85, 8.3Hz, 1H, H-6), 6.89 (d, J = 8.3Hz, 1H, H-7), 3.74 (s, 3H, OCH<sub>3</sub>), 3.27-3.34 (m, 2H, 2x-CH), 1.22 (d, J = 6.7Hz, 12H, 4xCH<sub>3</sub>).

**Example 3: N-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide**

Tin chloride dihydrate (225 g) was added to a solution of 2,4-dinitrophenylacetic acid (22.6 g) in ethanol (450 ml). The mixture was heated at 90° C for 10 hours. The reaction mixture was cooled and adjusted to pH 11 with 12M sodium hydroxide. The solids were removed by filtration and the filtrate was concentrated. The residue was treated with ethanol (300 ml). Insoluble materials were removed by filtration and washed with ethanol (5 x 60 ml). The combined ethanol solutions were evaporated and the solid obtained was dried under vacuum to give 15g of 6-amino-2-oxindole as a brown powder.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.03 (s, br, NH), 6.78 (d, J = 8.55Hz, 1H, H-4), 6.09-6.11 (m, 2H), 4.95 (s, br, 2H, NH<sub>2</sub>), 3.22 (s, 2H, H-3). MS (+ APCI) m/z (relative intensity, %) MS m/z 147 ([M-1]<sup>+</sup>, 100).

To a mixture of 6-amino-2-oxindole (1 g) and acetyl



WO 00/08202

130

PCT/US99/17845

chloride (0.79 g) in 10 ml of dichloromethane at 0° C was added triethylamine (1 g). The mixture was brought to room temperature and stirred for 4 hours protected from atmospheric moisture with a dry-tube. The solvent was then evaporated and the residue was recrystallized from methanol to give 0.98 g (77%) of 6-acetylamino-2-oxindole as a beige solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.28 (s, br, NH), 9.83 (s, br, NH), 7.34 (d, J = 1.4Hz, 1H, H-7), 7.06 (d, J = 7.9Hz, 1H, H-4), 6.97 (dd, J = 1.4, 7.9Hz, 1H, H-5), 3.37 (s, 2H, H-3), 2.01 (s, 3H, CH<sub>3</sub>).

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 6-acetamido-2-oxindole to give 0.3 g of N-[3-(3,5-diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.48 (s, 1H, CONH), 9.99 (s, 1H, CONH), 7.51 (d, J = 8Hz, 1H, H-4), 7.46 (s, 2H, aromatic), 7.44 (s, 1H, =CH-), 7.41 (d, J = 2Hz, 1H, H-7), 6.91 (dd, J = 2,8Hz, 1H, H-5), 3.73 (s, 3H, OCH<sub>3</sub>), 3.26-3.34 (m, 2H), 2.03 (s, 3H, CH<sub>3</sub>), 1.21 (d, J = 7Hz, 12H, 4xCH<sub>3</sub>).

MS m/z 392.

**Example 4: 3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-hydroxy-1,3-dihydroindol-2-one**

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 6-hydroxy-2-oxindole to give 0.3 g of 3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-hydroxy-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.34 (s, 1H, CONH), 9.79 (s, 1H, OH), 7.42 (d, J = 8Hz, 1H, H-4), 7.43 (s, 2H, aromatic), 7.33 (s, 1H, =CH-), 6.32 (d, J = 2Hz, 1H, H-7), 6.21 (dd, J = 2,8Hz, 1H, H-5), 3.72 (s, 3H, OCH<sub>3</sub>), 3.25-3.33 (m, 2H), 1.2 (d, J = 7Hz, 12H, 4xCH<sub>3</sub>).

MS m/z 351.

**Example 5: 5-Acetyl-3-(3,5-diisopropyl-4-methoxybenzylidene)-**

WO 00/08202

131

PCT/US99/17845 ,

1,3-dihydroindol-2-one

2-Oxindole (3 g) was suspended in 1,2-dichloroethane and slowly treated with 3.2 ml of acetyl chloride. The resulting suspension was heated at 50° C for 5 hours, cooled, and poured  
5 into water. The resulting precipitate was collected by vacuum filtration, washed copiously with water and dried under vacuum to give 2.9 g (73% yield) of 5-acetyl-2-oxindole as a brown solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.75 (s, br, NH), 7.83 (d, J = 8.23Hz, 1H), 7.78 (s, 1H, H-4), 6.88 (d, J = 8.23Hz, 1H),  
10 3.53 (s, 2H, CH<sub>2</sub>), 2.49 (s, 3H, CH<sub>3</sub>).

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-acetyl-2-oxindole to give 0.3 g of 5-Acetyl-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a  
15 yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.95 (s, 1H, CONH), 8.22 (d, J = 2Hz, 1H, H-4), 7.93 (dd, J = 2,8Hz, 1H, H-6), 7.69 (s, 1H, =CH-),  
7.53 (s, 2H, aromatic), 6.97 (d, J = 8Hz, 1H, H-7), 3.75 (s, 3H, OCH<sub>3</sub>), 3.24-3.34 (m, 2H), 2.48 (s, 3H, CH<sub>3</sub>), 1.24 (d, J =  
20 7Hz, 12H, 4xCH<sub>3</sub>).

MS m/z 377.

**Example 6:**     3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid methyl ester

2-Oxindole (82.9 g) was suspended in 630 ml of acetic  
25 acid in a reaction vessel equipped with mechanical stirring and the mixture cooled to 10° C in an ice water bath. Solid N-iodosuccinimide (175 g) was added in portions over 10 minutes. After the addition was complete the mixture was stirred for 1 hour at 10° C. The suspended solid which was always present  
30 became very thick at this time. The solid was collected by vacuum filtration, washed with 100 ml of 50% acetic acid in water and then with 200 ml of water and subjected to vacuum filtration for 20 minutes to partially air dry it. The product

WO 00/08202

132

PCT/US99/17845

was then dried under vacuum to give 93.5 g (36%) of 5-iodo-2-oxindole containing about 5% 2-oxindole by proton NMR.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.45 (s, 1H, NH-1), 7.49 (s, 1H, H-4), 7.48 (d, *J* = 8.10 Hz, 1H, H-6), 6.64 (d, *J* = 8.10 Hz, 1H, H-7), and 3.46 (s, 2H, CH<sub>2</sub>-3).

MS (*m/z* (relative intensity, %)) 258 ([M-1]<sup>+</sup>, 13).

5-Iodo-2-oxindole (17 g) was refluxed with 2 g of palladium diacetate, 18.2 g of triethylamine, 150 ml of methanol, 15 ml of dimethylsulfoxide and 2.6 g of DPPP in an atmosphere saturated with carbon monoxide. After 24 hours, the reaction was filtered to remove the catalyst and the filtrate was concentrated. The concentrate was chromatographed on silica gel using 30% ethyl acetate/hexane as the eluent. The fractions containing product were concentrated and allowed to stand. The precipitated product was collected by vacuum filtration to give 0.8 g (7%) of 5-methoxycarbonyl-2-oxindole as an off-white solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.70 (s, br, 1H, NH-1), 7.83 (dd, *J* = 1.77, 8.29 Hz, 1H, H-6), 7.77 (s, br, 1H, H-4), 6.89 (d, *J* = 8.29 Hz, 1H, H-7), 3.80 (s, 3H, COOCH<sub>3</sub>-5), 3.51 (s, 2H, CH<sub>2</sub>-3).

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-methoxycarbonyl-2-oxindole to give 0.25 g of 3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid methyl ester as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.96 (s, 1H, CONH), 8.29 (d, *J* = 1Hz, 1H, H-4), 7.88 (dd, *J* = 1,8Hz, 1H, H-6), 7.7 (s, 1H, =CH-), 7.55 (s, 2H, aromatic), 7.0 (d, *J* = 8Hz, 1H, H-7), 3.76 (s, 3H, OCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 3.26-3.35 (m, 2H), 1.23 (d, *J* = 7Hz, 12H, 4xCH<sub>3</sub>).

MS *m/z* 393.

**Example 7:**     3-(3-Isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

WO 00/08202

133

PCT/US99/17845

2-Isopropylphenol was methylated and then formylated to give 3-isopropyl-4-methoxybenzaldehyde. 3-Isopropyl-4-methoxybenzaldehyde was condensed with 2-oxindole to give 0.3 g of 3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.5 (s, 1H, CONH), 7.7, 7.6, 7.2, 7.1, 6.8 (multiplets, 8H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 1H, CH), 1.2 (d, 6H, 2xCH<sub>3</sub>).

**Example 8:**     3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one

5-Isopropyl-4-methoxy-2-methylbenzaldehyde was condensed with 2-oxindole to give 0.25 g of 3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.5 (s, 1H, CONH), 7.6, 7.4, 7.3, 7.2, 6.9, 6.8, 6.8 (multiplets, 7H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 1H, CH), 2.3 (s, 3H, CH<sub>3</sub>), 1.2 (d, 6H, 2xCH<sub>3</sub>).

**Example 9:**     5-Chloro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one

2-Isopropyl-5-methylphenol was methylated and then formylated to give 5-isopropyl-4-methoxy-2-methylbenzaldehyde.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.1 (s, 1H, CHO), 7.6, 6.9 (2xs, 2H, aromatic), 3.9 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 1H, CH), 2.6 (s, 3H, CH<sub>3</sub>), 1.2 (d, 6H, 2xCH<sub>3</sub>).

5-Isopropyl-4-methoxy-2-methylbenzaldehyde was condensed with 5-chloro-2-oxindole to give 0.3 g of 5-Chloro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.65 (s, 1H, CONH), 7.73 (s, 1H, ), 7.49 (s, 1H, ), 7.4 (d, J = 2Hz, 1H, H-4), 7.23 (dd, J = 2,8Hz, 1H, H-6), 6.98 (s, 1H), 6.86 (d, J = 8Hz, 1H, H-7), 3.86 (s, 3H, OCH<sub>3</sub>), 3.23-3.3 (m, 2H), 2.33 (s, 3H, CH<sub>3</sub>), 1.15 (d, J = 7Hz, 6H, 2xCH<sub>3</sub>).

WO 00/08202

134

PCT/US99/17845

MS m/z 341.

**Example 10:**     3-(3-Cyclopentyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

3-Cyclopentyl-4-methoxybenzaldehyde was condensed with 2-oxindole to give 0.25 g of 3-(3-Cyclopentyl-4-methoxybenzylidene)-1,3-dihydro-indol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.5 (s, 1H, CONH), 7.6, 7.5, 7.2, 7.1, 6.8 (multiplets, 8H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 1H, CH), 1.8 (multiplets, 8H, 4xCH<sub>2</sub>).

**Example 11:**     3-(3-Cyclopentyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one

2-Cyclopentylphenol was methylated and then formylated to give 3-cyclopentyl-4-methoxybenzaldehyde.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 9.9 (s, 1H, CHO), 7.8, 7.7, 7.2 (multiplets, 3H, aromatic), 3.9 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 1H, CH), 1.2 (multiplets, 8H, 4xCH<sub>2</sub>).

3-Cyclopentyl-4-methoxybenzaldehyde was condensed with 5-fluoro-2-oxindole to give 0.25 g of 3-(3-cyclopentyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.5 (s, 1H, CONH), 7.6, 7.6, 7.6, 7.4, 7.1, 7.0, 6.8 (multiplets, 7H, aromatic, vinyl CH), 3.9 (s, 3H, CH<sub>3</sub>), 3.3 (m, 1H, CH), 2.0-1.5 (multiplets, 8H, 4xCH<sub>2</sub>).

MS m/z 337.

**Example 12:**     3-(3-Cyclohexyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

2-Cyclohexylphenol was methylated and then formylated.

The aldehyde was condensed with oxindole to give 0.3 g of 3-(3-Cyclohexyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.5 (s, 1H, CONH), 7.7, 7.6, 7.6, 7.2, 7.1, 6.8, 6.8 (multiplets, 7H, aromatic, =CH-), 3.9 (s, 3H,

WO 00/08202

135

PCT/US99/17845 ,

OCH<sub>3</sub>), 3.0 (m, 1H, CH), 1.8, 1.4 (multiplets, 10H, 5xCH<sub>2</sub>).

**Example 13:** 5-Bromo-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one

2-Phenylanisole was formylated to give 4-methoxy-3-phenylbenzaldehyde.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 9.9 (s, 1H, CHO), 7.9, 7.8, 7.5, 7.4, 7.4, 7.3 (multiplets, 8H, aromatic), 3.9 (s, 3H, OCH<sub>3</sub>).

MS m/z 213.

4-Methoxy-3-phenylbenzaldehyde was condensed with 5-bromo-2-oxindole to give 0.3 g of 5-Bromo-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.6 (s, 1H, CONH), 7.8-6.8 (multiplets, 12H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>).

**Example 14:** 5-Chloro-3-(2,3-dihydro-benzofuran-5-ylmethylene)-1,3-dihydroindol-2-one

2,3-Dihydro-5-formylbenzofuran was condensed with 5-chloro-2-oxindole to give 0.25 g of 5-Chloro-3-(2,3-dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.6 (s, 1H, CONH), 8.6, 8.2, 7.7, 7.6, 7.2, 6.9, 6.8 (multiplets, 7H, aromatic, =CH-), 4.6 (t, 2H, OCH<sub>2</sub>), 3.2 (t, 2H, OCH<sub>2</sub>).

**Example 15:** 5-Chloro-3-(2,2-dimethyl-chroman-6-ylmethylene)-1,3-dihydroindol-2-one

2,2-Dimethyl-6-formylchromane was condensed with 5-chloro-2-oxindole to give 0.3 g of 5-Chloro-3-(2,2-dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.61 (s, 1H, CONH), 8.31, 8.27, 7.76, 7.17, 6.79 (multiplets, 6H, aromatic), 7.8 (s, 1H, =CH-), 2.78 (t, 2H, CH<sub>2</sub>), 1.81 (t, 2H, CH<sub>2</sub>), 1.31 (s, 6H, 2xCH<sub>3</sub>).

MS m/z 340.5 (M+1).

WO 00/08202

136

PCT/US99/17845

**Example 16: N-{3-[3-Cyclohexyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide**

Triphenylphosphine (7.47 g, 19.58 mmol) was added to a solution of 2-cyclohexyl-4-chlorophenol (6 g, 28.48 mmol) and  
5 2-hydroxyethylmorpholine (3.5 g, 28.48 mmol) in tetrahydrofuran (50 ml), followed by the dropwise addition of diethylazodicarboxylate (4.5 ml, 28.48 mmol). The mixture stirred at room temperature for 12 hours. The reaction mixture was concentrated under reduced pressure and triturated with  
10 dichloromethane/ hexanes and then washed with more hexanes. Chromatography (silica, 10-20-30% ethyl acetate/hexanes) afforded 8.4 g (95%) of 3-cyclohexyl-4-morpholinoethoxy-chlorobenzene as a yellowish oil.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 7.12 (m, 2H, 2xAr-H), 6.93 (d,  
15 J = 9Hz, 1H, Ar-H), 4.04 (m, 2H, O-CH<sub>2</sub>CH<sub>2</sub>), 3.55 (m, 4H, 2x ringO-CH<sub>2</sub>CH<sub>2</sub>), 2.84 (m, 1H, CHcyclohexyl), 2.68 (m, 2H, N-CH<sub>2</sub>CH<sub>2</sub>), 2.47 (m, 4H, 2x ringN-CH<sub>2</sub>CH<sub>2</sub>), 1.7 (m, 5H, cyclohexyl), 1.3 (m, 5H, cyclohexyl).

MS m/z 324.4 and 325.4, [M+1]<sup>+</sup> and [M+3]<sup>+</sup>.

20 Naphthalene (3.48 g, 27.2 mmol) was added to a suspension of 30% lithium dispersion (3.2 g, 138 mmol, pre-washed with tetrahydrofuran) in tetrahydrofuran (80 ml). The suspension was stirred until a green color appeared (approximately 1 hour). The reaction was cooled to -78°C, and a solution of the  
25 3-cyclohexyl-4-morpholinoethoxychlorobenzene (8 g, 24.7 mmol) in tetrahydrofuran (20ml) was added. After 1 hour, the reaction was warmed to 0°C and stirred for an additional hour. Dimethylformamide (9.6 ml, 123.5 mmol) was added, the reaction was stirred at 0°C for one more hour and then warmed to room  
30 temperature and stirred for another hour. The reaction was quenched with methanol (30 ml), added to 1N hydrochloric acid (300 ml) and extracted with ethyl acetate (400 ml). The organic layer was washed with water (300 ml), saturated sodium

WO 00/08202

137

PCT/US99/17845

bicarbonate solution (300 ml) and brine (300 ml), dried over magnesium sulfate and concentrated. Chromatography (silica, 40-60% ethyl acetate/hexanes) afforded 3 g (34%) of 3-cyclohexyl-4-morpholinoethoxybenzaldehyde as a light yellow oil.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 9.83 (s, 1H, CHO), 7.12 (m, 2H, 2xAr-H), 7.13 (d, J = 8Hz, 1H, Ar-H), 4.19 (m, 2H, O-CH<sub>2</sub>CH<sub>2</sub>), 3.55 (m, 4H, 2x ringO-CH<sub>2</sub>CH<sub>2</sub>), 2.88 (m, 1H, CHcyclohexyl), 2.74 (m, 2H, N-CH<sub>2</sub>CH<sub>2</sub>), 2.49 (m, 4H, 2x ringN-CH<sub>2</sub>CH<sub>2</sub>), 1.7 (m, 5H, cyclohexyl), 1.3 (m, 5H, cyclohexyl).

MS m/z 318.4 [M+1]<sup>+</sup>.

A mixture of 3-cyclohexyl-4-morpholinoethoxybenzaldehyde (3 g, 9.45 mmol), 6-acetamido-2-oxindole (3.5g, 18.9 mmol) and piperidine (5 ml, 50 mmol) in ethanol (35 ml) was held in a sealed tube at 100°C for 6 hours. The reaction mixture was cooled, diluted with ethyl acetate (500 ml) and filtered to remove excess 6-acetamido-2-oxindole. The filtrate was extracted with 0.5N hydrochloric acid (200 ml) and brine (2 X 200 ml). The acid wash was basified with solid sodium bicarbonate to pH 9 and extracted with ethyl acetate (2 x 200 ml). The combined organic layers were washed with brine (200 ml), dried over magnesium sulfate and concentrated. The combined solids were dissolved in dichloromethane (25 ml) and precipitated with diethyl ether (500 ml) to afford 1.6g of a mustard yellow solid. The solid was dissolved in methanol (100 ml) and filtered, affording a yellow solid (which was set aside) and then precipitated with water (400 ml) and filtered. The solid residue was redissolved in methanol and concentrated to a solid. This process was repeated twice to afford ~82% pure isomer 1 by HPLC. The filtered yellow solid was 77% isomer 2 by HPLC.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.45 (s, 1H, NH), 10.0 (s, 1H, NHAc), 7.55 (m, 3H, Ar-H), 7.44 (m, 2H, Ar-H and Ar-CH=C),



WO 00/08202

138

PCT/US99/17845

7.07 (d, 1H,  $J = 8\text{Hz}$ , Ar-H), 6.89 (dd, 1H,  $J = 2$  and  $8\text{Hz}$ , Ar-H), 4.16 (m, 2H, Ar-OCH<sub>2</sub>), 3.57 (m, 4H, OCH<sub>2</sub>CH<sub>2</sub>N), 2.92 (m, 1H, chexCH), 2.75 (m, 2H, ArOCH<sub>2</sub>-CH<sub>2</sub>N), 2.51 (m, 4H, NCH<sub>2</sub>CH<sub>2</sub>O), 2.02 (s, 3H, NHCOCH<sub>3</sub>), 1.8 (m, 5H, chexCH<sub>2</sub>), 1.38 (m, 5H, chexCH<sub>2</sub>).

5 20% isomer 1; 77% isomer 2

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>)  $\delta$ : 10.39 (s, 1H, NH), 9.92 (s, 1H, NHAc), 8.37 (d, 1H,  $J = 2\text{Hz}$ , Ar-H), 8.24 (m, 1H, Ar-H), 7.55 (s, 1H, Ar-CH=C), 7.52 (d, 1H,  $J = 8\text{Hz}$ , Ar-H), 7.36 (m, 1H, Ar-H), 7.03 (m, 2H, Ar-H), 4.16 (m, 2H, Ar-OCH<sub>2</sub>), 3.57 (m, 10 4H, OCH<sub>2</sub>CH<sub>2</sub>N), 2.88 (m, 1H, chexCH), 2.74 (m, 2H, ArOCH<sub>2</sub>-CH<sub>2</sub>N), 2.51 (m, 4H, NCH<sub>2</sub>CH<sub>2</sub>O), 2.03 (s, 3H, NHCOCH<sub>3</sub>), 1.8 (m, 5H, chexCH<sub>2</sub>), 1.38 (m, 5H, chexCH<sub>2</sub>).

**Example 17: 3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-methoxy-1,3-dihydroindol-2-one**

15 3,5-Diisopropyl-4-hydroxybenzaldehyde was methylated to give 3,5-diisopropyl-4-methoxybenzaldehyde.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>)  $\delta$ : 10.1 (s, 1H, CHO), 6.9 (s, 2H, aromatic), 3.9 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 2H, 2xCH), 1.2 (d, 12H, 4xCH<sub>3</sub>).

20 A solution of 11.4 g of hydroxylamine hydrochloride in water (50 ml) was added to a solution of chloral hydrate (9.6 g) and sodium sulfate (83 g) in water (200 ml) at 60°C. The mixture was held at 60°C while, in a separate flask, a solution of 4-anisidine (6.4 g) and concentrated hydrochloric acid (4.3 25 ml) in water (80 ml) was warmed to 80°C. The first solution was then added to the second and the reaction was refluxed for 2 minutes, cooled slowly to room temperature and then cooled in an ice bath. The tan precipitate which formed was collected by vacuum filtration, washed with water and dried under vacuum to 30 give 8.6 g (85 % yield) of N-(2-hydroximinoacetyl)anisidine.

Concentrated sulfuric acid (45 ml) containing water (5 ml) was warmed to 60°C and 8.6 g of N-(2-hydroximinoacetyl)-anisidine was added in one portion. The mixture was stirred

WO 00/08202

139

PCT/US99/17845

at 93°C for 10 minutes and then allowed to cool to room temperature. The mixture was poured into 500 g of ice and extracted 3 times with ethyl acetate. The combined extracts were dried over anhydrous sodium sulfate and concentrated to give 5.1 g (65 % yield) of 5-methoxyisatin as a dark red solid.

A mixture of 5-methoxyisatin (5.0 g) and 30 ml of hydrazine hydrate was refluxed for 15 minutes. The reaction mixture was cooled to room temperature and 50 ml of water was added. The mixture was extracted 3 times with 25 ml of ethyl acetate, the organic layers combined, dried over anhydrous sodium sulfate and concentrated to give a yellow solid. The solid was stirred in ethyl acetate and 1.1 g of insoluble material removed by vacuum filtration and saved. This material proved to be 2-hydrazinocarbonylmethyl-4-anisidine. The filtrate was concentrated and chromatographed on silica gel eluting with ethyl acetate:hexane 1:1 to give 0.7 g of 5-methoxy-2-oxindole as a dirty yellow solid. The 1.1 g of 2-hydrazinocarbonylmethyl-4-anisidine was refluxed for 1 hour in 20 ml of 1 N sodium hydroxide. The mixture was cooled, acidified to pH 2 with concentrated hydrochloric acid and extracted 3 times with 25 ml of ethyl acetate. The organic extracts were combined, washed with brine, dried over anhydrous sodium sulfate and concentrated to give 0.8 g of 5-methoxy-2-oxindole as a dirty yellow solid. The combined yield was 1.5 g or 33 %.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.13 (s, 1H, NH-1), 6.84 (s, 1H, H-4), 6.72 (d, J = 9Hz, 1H, H-6), 6.69 (d, J = 9Hz, 1H, H-7), 3.68 (s, 3H, OCH<sub>3</sub>-5), 3.41 (s, 2H, CH<sub>2</sub>-3).

MS m/z 163 [M+1]<sup>+</sup>.

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-methoxy-2-oxindole to give 0.3 g of 3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-methoxy-1,3-dihydroindol-2-one as a

WO 00/08202

140

PCT/US99/17845

yellow-orange solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.34 (s, 1H, NH), 7.59 (s, 2H), 7.47 (s, 2H), 7.15 (d, J = 2Hz, 1H), 6.85 (dd, J = 2, 8Hz, 1H), 6.58 (d, J = 8Hz, 1H), 3.73 (s, 3H, OCH<sub>3</sub>), 3.63 (s, 3H, OCH<sub>3</sub>), 3.25-3.39 (m, 2H, 2 x (CH<sub>3</sub>)<sub>2</sub>CH), 1.2 (d, 12H, 2 x (CH<sub>3</sub>)<sub>2</sub>CH).

MS APCI m/z 366.1 [M+1]<sup>+</sup>.

**Example 18: N-[3-(4-Methoxy-3-thiophene-3-ylbenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide**

Tetrakis(triphenylphosphine)palladium(0) (1.35 g) was added to a solution of 4-methoxy-3-bromobenzaldehyde (6.72 g) in toluene (45 ml) and ethanol (45 ml), followed by addition of 2M aqueous sodium carbonate (80 ml). To this mixture was added thiophene-3-boronic acid (5 g). The mixture was then refluxed for 12 hours. The reaction mixture was poured into water (200 ml) and extracted into ethyl acetate (2 x 150 ml).

The organic layer was washed with saturated aqueous sodium bicarbonate (150 ml) and brine (150 ml), dried over magnesium sulfate and concentrated. Chromatography (silica, 20% ethyl acetate/hexanes) afforded 6 g (88%) of 3(3-thiophene)-4-methoxybenzaldehyde as a yellow oil.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 9.92 (s, 1H, CHO), 8.04 (d, 1H, J = 2Hz, 1xAr-H), 7.8 (m, 2H, Ar-H), 7.58 (dd, 1H, J = 3 and 5Hz, thiophene and Ar-H), 7.50 (dd, 1H, J = 2 and 5Hz, Ar-H), 7.28 (d, 1H, J = 9Hz, thiophene), 3.94 (s, 3H, OCH<sub>3</sub>).

MS m/z 219.2 [M+1]<sup>+</sup>.

A mixture of 3(3-thiophene)-4-methoxybenzaldehyde (1.72 g), 6-acetamido-2-oxindole (1.5g) and piperidine (4 ml) in ethanol (26 ml) was held in a sealed tube for 12 hours at 100°C. The reaction was cooled and poured into diethyl ether.

The precipitate which formed was removed by filtration, washed with diethyl ether and then hexanes and dried to afford 0.9 g (29%) of N-[3-(4-methoxy-3-thiophene-3-ylbenzylidene)-2-

WO 00/08202

141

PCT/US99/17845

oxo-2,3-dihydro-1H-indol-6-yl]-acetamide as a tan solid.

<sup>1</sup>HNMR 360 MHz, DMSO-d<sub>6</sub>) δ: 10.52 (s, 1H, NH), 10.22 (s, 1H, NHAc), 7.85 (m, 2H, 2xAr-H), 7.7 (dd, J = 2 and 9Hz, 1H, Ar-H), 7.6 (m, 2H, 2xAr-H), 7.5 (dd, 1H, J = 1 and 5Hz, Ar-H),  
5 7.47 (s, 2H, Ar-H and Ar-CH=C), 7.25 (d, 1H, J = 9Hz, Ar-H),  
6.96 (dd, 1H, J = 2 and 9Hz, Ar-H), 3.92 (s, 3H, OCH<sub>3</sub>), 2.04 (s, 3H, NHCOCH<sub>3</sub>).

MS m/z 391.2 [M+1]<sup>+</sup>.

**Example 19:**     3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-methyl-1,3-dihydroindol-2-one

A mixture of 5-methylisatin (15.0 g) and 60 ml of hydrazine hydrate was stirred at 140 - 160°C for 4 hours. The reaction mixture was cooled to room temperature, poured into 300 ml of ice water and acidified to pH 2 with 6 N  
15 hydrochloric acid. After standing at room temperature for 2 days, a precipitate formed which was collected by vacuum filtration, washed with water and dried under vacuum to give 6.5 g (47 % yield) of 5-methyl-2-oxindole.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.20 (s, br, 1H, NH-1), 6.99  
20 (s, 1H, H-4), 6.94 (d, J = 8 Hz, 1H, H-6), 6.68 (d, J = 8Hz, 1H, H-7), 3.39 (s, 2H, CH<sub>2</sub>-3), and 2.22 (s, 3H, CH<sub>3</sub>-5).

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-methyl-2-oxindole to give 0.3 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-5-methyl-1,3-dihydroindol-2-one as a  
25 yellow-orange solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.41 (s, br, 1H, NH), 7.56 (s, 1H), 7.51 (s, 2H), 7.50 (s, br, 1H), 7.03 (d, br, J = 8Hz, 1H), 6.76 (d, J = 8Hz, 1H), 3.74 (s, 3H, OCH<sub>3</sub>), 3.26-3.35 (m, 2H, 2 x (CH<sub>3</sub>)<sub>2</sub>CH), 2.15 (s, 3H, CH<sub>3</sub>), 1.22 (d, J = 7Hz, 12H, 2  
30 x (CH<sub>3</sub>)<sub>2</sub>CH).

MS APCI m/z 350.2 [M+1]<sup>+</sup>.

**Example 20:**     5-Amino-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

WO 00/08202

142

PCT/US99/17845

2-Oxindole (6.5 g) was dissolved in concentrated sulfuric acid (25 ml) and the mixture maintained at -10 to 15°C while fuming nitric acid (2.1 ml) was added dropwise. After the addition of the nitric acid the reaction mixture was stirred  
5 at 0°C for 0.5 hour and poured into ice-water. The precipitate which formed was collected by filtration, washed with water and crystallized from 50% acetic acid. The final crystalline product was then filtered, washed with water and dried under vacuum to give 6.3 g (70%) of 5-nitro-2-oxindole.

10 5-Nitro-2-oxindole (6.3 g) was hydrogenated in methanol over 10% palladium on carbon to give 3.0 g (60% yield) of 5-amino-2-oxindole as a white solid.

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-amino-2-oxindole to give 0.3 g of 5-Amino-3-(3,5-  
15 diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.62 (br s, 1H, NH), 8.31 (s, 2H), 7.8 (s, 1H), 7.65 (s, 1H), 7.55 (d, 1H), 7.47 (s, 1H), 7.09 (dd, J = 2, 8Hz, 1H), 6.88 (d, J = 8Hz, 1H), 3.72 (3H, OCH<sub>3</sub>), 3.24-3.32 (m, 2H, 2 x (CH<sub>3</sub>)<sub>2</sub>CH), 1.23 (d, J = 6.5Hz, 12H, 2 x (CH<sub>3</sub>)<sub>2</sub>CH).

**Example 21: 5-Chloro-3-(4-methoxy-3,5-dimethylbenzylidene)-1,3-dihydroindol-2-one**

2,6-Dimethylphenol was methylated and then formylated.

25 <sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.1 (s, 1H, CHO), 6.9 (s, 2H, aromatic), 3.9 (s, 3H, OCH<sub>3</sub>), 2.6 (s, 6H, 2xCH<sub>3</sub>).

3,5-Dimethyl-4-methoxybenzaldehyde was condensed with 5-chloro-2-oxindole to give 0.3 g of 5-chloro-3-(4-methoxy-3,5-dimethylbenzylidene)-1,3-dihydroindol-2-one as a yellow-orange  
30 solid.

**Example 22: 3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-fluoro-1,3-dihydroindol-2-one**

A suspension of sodium hydride (2.6 g) and

WO 00/08202

143

PCT/US99/17845

dimethylmalonate (14.5 g) in dimethylsulfoxide (160 ml) was stirred at 100°C for 1 hour. The mixture was cooled to room temperature and 7.95 g of 2,5-difluoronitrobenzene was added and stirring was continued for 30 minutes. The mixture was  
5 then heated to 100°C and stirred for 1 additional hour, then cooled to room temperature and poured into 400 ml of saturated ammonium chloride. The mixture was extracted with ethyl acetate (200 ml) and the organic layer washed with brine, dried over anhydrous sodium sulfate and concentrated under  
10 vacuum. The residue was crystallized from methanol to give 24.4 g (80% yield) of dimethyl 4-fluoro-2-nitrophenylmalonate as a white solid. The filtrate was concentrated and chromatographed on a column of silica gel (ethyl acetate:hexane 1:8) to give 5.03 g of dimethyl 4-fluoro-2-  
15 nitrophenylmalonate, for a total of 29.5 g (96 % yield).

Dimethyl 4-fluoro-2-nitrophenylmalonate (5.0 g) was refluxed in 20 ml of 6 N hydrochloric acid for 24 hours. The reaction was cooled and the white solid which formed was collected by vacuum filtration, washed with water and dried to  
20 give 3.3 g (87% yield) of 4-fluoro-2-nitrophenylacetic acid.

4-Fluoro-2-nitrophenylacetic acid (3.3 g) dissolved in acetic acid (15 ml) was hydrogenated over 0.45 g of 10% palladium on carbon under 60 psi for 2 hours. The catalyst was removed by filtration and washed with 15 ml of methanol.  
25 The combined filtrates were concentrated and diluted with water. The precipitate was collected by vacuum filtration, washed with water and dried to give 1.6 g (70% yield) of 6-fluoro-2-oxindole. The filtrate was concentrated to give a purple solid with an NMR spectrum similar to the first crop.  
30 Chromatography of the purple solid (ethyl acetate:hexane 1:2, silica gel) gave a second crop of 6-fluoro-2-oxindole as a white solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.43 (s, 1H, NH-1), 7.17 (t,

WO 00/08202

144

PCT/US99/17845

$J = 8\text{ Hz}$ , H-4), 6.69 (ddd,  $J = 2, 8, 10\text{ Hz}$ , 1H, H-5), 6.6 (dd,  $J = 2, 9\text{ Hz}$ , 1H, H-7), and 3.42 (s, 2H,  $\text{CH}_2\text{-3}$ ).

MS  $m/z$  152.8  $[\text{M}+1]^+$ .

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with  
5 6-fluoro-2-oxindole to give 0.3 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-6-fluoro-1,3-dihydroindol-2-one as a yellow-orange solid.

$^1\text{H NMR}$  (360 MHz,  $\text{DMSO-d}_6$ )  $\delta$ : 10.69 (s, br, 1H, NH), 7.58 (s, 1H, vinyl), 7.55-7.59 (m, 1H), 7.47 (s, 2H), 6.68 (m, 2H),  
10 3.73 (s, 3H,  $\text{OCH}_3$ ), 3.25-3.34 (m, 2H, 2 x  $(\text{CH}_3)_2\text{CH}$ ), 1.2 (d,  $J = 7\text{ Hz}$ , 12H, 2 x  $(\text{CH}_3)_2\text{CH}$ ).

MS  $m/z$  353.

**Example 23:** 3-(2,2-Dimethylchroman-6-ylmethylene)-5-fluoro-1,3-dihydroindol-2-one

15 2,2-Dimethyl-6-formylchromane (commercially available) was condensed with 5-fluoro-2-oxindole to give 0.3 g of 3-(2,2-Dimethylchroman-6-ylmethylene)-5-fluoro-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 24:** 5-Chloro-3-[3,5-diisopropyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-1,3-dihydroindol-2-one

20 Triphenylphosphine (5.14 g, 19.58 mmol) was added to a solution of 3,5-diisopropyl-4-hydroxybenzaldehyde (4 g, 19.58 mmol) in tetrahydrofuran (40 ml) followed by addition of 2-hydroxyethylmorpholine (2.57 g, 19.58 mmol) and then the  
25 dropwise addition of diethylazodicarboxylate (3.41 g, 19.58 mmol). The mixture was stirred at room temperature for 12 hours. The reaction was concentrated under reduced pressure and partitioned between 2N hydrochloric acid (200 ml) and ethyl acetate (150 ml). The aqueous layer was extracted with  
30 ethyl acetate (2 x 150ml), basified to pH 9 with solid sodium bicarbonate and extracted with ethyl acetate (3 x 150 ml). The organic layers were combined, dried over magnesium sulfate and concentrated to afford 900 mg (14%) of 3,5-diisopropyl-4-(2-

WO 00/08202

145

PCT/US99/17845

morpholin-4-ylethoxy)-benzaldehyde as a yellowish oil.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 9.91 (s, 1H, CHO), 9.37 (s, 2H, 2xAr-H), 3.85 (t, *J* = 5Hz, 2H, O-CH<sub>2</sub>CH<sub>2</sub>), 3.59 (m, *J* = 5 and 4Hz, 4H, 2x ringO-CH<sub>2</sub>CH<sub>2</sub>), 3.39 (sept, *J* = 7Hz, 2H, 2xCHCH<sub>3</sub>), 2.72 (t, *J* = 5Hz, 2H, N-CH<sub>2</sub>CH<sub>2</sub>), 2.4 (m on DMSO, 4H, 2x ringN-CH<sub>2</sub>CH<sub>2</sub>), 1.2 (d, *J* = 7Hz, 12H, 4xCH<sub>3</sub>).

A mixture of 3,5-diisopropyl-4-(2-morpholin-4-ylethoxy)-benzaldehyde (0.4 g, 1.26 mmol), 5-chloro-2-oxindole (0.21g, 1.26 mmol) and pyrrolidine (0.5 ml, 6.3 mmol) in ethanol (2 ml) was held in sealed tube at 100°C for 12 hours. The mixture was then poured into 1N hydrochloric acid (100 ml) and the solid which remained were filtered and washed with more water (50 ml). The solid was then dissolved in ethyl acetate (200 ml), the solution dried over magnesium sulfate, filtered and then concentrated. The solid which remained was chromatographed (silica, 4/4/1 dichloromethane/ hexanes/ methanol) to give 60 mg (10%) of 5-chloro-3-[3,5-diisopropyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-1,3-dihydroindol-2-one as a brownish-orange solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.54 (s, 1H, NH), 9.43 (d, *J* = 10Hz, 1H, Ar-H), 7.58 (d, *J* = 9Hz, 1H, Ar-H), 7.46 (d, *J* = 5Hz, 2H, Ar-H), 7.3 (m, 1H, Ar-H), 7.19 (m, *J* = 9 and 10Hz, 1H, Ar-H), 6.99 (m, *J* = 9 and 9Hz, 1H, Ar-H), 6.83 (m, *J* = 9 and 1Hz, 1H, Ar-H), 3.61 (dd, *J* = 9 and 22 Hz, 1H, 1xCH<sub>2</sub>CH), 3.43 (m, 1H, CHCH<sub>3</sub>), 2.86 (dd, *J* = 4 and 22Hz, 1H, 1xCH<sub>2</sub>CH), 1.35 (d, *J* = 9Hz, 3H, CH<sub>3</sub>).

MS *m/z* 262.0 [M+1]<sup>+</sup>.

**Example 25: 3-(3,5-Diisopropyl-4-methoxybenzylidene)-7-fluoro-1,3-dihydroindol-2-one**

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 7-fluoro-2-oxindole to give 0.25 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-7-fluoro-1,3-dihydroindol-2-one as a yellow-orange solid.



WO 00/08202

146

PCT/US99/17845 ,

**Example 26:    3-(4-Methoxy-3-thiophen-3-ylbenzylidene)-5-(2-morpholin-4-yl-ethyl)-1,3-dihydroindol-2-one**

A solution of 5-chloroacetyl-2-oxindole (4.18 g) in trifluoroacetic acid (30 ml) was cooled in an ice bath and  
5 4.65 g of triethylsilane added. The mixture was then stirred at room temperature for 3 hours. The mixture was poured into 150 ml of water and the precipitate which formed was collected by vacuum filtration, washed with 50 ml of water and dried to give 2.53 g (65 % yield) of 5-(2-chloroethyl)-2-oxindole as a  
10 reddish-brown solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.26 (s, br, NH), 7.11 (s, 1H, H-4), 7.05 (d, J = 8Hz, 1H), 6.73 (d, J = 8Hz, 1H), 3.77 (t, J = 7Hz, 2H, CH<sub>2</sub>), 3.42 (s, 2H, CH<sub>2</sub>), 2.94 (t, J = 7Hz, 2H, CH<sub>2</sub>).

15 A solution of 5-(2-chloroethyl)-2-oxindole (2.3 g), morpholine (1.2 ml) and diisopropylethylamine (1.2 ml) in dimethylsulfoxide (10 ml) was heated overnight at 100°C. The mixture was cooled, poured into water and extracted with ethyl acetate. The organic layer was washed with brine, dried and  
20 evaporated. The residue was chromatographed (silica gel, 5 % methanol in chloroform) to give 0.9 g (31 %) of 5-(2-morpholin-4-yl-ethyl)-2-oxindole as a white solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.21 (s, br, 1H, NH), 7.06 (s, br, 1H, H-4), 7.0 (d, J = 8Hz, 1H, H-6), 6.71 (d, J = 8Hz, 1H, H-7), 3.55-3.58 (m, 4H), 3.41 (s, 2H, H-3), 2.63-2.68 (m, 2H), 2.39-2.47 (m, 6H).

A mixture of 3(3-thiophene)-4-methoxybenzaldehyde (0.27 g, 1.22 mmol), 5-(2-morpholin-4-yl-ethyl)-2-oxindole (0.3g, 1.22 mmol) and piperidine (0.6 ml, 6.1 mmol) in ethanol (4 ml)  
30 was held in a sealed tube at 100°C for 12 hours. The reaction was cooled and poured into diethyl ether (150 ml) and hexanes (150 ml). The solid which remained was removed by filtration, washed with diethyl ether and then hexanes and dried to afford

WO 00/08202

147

PCT/US99/17845

0.27 g (50%) of 3-(4-methoxy-3-thiophen-3-ylbenzylidene)-5-(2-morpholin-4-yl-ethyl)-1,3-dihydroindol-2-one as an orange-yellow solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.41 (s, 1H, NH), 8.86 (d, J = 2.15Hz, 1H, Ar-H), 8.39 (dd, J = 2 and 9Hz, 1H, Ar-H), 7.8 (dd, 1H, J = 1 and 3Hz, Ar-H), 7.7 (s, 1H, Ar-CH=C), 7.6 (m, 1H, ArH), 7.5 (m, 2H, 2xArH), 7.21 (d, J = 9Hz, 1H, Ar-H), 7.02 (dd, 1H, J = 1 and 8Hz, Ar-H), 6.73 (d, 1H, J = 8Hz, Ar-H), 3.93 (s, 3H, OCH<sub>3</sub>), 3.58 (m, 4H, 2x OCH<sub>2</sub>CH<sub>2</sub>N), 2.7 (m, 2H, ArCH<sub>2</sub>CH<sub>2</sub>), 2.5 (m, 2H, ArCH<sub>2</sub>CH<sub>2</sub>N), 2.4 (m, 4H, 2xOCH<sub>2</sub>CH<sub>2</sub>N).

MS m/z 447.2 [M+1]<sup>+</sup>.

**Example 27: N-[3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide**

A mixture of 5-isopropyl-4-methoxy-2-methylbenzaldehyde (0.25 g, 1.3 mmol), 6-acetamido-2-oxindole (0.25g, 1.3 mmol) and pyrrolidine (0.54 ml, 6.5 mmol) in ethanol (4 ml) was held in a sealed tube at 100°C for 12 hours. The reaction was cooled and then poured into water (100 ml). The solid which formed was removed by filtration, washed with water and then dissolved in ethyl acetate (200 ml). The ethyl acetate solution was dried over magnesium sulfate and concentrated. The resulting solid was triturated with dichloromethane/hexanes to afford 0.33 g (70%) of N-[3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide as a brownish yellow solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.46 (s, 1H, NH), 7.49 (s, 1H, Ar-H), 7.46 (s, 1H, Ar-H), 7.41 (s, 1H, Ar-H), 6.93 (m, 1H, Ar-CH=C), 6.83 (dd, 1H, J = 2 and 8Hz, Ar-H), 3.83 (s, 3H, OCH<sub>3</sub>), 2.30 (s, 3H, Ar-CH<sub>3</sub>), 2.02 (s, 3H, NHCOCH<sub>3</sub>), 1.2 (m, 1H, CH(CH<sub>3</sub>)<sub>2</sub>), 1.1 (d, J = 7Hz, 6H, CH(CH<sub>3</sub>)<sub>2</sub>).

MS m/z 365.2 [M+1]<sup>+</sup>.

**Example 28: 3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-ethyl-1,3-dihydroindol-2-one**

WO 00/08202

148

PCT/US99/17845

2-Oxindole (3 g) suspended in 1,2-dichloroethane was slowly treated with 3.2 ml of acetyl chloride. The resulting suspension was stirred at 50°C for 5 hours, cooled, and poured into water. The resulting precipitate was collected by vacuum  
5 filtration, washed copiously with water and dried under vacuum to give 2.9 g (73 % yield) of 5-acetyl-2-oxindole as a brown solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ : 10.75 (s, br, NH), 7.83 (d, J = 8Hz, 1H), 7.78 (s, 1H, H-4), 6.88 (d, J = 8Hz, 1H), 3.53 (s,  
10 2H, CH<sub>2</sub>), 2.49 (s, 3H, CH<sub>3</sub>).

5-Acetyl-2-oxindole (2 g) and 15 ml of trifluoroacetic acid in an ice bath was slowly treated with 1.8 g of triethylsilane and then stirred at room temperature for 5 hours. One ml of triethylsilane was added and stirring  
15 continued overnight. The reaction mixture was poured into ice water and the resulting precipitate collected by vacuum filtration, washed copiously with water and dried under vacuum to give 1.3 g (71 % yield) of 5-ethyl-2-oxindole as a yellow solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.25 (s, br, NH-1), 7.03 (s, 1H, H-4), 6.97 (d, J = 8Hz, 1H, H-6), 6.69 (d, J = 8Hz, 1H, H-7), 3.40 (s, 2H, CH<sub>2</sub>-3), 2.51 (q, J = 8Hz, 2H, CH<sub>2</sub>CH<sub>3</sub>-5), and 1.12 (t, J = 7Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>-5).

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with  
25 5-ethyl-2-oxindole to give 0.3 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-5-ethyl-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.42 (s, 1H, NH), 7.56 (s, 1H), 7.51 (s, 2H), 7.49 (s, br, 1H), 7.08 (dd, J = 8Hz, 1H),  
30 6.79 (d, J = 8Hz, 1H), 3.74 (s, 3H, OCH<sub>3</sub>), 3.25-3.55 (m, 2H, 2 x (CH<sub>3</sub>)<sub>2</sub>CH), 2.44 (q, J = 8Hz, 2H, CH<sub>3</sub>CH<sub>2</sub>), 1.22 (d, J = 7Hz, 12H, 2 x (CH<sub>3</sub>)<sub>2</sub>CH), 1.08 (t, J = 8Hz, 3H, CH<sub>3</sub>CH<sub>2</sub>).

MS APCI m/z 364.3 [M+1]<sup>+</sup>.

WO 00/08202

149

PCT/US99/17845

**Example 29: N-[2'-Methoxy-5'-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-biphenyl-3-yl]-acetamide**

A mixture of 3-bromo-p-anisaldehyde (1 g, 4.65 mmol), pyrrolidine (1.9 ml, 23.2 mmol) and oxindole (0.62 g, 4.65 mmol) in dimethylformamide (7 ml) was held in a sealed tube at 100°C for 12 hours. The mixture was cooled to room temperature and poured into 1N hydrochloric acid (100 ml). The precipitate which formed was filtered and washed with water. The precipitate was then dissolved in ethyl acetate (200 ml), the solution washed with brine, dried over magnesium sulfate and then concentrated. The solid obtained was chromatographed (silica, 30-40% ethyl acetate/ hexanes) to give 0.63 g (42%) of 3-(3-bromo-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.5 (m, 1H, NH), 7.7 (m, 1H, Ar-H), 7.6 (m, 1H, Ar-CH=C), 7.5 (m, 1H, Ar-H), 7.2 (m, 2H, 2xAr-H), 6.9 (m, 1H, Ar-H), 6.8 (m, 1H, Ar-H), 3.9 (s, 3H, OCH<sub>3</sub>). MS m/z 330.0 and 332.0.

[M]<sup>+</sup> and [M+2]<sup>+</sup>.

Tetrakis(triphenylphosphine)palladium(0) (0.02 g, 0.02 mmol) was added to a solution of 3-(3-Bromo-4-methoxybenzylidene)-1,3-dihydroindol-2-one (0.2 g, 0.61 mmol) in toluene (1 ml) and ethanol (2 ml), followed by addition of 2M aqueous sodium carbonate (1.2 ml, 2.4 mmol). To this mixture was added 3-acetamidophenylboronic acid (0.12 g, 0.67 mmol), and the mixture was heated to 100°C in a sealed tube and held there for 12 hours. The reaction was then poured into water (50 ml) and extracted with ethyl acetate (2x100 ml). The combined organic layers were washed with saturated aqueous sodium bicarbonate (50 ml) and brine (50 ml). The organic layer was dried over magnesium sulfate and concentrated. Chromatography (silica, 30-40% ethyl acetate/hexanes) afforded a waxy solid that was triturated with diethyl ether/hexanes

WO 00/08202

150

PCT/US99/17845

and then dried to afford 0.05 g (22%) of *N*-[2'-methoxy-5'-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-biphenyl-3-yl]-acetamide as a reddish orange solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.6 (m, 1H, NH), 9.88 (s, 1H, NHAc), 7.7 (m, 1H, Ar-H), 7.5 (m, 3H, 2xAr-H and Ar-CH=C), 7.2 (m, 4H, Ar-H), 7.0 (m, 2H, Ar-H), 6.8 (m, 2H, Ar-H), 3.9 (two s, 3H, OCH<sub>3</sub>), 2.02 (2, 3H, COCH<sub>3</sub>).

MS m/z 385.2 [M+1]<sup>+</sup>.

**Example 30:** 5-Fluoro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one

5-Isopropyl-4-methoxy-2-methylbenzaldehyde was condensed with 5-fluoro-2-oxindole to give 0.25 g of 5-fluoro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.54 (s, 1H, NH), 7.72 (s, 1H), 7.47 (s, 1H), 7.1 (dd, 1H), 7.04 (ddd, 1H), 6.98 (s, 1H), 6.82-6.86 (m, 1H), 3.86 (s, 3H, OCH<sub>3</sub>), 3.25-3.28 (m, 1H, (CH<sub>3</sub>)<sub>2</sub>CH), 2.32 (s, 3H, CH<sub>3</sub>), 1.14 (d, 6H, (CH<sub>3</sub>)<sub>2</sub>CH).

EI MS m/z 325 [M]<sup>+</sup>.

**Example 31:** *N*-[3-(4-Methoxy-3-thiophen-2-ylbenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide

Tetrakis(triphenylphosphine)palladium(0) (0.24 g) was added to a solution of 4-methoxy-3-bromobenzaldehyde (1.5 g) in toluene (15 ml) and ethanol (15 ml), followed by addition of 2M aqueous sodium carbonate (14 ml). To this mixture was added thiophene-2-boronic acid (0.98 g), and the mixture was heated to reflux. After 3 hours, the reaction was partitioned between water (100 ml) and ethyl acetate (250 ml). The organic layer was washed with saturated aqueous sodium bicarbonate (75 ml) and brine (75 ml), dried over magnesium sulfate and concentrated. Chromatography (silica, 20% ethyl acetate/hexanes) afforded 1.3 g (87%) of 3-(2-thiophene)-4-methoxybenzaldehyde as a yellow oil.

WO 00/08202

151

PCT/US99/17845

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 9.92 (s, 1H, CHO), 8.2 (d, 1H, J = 3Hz, 1xAr-H), 7.8 (dd, 1H, J = 10 and 10Hz, SCHCHCH), 7.65 (m, 1H, Ar-H), 7.60 (dd, 1H, J = 2 and 6Hz, Ar-H), 7.32 (d, 1H, J = 10Hz, SCHCHCH), 7.13 (d, 1H, J = 10 and 6Hz, SCHCHCH),  
5 3.99 (s, 3H, OCH<sub>3</sub>).

A mixture of 3-(2-thiophene)-4-methoxybenzaldehyde (0.25 g, 1.15 mmol), 6-acetamido-2-oxindole (0.22g, 1.15 mmol) and pyrrolidine (0.48 ml, 5.75 mmol) in ethanol (4 ml) was held in a sealed tube at 100°C for 12 hours. The reaction was cooled  
10 and the precipitate which formed was removed by filtration, washed with ethanol and then hexanes and dried to afford 0.15 g (33%) of N-[3-(4-methoxy-3-thiophen-2-ylbenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide as a yellow solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.49 (s, 1H, NH), 9.95 (s, 1H, NHAc), 8.99 (d, J = 2Hz, 1H, Ar-H), 8.28 (dd, J = 2 and 9Hz, 1H, Ar-H), 7.6 (m, 2H, 2xAr-H), 7.5 (m, 2H, Ar-H and Ar-CH=C), 7.38 (d, 1H, J = 1Hz, Ar-H), 7.22 (d, 1H, J = 9Hz, Ar-H), 7.14 (dd, 1H, J = 4 and 5Hz, Ar-H), 7.05 (dd, 1H, J = 2 and 9Hz, Ar-H), 3.97 (s, 3H, OCH<sub>3</sub>), 2.04 (s, 3H, NHCOCH<sub>3</sub>).

20 MS m/z 391.3 [M]<sup>+</sup> and [M+2]<sup>+</sup>.

**Example 32: 6-Amino-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one**

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 6-amino-2-oxindole to give 0.3 g of 6-amino-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a  
25 yellow-orange solid.

**Example 33: N-[3-(2,2-Dimethylchroman-6-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide**

A mixture of 6-acetamido-2-oxindole (285 mg), 2,2-dimethylchroman-6-carboxaldehyde (285 mg) and piperidine (0.2 ml) was stirred at 90°C overnight. The reaction mixture was cooled and diluted with 20 ml of water. The sticky solid which formed was filtered and chromatographed (2% ethanol in ethyl  
30

WO 00/08202

152

PCT/US99/17845

acetate) to give 240 mg (44%) of N-[3-(2,2-dimethylchroman-6-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide as an orange red solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.44 (s, 1H, NH), 10.0 (s, 1H, NH), 7.6, 7.46, 7.42, 7.37, 6.93, 6.82 (m, 7H, 6 Ar-H and 1 vinyl-H), 2.8 (t, J = 6Hz, 2H, CH<sub>2</sub>), 2.03 (s, 3H, CH<sub>3</sub>), 1.83 (t, J = 6Hz, 2H, CH<sub>2</sub>), 1.32 (s, 6H, 2x CH<sub>3</sub>).

MS m/z 363.1 [M+1]<sup>+</sup>.

**Example 34: 5-Bromo-3-(2,2-dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one**

2-Oxindole (1.3 g) in 20 ml of acetonitrile was cooled to -10°C and 2.0 g of N-bromosuccinimide was slowly added with stirring. The reaction was stirred for 1 hour at -10°C and then 2 hours at 0°C. The precipitate which formed was collected, washed with water and dried to give 1.9 g (90% yield) of 5-bromo-2-oxindole.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.44 (s, br, 1H, NH-1), 7.32-7.36 (m, 2H), 6.76 (d, J = 8.5 Hz, 1H, H-7), 3.5 (s, 2H, CH<sub>2</sub>).

MS m/z 212.1 and 214.1, [M]<sup>+</sup> and [M+2]<sup>+</sup>.

2,2-Dimethyl-6-formylchromane (commercially available) was condensed with 5-bromo-2-oxindole to give 0.3 g of 5-bromo-3-(2,2-dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one as a yellow-orange solid.

Isomer 1 (67%): <sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.65 (s, br, 1H, NH), 7.73 (d, 1H), 7.6 (s, 1H, vinyl), 7.5 (s, br, 1H), 7.47 (dd, 1H), 7.37 (dd, 1H), 6.86 (d, 1H), 6.82 (d, 1H), 2.77-2.81 (m, 4H, 2 x CH<sub>2</sub>), 1.81-1.84 (m, 4H, 2 x CH<sub>2</sub>), 1.32 (s, 6H, 2 x CH<sub>3</sub>).

Isomer 2 (33%): <sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.63 (s, br, 1H, NH), 8.32 (dd, 1H), 8.27 (s, br, 1H), 7.89 (d, 1H), 7.81 (s, 1H, vinyl), 7.3 (dd, 1H), 6.79 (d, 1H), 6.76 (d, 1H), 2.77-2.81 (m, 4H, 2 x CH<sub>2</sub>), 1.81-1.84 (m, 4H, 2 x CH<sub>2</sub>), 1.31 (s, 6H, 2 x CH<sub>3</sub>).

WO 00/08202

153

PCT/US99/17845

MS APCI m/z 384.1 and 385.9, [M]<sup>+</sup> and [M+2]<sup>+</sup>.

**Example 35:**     3-(4-Methoxy-3-thiophen-3-ylbenzylidene)-1,3-dihydroindol-2-one

Tetrakis(triphenylphosphine)palladium(0) (0.02 g, 0.02  
5 mmol) was added to a solution of 3-(3-bromo-4-methoxy-  
benzylidene)-1,3-dihydroindol-2-one (0.2 g, 0.61 mmol) in  
toluene (1 ml) and ethanol (1 ml), followed by addition of 2M  
aqueous sodium carbonate (1.2 ml, 2.4 mmol). To this mixture  
was added thiophene-3-boronic acid (0.09 g, 0.67 mmol) and the  
10 mixture was held at 100°C in a sealed tube for 12 hours. The  
reaction was then poured into water (50 ml) and extracted with  
ethyl acetate (2x100 ml). The combined organic layers were  
washed with saturated aqueous sodium bicarbonate (50 ml) and  
brine (50 ml), dried over magnesium sulfate and concentrated.  
15 Chromatography (silica, methylene chloride then 30-40% ethyl  
acetate/hexanes) afforded 0.1 g (50%) of 3-(4-methoxy-3-  
thiophen-3-ylbenzylidene)-1,3-dihydroindol-2-one as a yellow  
solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.54 (s, 1H, NH), 7.8 (m, 2H,  
20 2xAr-H), 7.6 (m, 5H, 4xAr-H and Ar-CH=C), 7.2 (m, 2H, Ar-H),  
6.8 (m, 2H, Ar-H), 3.92 (s, 3H, OCH<sub>3</sub>).

MS m/z 334.3 [M+1]<sup>+</sup>.

**Example 36:**     5-Bromo-3-(5-isopropyl-4-methoxy-2-methyl-  
benzylidene)-1,3-dihydroindol-2-one

25 5-Isopropyl-4-methoxy-2-methylbenzaldehyde was condensed  
with 5-bromo-2-oxindole to give 0.3 g of 5-bromo-3-(5-  
isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-  
one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.5 (s, 1H, CONH), 7.7, 7.6, 7.5, 7.4,  
30 7.0, 6.8 (m, 6H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 3.3 (m,  
1H, CH), 2.3 (s, 3H, CH<sub>3</sub>), 1.2 (d, 6H, 2xCH<sub>3</sub>).

**Example 37:**     5-Fluoro-3-(6-methoxybiphenyl-3-ylmethylene)-  
1,3-dihydroindol-2-one



WO 00/08202

154

PCT/US99/17845

4-Methoxy-3-phenylbenzaldehyde was condensed with 5-fluoro-2-oxindole to give 0.3 g of 5-fluoro-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one as a yellow-orange solid.

5

**Example 38: 3-(3-Isopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one**

Diethyl oxalate (30 ml) in 20 ml of dry ether was added with stirring to 19 g of potassium ethoxide suspended in 50 ml of dry ether. The mixture was cooled in an ice bath and 20 ml of 3-nitro-o-xylene in 20 ml of dry ether was slowly added. The thick dark red mixture which formed was refluxed for 0.5 hour, concentrated to a dark red solid, and treated with 10 % sodium hydroxide until almost all of the solid dissolved. The dark red mixture was treated with 30% hydrogen peroxide until the red color changed to yellow. The mixture was then treated alternately with 10% sodium hydroxide and 30% hydrogen peroxide until the dark red color no longer formed on addition of sodium hydroxide. The remaining solid was filtered and the filtrate acidified with 6 N hydrochloric acid. The resulting precipitate was collected by vacuum filtration, washed with water, and dried under vacuum to give 9.8 g (45% yield) of 1-methyl-6-nitrophenylacetic acid as an off-white solid. The solid was hydrogenated in methanol over 10% palladium on carbon to give 9.04 g of 4-methyl-2-oxindole as a white solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.27 (s, br, 1H, NH), 7.06 (t, J = 8Hz, 1H, H-6), 6.74 (d, J = 8Hz, H-5), 6.63 (d, J = 8Hz, 1H, H-7), 3.36 (s, 2H, CH<sub>2</sub>), 2.18 (s, 3H, CH<sub>3</sub>).

3-Isopropyl-4-methoxybenzaldehyde was condensed with 4-methyl-2-oxindole to give 0.25 g of 3-(3-isopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.4(s, 1H, NH), 8.2, 8.1, 7.6, 7.2,

WO 00/08202

155

PCT/US99/17845

7.1, 6.8, 6.7 (m, 7H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 1H, CH), 2.6 (m, 1H, CH), 1.2 (d, 6H, 2xCH<sub>3</sub>).

**Example 39: 3-(4,5-Dimethoxy-2-thiophen-2-ylbenzylidene)-1,3-dihydroindol-2-one**

5 A mixture of 6-bromoveratraldehyde (1 g, 4.08 mmol), oxindole (0.54 g, 4.08 mmol) and pyrrolidine (1.7 ml, 20.4 mmol) in dimethylformamide (6 ml) was held in a sealed tube at 100°C for 12 hours. The reaction mixture was cooled and added to 1N hydrochloric acid (100 ml). The precipitate which formed  
10 was filtered and washed with water. The precipitate was then dissolved in ethyl acetate (200 ml), the solution washed with brine, dried over magnesium sulfate and concentrated. The solid obtained was chromatographed (silica, 30-4-% ethyl acetate/ hexanes) to give 1.2 g (81%) of 3-(3-bromo-4,5-  
15 dimethoxybenzylidene)-1,3-dihydroindol-2-one as a yellow solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.5 (s, 1H, NH), 7.4 (s, 1H, Ar-CH=C), 7.3 (m, 2H, 2xAr-H), 7.2 m, 1H, Ar-H), 6.8 (m, 2H, 2xAr-H), 3.8 (s, 3H, OCH<sub>3</sub>), 3.7 (s, 3H, OCH<sub>3</sub>).

20 MS m/z 360.5 and 362.5, [M]<sup>+</sup> and [M+2]<sup>+</sup>.

Tetrakis(triphenylphosphine)palladium(0) (0.03 g, 0.03 mmol) was added to a solution of 3-(3-bromo-4,5-dimethoxybenzylidene)-1,3-dihydroindol-2-one (0.32 g, 0.89 mmol) in toluene (2 ml) and ethanol (2 ml), followed by  
25 addition of 2M aqueous sodium carbonate (1.8 ml, 3.6 mmol). To this mixture was added thiophene-2-boronic acid (0.13 g, 0.98 mmol), and the mixture was held at 100°C in a sealed tube for 12 hours. The reaction mixture was poured into water (50 ml) and extracted with ethyl acetate (2x100 ml). The combined  
30 organic layers were washed with saturated aqueous sodium bicarbonate (50 ml) and brine (50 ml), dried over magnesium sulfate and concentrated. Chromatography (silica, 30-40% ethyl acetate/hexanes) afforded an orange solid that was

WO 00/08202

156

PCT/US99/17845

trituated with methylene chloride and dried to afford 0.19 g (59%) of 3-(4,5-dimethoxy-2-thiophen-2-ylbenzylidene)-1,3-dihydroindol-2-one as an orange solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.5 (m, 1H, NH), 7.6 (m, 1H, Ar-H), 7.5 (m, 2H, Ar-H), 7.3 (m, 1H, Ar-CH=C), 7.2 (m, 4H, Ar-H), 6.8 (m, 2H, Ar-H), 6.8 (m, 2H, Ar-H), 3.9 (two s, 3H, OCH<sub>3</sub>), 3.7 (two s, 3H, COCH<sub>3</sub>).

MS m/z 364.0 [M+1]<sup>+</sup>.

10 **Example 40:**     N-{3-[4-(2-Morpholin-4-ylethoxy)-3-thiophen-2-ylbenzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide

Triphenylphosphine (6.85 g) was added to a solution of 3-bromo-4-hydroxybenzaldehyde (5 g) in tetrahydrofuran (40 ml), followed by addition of 2-hydroxyethylmorpholine (3.01 ml) and  
15 then the dropwise addition of diethylazodicarboxylate (4.11 ml). After 12 hours, the reaction was concentrated under reduced pressure, poured into 2N hydrochloric acid (200 ml) and extracted with ethyl acetate (2x 150 ml). The aqueous layer was basified to pH 9 with solid sodium bicarbonate and  
20 extracted with ethyl acetate (2 x 150 ml). The organic layers were dried over magnesium sulfate and concentrated to afford 4.4 g (56%) of 4-morpholinoethoxy-3-bromobenzaldehyde as a reddish oil.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 9.84 (s, 1H, NH), 8.07 (d, 1H, J = 2Hz, Ar-H), 7.90 (dd, 1H, J = 2 and 8Hz, Ar-H), 7.32 (d, 1H, J = 8Hz, Ar-H), 4.29 (t, 2H, J = 6Hz, Ar-OCH<sub>2</sub>), 3.56 (m, 4H, OCH<sub>2</sub>CH<sub>2</sub>N), 2.76 (t, 2H, J = 6Hz, ArOCH<sub>2</sub>-CH<sub>2</sub>N), 2.50 (m, 4H, NCH<sub>2</sub>CH<sub>2</sub>O).

Tetrakis(triphenylphosphine)palladium(0) (0.23) was added  
30 to a solution of 4-morpholinoethoxy-3-bromobenzaldehyde (2.13 g) in toluene (10 ml) and ethanol (10 ml), followed by addition of 2M aqueous sodium carbonate (13 ml). To this mixture was added thiophene-2-boronic acid (1.13 g), and the mixture was

WO 00/08202

157

PCT/US99/17845

refluxed for 12 hours. The reaction mixture was then poured into water (100 ml) and extracted into ethyl acetate (3 x 100 ml). The product was extracted into 1N hydrochloric acid (2 x 100 ml), the aqueous layers basified to pH 9 with solid sodium bicarbonate and extracted with ethyl acetate (2 x 100ml). The organic layers were washed with brine (75 ml), dried over magnesium sulfate and concentrated to afford 1.7 g (75%) of 3-(2-thiophene)-4-morpholinoethoxybenzaldehyde as a greenish/yellow oil.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>)  $\delta$ : 9.93 (s, 1H, CHO), 8.2 (d, 1H,  $J$  = 2Hz, Ar-H), 7.82 (dd, 1H,  $J$  = 2 and 8Hz, Ar-H), 7.76 (dd, 1H,  $J$  = 1 and 4Hz, Ar-H), 7.59 (dd, 1H,  $J$  = 1 and 5Hz, Ar-H), 7.35 (d, 1H,  $J$  = 9Hz, Ar-H), 7.14 (dd, 1H,  $J$  = 4 and 5Hz, Ar-H), 4.34 (t,  $J$  = 6Hz, 2H, O-CH<sub>2</sub>CH<sub>2</sub>), 3.56 (m, 4H, 2x ringO-CH<sub>2</sub>CH<sub>2</sub>), 2.82 (t,  $J$  = 6Hz, 2H, N-CH<sub>2</sub>CH<sub>2</sub>), 2.5 (m on DMSO, 4H, 2x ringN-CH<sub>2</sub>CH<sub>2</sub>).

MS m/z 318.2 [M+1]<sup>+</sup>.

A mixture of 3(2-thiophene)-4-morpholinoethoxybenzaldehyde (1.74 g), 6-acetamido-2-oxindole(1g) and piperidine (2.6 ml) in ethanol (11 ml) was held in sealed tube at 100°C for 12 hours. The reaction was cooled and added to diethyl ether and the ether decanted. The oily residue was dissolved in methylene chloride (10 ml) and precipitated in ether (200 ml) and hexanes (100 ml). The precipitate was removed by filtration, washed with ether and then hexanes and dried to afford a brownish solid. Chromatography (silica, 5% methanol/dichloromethane) afforded N-{3-[4-(2-morpholin-4-ylethoxy)-3-thiophen-2-ylbenzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide as a dark yellow solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>)  $\delta$ : 10.49 (s, 1H, NH), 10.01 (s, 1H, NHAc), 8.05 (d,  $J$  = 2Hz, 1H, Ar-H), 7.72 (m, 1H, Ar-H), 7.65 (m, 1H, Ar-H), 7.48 (m, 2H, 2xAr-H), 7.47 (s, 1H, Ar-CH=C), 7.45 (d, 1H,  $J$  = 2Hz, Ar-H), 7.29 (d, 1H,  $J$  = 9Hz, Ar-

WO 00/08202

158

PCT/US99/17845 ,

H), 7.13 (m, 1H, Ar-H), 6.90 (dd, 1H,  $J = 2$  and 9Hz, Ar-H), 4.32 (m, 2H,  $\text{ArOCH}_2\text{CH}_2\text{N}$ ), 3.58 (m, 4H,  $2 \times \text{OCH}_2\text{CH}_2\text{N}$ ), 2.84 (m, 2H,  $\text{ArOCH}_2\text{CH}_2\text{N}$ ), 2.5 (m under DMSO, 4H,  $2 \times \text{OCH}_2\text{CH}_2\text{N}$ ), 2.03 (s, 3H,  $\text{NHCOCH}_3$ ).

5 MS m/z 490.0  $[\text{M}+1]^+$ .

**Example 41:** 3-(2,2-Dimethylchroman-6-ylmethylene)-4-methyl-1,3-dihydroindol-2-one

2,2-Dimethyl-6-formylchromane (commercially available) was condensed with 4-methyl-2-oxindole to give 0.25 g of 3-(2,2-dimethylchroman-6-ylmethylene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

$^1\text{H}$ NMR (360 MHz, DMSO- $d_6$ )  $\delta$ : 10.45 (s, br, 1H, NH), 8.17 (dd,  $J = 2$ , 8Hz, 1H), 8.02 (d,  $J = 2$ Hz, 1H), 7.6 (s, 1H, vinyl), 7.05 (t,  $J = 8$ Hz, 1H), 6.75 (t, 2H), 6.67 (d,  $J = 8$ Hz, 1H), 2.77 (t,  $J = 7$ Hz, 2H,  $\text{CH}_2$ ), 2.55 (s, 3H,  $\text{CH}_3$ ), 1.8 (t,  $J = 7$ Hz, 2H,  $\text{CH}_2$ ), 1.31 (s, 6H,  $2 \times \text{CH}_3$ ).

MS APCI m/z 320.2  $[\text{M}+1]^+$ .

**Example 42:** 3-(2,3-Dihydrobenzofuran-5-ylmethylene)-5-fluoro-1,3-dihydroindol-2-one

20 2,3-Dihydro-5-formylbenzofuran (commercially available) was condensed with 5-fluoro-2-oxindole to give 0.25 g of 3-(2,3-dihydrobenzofuran-5-ylmethylene)-5-fluoro-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 43:** 3-(3-Cyclohexyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one

2-Cyclohexylphenol was methylated then formylated. The aldehyde was condensed with 5-fluoro-2-oxindole to give 0.3 g of 3-(3-cyclohexyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one as a yellow-orange solid.

30 **Example 44:** 5-Fluoro-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

3-Isopropyl-4-methoxybenzaldehyde was condensed with 5-fluoro-2-oxindole to give 0.25 g of 5-fluoro-3-(3-isopropyl-4-

WO 00/08202

159

PCT/US99/17845

methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 45:** 3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydro-pyrrolo[2,3-b]pyridin-2-one

5 Zinc dust (6.5 g) was added to a solution of 3,3-dibromo-7-azaoxindole (2.9 g) in acetic acid (20 ml) and acetonitrile (30 ml). The mixture was stirred for 2 hours at room temperature. The solids were removed by filtration and the solvent evaporated. The residue was treated with ethyl  
10 acetate. The ethyl acetate solution containing insoluble solid was passed through a short column of silica gel. The collected ethyl acetate solution was evaporated and the residue dried under vacuum to give 1.8 g (91% yield) of 7-aza-2-oxindole acetic acid salt.

15 <sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.90 (s, br, 1H, NH), 8.01, 7.52, 6.91 (m, 3H), 3.52 (s, 2H, CH<sub>2</sub>).

MS APCI 135 (M+1).

5-Isopropyl-4-methoxy-2-methylbenzaldehyde condensed with 7-aza-2-oxindole to give 0.25 g of 3-(5-isopropyl-4-methoxy-2-  
20 methylbenzylidene)-1,3-dihydro-pyrrolo[2,3-b]pyridin-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.5 (s, 1H, CONH), 8.3-6.8 (m, 6H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 3.3 (m, 1H, CH), 2.3 (s, 3H, CH<sub>3</sub>), 1.2 (d, 6H, 2xCH<sub>3</sub>).

25 **Example 46:** 3-(3'-Ethoxy-6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one

Tetrakis(triphenylphosphine)palladium(0) (0.02 g, 0.02 mmol) was added to a solution of 3-(bromo-4-methoxybenzylidene)-1,3-dihydroindol-2-one (0.2 g, 0.61 mmol) in  
30 toluene (1 ml) and ethanol (1 ml), followed by addition of 2M aqueous sodium carbonate (1.2 ml, 2.4 mmol). To this mixture was added 3-ethoxyphenylboronic acid (0.11 g, 0.67 mmol), and the mixture was held at 100°C in a sealed tube for 12 hours.

WO 00/08202

160

PCT/US99/17845

The reaction mixture was added to water (40 ml) and extracted with ethyl acetate (75 ml). The combined organic layers were washed with saturated aqueous sodium bicarbonate (50 ml) and brine (50 ml). The organic layer was dried over magnesium sulfate and concentrated. The resulting solid was triturated with dichloromethane/ hexanes to afford 0.9 g (39%) of 3-(3'-ethoxy-6-methoxy-biphenyl-3-ylmethylene)-1,3-dihydroindol-2-one as a yellow solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.57 (s, 1H, NH), 7.65 (m, 5H, 4xAr-H and Ar-CH=C), 7.3 (m, 3H, 4xAr-H), 7.15 (m, 2H, Ar-H), 6.9 (m, 2H, Ar-H), 4.05 (q, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.85 (s, 3H, OCH<sub>3</sub>), 1.35 (t, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS m/z 372.5 [M+1]<sup>+</sup>.

**Example 47:** 3-(3-Cyclopentyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one

3-Cyclopentyl-4-methoxybenzaldehyde was condensed with 7-aza-2-oxindole to give 0.25 g of 3-(3-cyclopentyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 1.1 (s, 1H, CONH), 8.3-6.8 (m, 7H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 1H, CH), 1.8 (m, 8H, 4xCH<sub>2</sub>).

**Example 48:** 3-(3-Cyclopentyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

3-Cyclopentyl-4-methoxybenzaldehyde was condensed with 4-methyl-2-oxindole to give 0.25 g of 3-(3-cyclopentyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.4 (s, 1H, CONH), 8.3, 8.1, 7.6, 7.0, 7.0, 6.8, 6.6 (m, 7H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 1H, CH), 1.8 (m, 8H, 4xCH<sub>2</sub>).

**Example 49:** 3-(4,5,2'-Trimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one

WO 00/08202

161

PCT/US99/17845

Tetrakis(triphenylphosphine)palladium(0) (0.02 g, 0.02 mmol) was added to a solution of 3-(3-bromo-4,5-dimethoxybenzylidene)-1,3-dihydroindol-2-one (0.20 g, 0.56 mmol) in toluene (1ml) and ethanol (1 ml), followed by addition of 2M aqueous sodium carbonate (1.1 ml, 2.2 mmol). To this mixture was added 2-methoxyphenylboronic acid (0.09 g, 0.62 mmol), and the mixture was held at 100°C in a sealed tube for 12 hours. The reaction mixture was added to water (50 ml) and extracted with ethyl acetate (2x100 ml). The combined organic layers were washed with saturated aqueous sodium bicarbonate (50 ml) and brine (50 ml). The organic layer was dried over magnesium sulfate and concentrated. Chromatography (silica, 30-40% ethyl acetate / hexanes) afforded 0.14 g (64%) of 3-(4,5,2'-trimethoxy-biphenyl-2-ylmethylene)-1,3-dihydroindol-2-one as an orange solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.4 (m, 1H, NH), 7.5 (m, 1H, Ar-H), 7.3 (m, 2H, Ar-H and Ar-CH=C), 7.2 (m, 4H, Ar-H), 6.9 (m, 4H, Ar-H), 3.8 (two s, 3H, COCH<sub>3</sub>), 3.7 (two s, 3H, OCH<sub>3</sub>), 3.5 (two s, 3H, COCH<sub>3</sub>).

MS m/z 388.2 [M+1]<sup>+</sup>.

**Example 50:** N-{3-[4-(2-Morpholin-4-ylethoxy)-3-thiophen-3-ylbenzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide

Tetrakis(triphenylphosphine)palladium(0) (0.21 g) was added to a solution of 4-morpholinoethoxy-3-bromobenzaldehyde (1.92 g) in toluene (10 ml) and ethanol (10 ml), followed by addition of 2M aqueous sodium carbonate (12 ml). To this mixture was added thiophene-3-boronic acid (1.02 g), and the mixture was heated to reflux. After 12 hours, the reaction mixture was added to water (100 ml) and extracted into ethyl acetate (3 x 100 ml). The product was then extracted into 1N hydrochloric acid (2 x 100 ml), the aqueous layers basified to pH 9 with solid sodium bicarbonate and extracted with ethyl acetate (2 x 100 ml). The organic layers were washed with



WO 00/08202

162

PCT/US99/17845 ,

brine (75 ml), dried over magnesium sulfate and concentrated to afford 1.7 g (83%) of 3-(3-thiophene)-4-morpholinoethoxy benzaldehyde a reddish oil.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 9.91 (s, 1H, CHO), 8.08 (m, 2H, Ar-H), 7.83 (dd, 1H, J = 2 and 9Hz, Ar-H), 7.76 (dd, 1H, J = 1 and 4Hz, Ar-H), 7.59 (dd, 1H, J = 1 and 5Hz, Ar-H), 7.35 (d, 1H, J = 9Hz, Ar-H), 7.64 (dd, 1H, J = 1 and 5Hz, Ar-H), 7.58 (dd, 1H, J = 3 and 5Hz, Ar-H), 7.31 (d, 1H, J = 9Hz, Ar-H), 4.27 (t, J = 5Hz, 2H, O-CH<sub>2</sub>CH<sub>2</sub>), 3.56 (m, 4H, 2x ringO-CH<sub>2</sub>CH<sub>2</sub>), 2.75 (t, J = 6Hz, 2H, N-CH<sub>2</sub>CH<sub>2</sub>), 2.48 (m on DMSO, 4H, 2x ringN-CH<sub>2</sub>CH<sub>2</sub>).

MS m/z 318.2 [M+1]<sup>+</sup>.

A mixture of 3-(3-thiophene)-4-morpholinoethoxy-benzaldehyde (1.74 g), 6-acetamido-2-oxindole (1 g) (prepared as described in example 1) and piperidine (2.6 ml) in ethanol (11 ml) was held in a sealed tube at 100°C for 12 hours. The reaction was cooled and added to diethyl ether (200 ml). The solvent was decanted, and the oily solid which remained was dissolved in dichloromethane (15 ml) and precipitated in diethyl ether (200 ml) and hexanes (200 ml), filtered, washed with diethyl ether and then hexanes and dried to afford a mustard yellow solid. Chromatography (silica, 5% methanol/dichloromethane) afforded N-{3-[4-(2-morpholin-4-ylethoxy)-3-thiophen-3-ylbenzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide as a yellow foam.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.47 (s, 1H, NH), 10.00 (s, 1H, NHAc), 8.07 (m, 1H, Ar-H), 7.91 (d, J = 2Hz, 1H, Ar-H), 7.63 (m, 2H, 2xAr-H), 7.58 (m, 2H, 2xAr-H), 7.48 (s, 1H, Ar-CH=C), 7.45 (d, 1H, J = 2Hz, Ar-H), 7.26 (d, 1H, J = 9Hz, Ar-H), 6.90 (dd, 1H, J = 2 and 9Hz, Ar-H), 4.26 (m, 2H, ArOCH<sub>2</sub>CH<sub>2</sub>N), 3.58 (m, 4H, 2xOCH<sub>2</sub>CH<sub>2</sub>N), 2.76 (m, 2H, ArOCH<sub>2</sub>CH<sub>2</sub>N), 2.48 (m under DMSO, 4H, 2xOCH<sub>2</sub>CH<sub>2</sub>N), 2.03 (s, 3H, NHCOCH<sub>3</sub>).

WO 00/08202

163

PCT/US99/17845 ,

MS m/z 490.0 [M+1]<sup>+</sup>.

**Example 51:**     5-Chloro-3-(3-cyclohexyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

2-Cyclohexylphenol was methylated and then formylated.

5 The aldehyde was condensed with 5-chloro-2-oxindole to give 0.3 g of 5-chloro-3-(3-cyclohexyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.6(s, 1H, CONH), 7.7-6.8(m, 7H, aromatic, =CH-), 3.9(s, 3H, OCH<sub>3</sub>), 3.0 (m, 1H, CH), 1.8, 1.4 (m, 10H, 5xCH<sub>2</sub>).

**Example 52:**     [3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-carbamic acid tert-butyl ester

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 6-tert-butyloxycarbonylamino-2-oxindole to give 0.35 g of [3-(3,5-diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-carbamic acid tert-butyl ester as a yellow-orange solid.

**Example 53:**     3-(3,5-Diisopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

20 3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 4-methyl-2-oxindole to give 0.3 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.41 (s, 1H, CONH), 8.0 (s, 2H), 7.66 (s, 1H), 7.07 (t, 1H), 6.76 (d, 1H), 6.67 (d, 1H), 3.7 (s, 3H, OCH<sub>3</sub>), 3.21-3.3 (m, 1H), 1.21 (d, 12H, 4 x CH<sub>3</sub>).

**Example 54:**     5-Bromo-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

30 3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-bromo-2-oxindole to give 0.3 g of 5-bromo-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

WO 00/08202

164

PCT/US99/17845 ;

**Example 55: N-{3-[3-tert-Butyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide**

Triphenylphosphine (5.89 g, 22.44 mmol) was added to a solution of 3-tert-butyl-4-hydroxybenzaldehyde (4 g, 22.44 mmol) in tetrahydrofuran (40 ml), followed by addition of 2-hydroxyethylmorpholine (2.94 g, 22.44 mmol) and then the dropwise addition of diethylazodicarboxylate (3.91 g, 22.44 mmol). The mixture was allowed to stir at room temperature for 3 days. The reaction was concentrated under reduced pressure and partitioned between 2N hydrochloric acid (200 ml) and ethyl acetate (150 ml). The aqueous layer was extracted with ethyl acetate (2 x 150 ml), basified to pH 9 with solid sodium bicarbonate, saturated with solid sodium chloride and extracted with ethyl acetate (3 x 150 ml). The combined organic layers were dried over magnesium sulfate and concentrated to afford 2.8 g (44%) of 3-tert-butyl-4-(2-morpholin-4-ylethoxy)benzaldehyde as a yellowish oil.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 9.8 (s, 1H, CHO), 7.75 (m, 2H, 2xAr-H), 7.16 (d, J = 10Hz, 1H, Ar-H), 4.19 (t, J = 7Hz, 2H, O-CH<sub>2</sub>CH<sub>2</sub>), 3.55 (m, 4H, 2x ringO-CH<sub>2</sub>CH<sub>2</sub>), 2.75 (t, J = 7Hz, 2H, N-CH<sub>2</sub>CH<sub>2</sub>), 2.4 (m on DMSO, 4H, 2x ringN-CH<sub>2</sub>CH<sub>2</sub>), 1.36 (s, 9H, C(CH<sub>3</sub>)<sub>3</sub>).

MS m/z 292.0 [M+1]<sup>+</sup>.

A mixture of 3-tert-butyl-4-(2-morpholin-4-ylethoxy)benzaldehyde (0.25 g), 6-acetamido-2-oxindole (0.16g) and piperidine (0.43 ml) in ethanol (3 ml) was held in a sealed tube at 100°C for 12 hours. The reaction was cooled, added to ethyl acetate (250 ml), washed with 1N hydrochloric acid (100 ml) and brine (100 ml), dried with magnesium sulfate and concentrated. Some product went into the acid wash, so this was basified with solid sodium bicarbonate to pH 9 and extracted with ethyl acetate (2 x 150 ml). The organic layers were dried over magnesium sulfate, combined with the oil from

WO 00/08202

165

PCT/US99/17845

the first crop of product and concentrated. Chromatography (silica, 40% ethyl acetate followed by 5-10% methanol / methylene chloride) afforded 100 mg (27%) of N-{3-[3-tert-butyl-4-(2-morpholin-4-ylethoxy)benzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide as a yellow solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.46 (s, 1H, NH), 9.99 (s, 1H, NHAc), 7.58 (m, 3H, Ar-H), 7.41 (m, 2H, Ar-H and Ar-CH=C), 7.12 (d, 1H, Ar-H), 6.92 (dd, 1H, J = 2 and 8Hz, Ar-H), 4.18 (m, 2H, Ar-OCH<sub>2</sub>), 3.57 (m, 4H, OCH<sub>2</sub>CH<sub>2</sub>N), 2.77 (m, 2H, ArOCH<sub>2</sub>-CH<sub>2</sub>N), 2.49 (m, 4H, NCH<sub>2</sub>CH<sub>2</sub>O), 2.02 (s, 3H, NHCOCH<sub>3</sub>), 1.38 (s, 9H, Ar-tBu).

MS m/z 464.0 [M+1]<sup>+</sup>.

**Example 56: 3-(4-Methoxy-3,5-dimethylbenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one**

2,6-Dimethylphenol was methylated and then formylated.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.1 (s, 1H, CHO), 6.9 (s, 2H, aromatic), 3.9 (s, 3H, OCH<sub>3</sub>), 2.6 (s, 6H, 2xCH<sub>3</sub>).

The aldehyde was condensed with 7-aza-2-oxindole to give 0.2 g of 3-(4-methoxy-3,5-dimethylbenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 1.5 (s, 1H, CONH), 8.2-6.8 (m, 6H, aromatic, =CH-), 3.8 (s, 3H, OCH<sub>3</sub>), 2.5 (s, 6H, 2xCH<sub>3</sub>), 2.4 (s, 3H, CH<sub>3</sub>).

**Example 57: 5-Bromo-3-[3,5-diisopropyl-4-(2-morpholin-4-ylethoxy)benzylidene]-1,3-dihydroindol-2-one**

A mixture of 3,5-diisopropyl-4-(2-morpholin-4-ylethoxy)-benzaldehyde (0.4 g), 5-bromo-2-oxindole (0.27g) and pyrrolidine (0.5 ml) in ethanol (2 ml) was held in a sealed tube at 100°C for 12 hours. The mixture was then added to 1N hydrochloric acid (100 ml) and the solids removed by filtration and washed with more water (50 ml). The solids were then dissolved in ethyl acetate (200 ml), the solution

WO 00/08202

166

PCT/US99/17845 ,

washed with 1N hydrochloric acid (75 ml) and brine (75 ml), dried with magnesium sulfate and concentrated. The crude solid was chromatographed (silica, 4:4:1 dichloromethane: hexanes: methanol) to give 80 mg (13%) of 5-bromo-3-[3,5-diisopropyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-1,3-dihydroindol-2-one as a reddish-brown solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.54 (s, 1H, NH), 9.43 (d, J = 10Hz, 1H, Ar-H), 7.58 (d, J = 9Hz, 1H, Ar-H), 7.46 (d, J = 5Hz, 2H, Ar-H), 7.3 (m, 1H, Ar-H), 7.19 (m, J = 9 and 10Hz, 1H, Ar-H), 6.99 (m, J = 9 and 9Hz, 1H, Ar-H), 6.83 (m, J = 9 and 1Hz, 1H, Ar-H), 3.61 (dd, J = 9 and 22Hz 1H, 1xCH<sub>2</sub>CH), 3.43 (m, 1H, CHCH<sub>3</sub>), 2.86 (dd, J = 4 and 22Hz, 1H, 1xCH<sub>2</sub>CH), 1.35 (d, J = 9 Hz, 3H, CH<sub>3</sub>).

MS m/z 262.0 [M+1]<sup>+</sup>.

15 **Example 58: 3-(3'-Ethoxy-4,5-dimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one**

Tetrakis(triphenylphosphine)palladium(0) (0.02 g, 0.02 mmol) was added to a solution of 3-(3-bromo-4,5-dimethoxybenzylidene)-1,3-dihydroindol-2-one (0.2 g, 0.56 mmol) in toluene (1 ml) and ethanol (1 ml), followed by addition of 2M aqueous sodium carbonate (1 ml, 2mmol). To this mixture was added 3-ethoxyphenylboronic acid (0.1 g, 0.62 mmol) and the mixture was held at 100°C in a sealed tube for 12 hours. The reaction mixture was then added to water (40 ml) and extracted with ethyl acetate (2 x 75 ml). The combined organic layers were washed with saturated aqueous sodium bicarbonate (50 ml) and brine (50 ml), dried over magnesium sulfate and concentrated. The resulting solid was triturated with methylene chloride / hexanes to afford 0.14 g (64%) of 3-(3'-ethoxy-4,5-dimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one as a yellow/orange solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.5 (s, 1H, NH), 7.6 (d, J = 9Hz, 1H, Ar-H), 7.2 (m, 4H, 3xAr-H and Ar-CH=C), 7.08 (s, 2H,

WO 00/08202

167

PCT/US99/17845

Ar-H), 6.8 (m, 5H, 5xAr-H), 4.0 (q,  $J = 8\text{Hz}$ , 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.89 (s, 3H,  $\text{OCH}_3$ ), 3.77 (s, 3H,  $\text{OCH}_3$ ), 1.29 (t,  $J = 8\text{Hz}$ , 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS  $m/z$  402.5  $[\text{M}+1]^+$ .

5 **Example 59:** 5-chloro-3-(4-methoxy-3-thiophen-2-ylbenzylidene)-1,3-dihydroindol-2-one

A mixture of 3-(2-thiophene)-4-methoxybenzaldehyde (0.21 g), 5-chloro-2-oxindole (0.16g) and pyrrolidine (0.4 ml) in ethanol (1 ml) was placed in a sealed tube and heated to  $100^\circ\text{C}$ .

10 A precipitate formed within 10 minutes so the reaction was diluted with another 10 ml of ethanol and heating resumed. After 4 hours, the reaction was cooled and the solid which formed was removed by filtration and washed with ethanol and then hexanes to afford 0.21 g (60%) of 5-chloro-3-(4-methoxy-15 3-thiophen-2-ylbenzylidene) -1,3-dihydroindol-2-one as a yellow-orange solid.

$^1\text{H}$ NMR (360 MHz,  $\text{DMSO}-d_6$ )  $\delta$ : 10.7 (s, 1H, NH), 7.8-9.1 (m, 3H, Ar-H), 7.6 (m, 3H, 2xAr-H and Ar-CH=C), 7.2 (m, 3H, Ar-H), 6.9 (m, 1H, Ar-H), 3.9 (m, 3H,  $\text{OCH}_3$ ).

20 MS  $m/z$  368.1 and 369.9  $[\text{M}]^+$  and  $[\text{M}+2]^+$ .

**Example 60:** 5-Chloro-3-4-methoxy-3-pyridin-3-ylbenzylidene)-1,3-dihydroindol-2-one

Tetrakis(triphenylphosphine)palladium(0) (0.2 g) was added to a solution of 3-bromo-4-methoxybenzaldehyde (1.25 g) 25 in toluene (10 ml) and ethanol (10 ml), followed by addition of 2M aqueous sodium carbonate (11 ml). To this mixture was added pyridine-3-boronic acid propane diol (1 g), and the mixture was heated to reflux. After 2 hours, the reaction was added to water (75 ml) and extracted with ethyl acetate (2x 75 30 ml). The combined organic layers were washed with saturated aqueous sodium bicarbonate (100 ml) and brine (100 ml), dried over magnesium sulfate and concentrated. Chromatography (silica, 20-50% ethyl acetate/hexanes) afforded 1.08 g (87%)

WO 00/08202

168

PCT/US99/17845 ,

of 3-(3-pyridyl)-4-methoxybenzaldehyde as a white solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 9.93 (s, 1H, CHO), 8.7 (d, 1H, J = 3Hz, Ar-H), 8.56 (dd, 1H, J = 1 and 6Hz, Ar-H), 7.9 (m, 3H, 3xAr-H), 7.46 (dd, 1H, J = 6 and 9Hz, Ar-H), 7.35 (d, 1H, J = 10Hz, Ar-H), 3.89 (s, 3H, OCH<sub>3</sub>).

A mixture of 3-(3-pyridyl)-4-methoxybenzaldehyde (0.25 g), 5-chloro-2-oxindole (0.2g) and pyrrolidine (0.5 ml) in ethanol (2 ml) was placed in a sealed tube and heated to 100°C.

A precipitate formed within 10 minutes so the reaction was diluted with another 10 ml of ethanol and heating resumed. After 4 hours, the reaction was cooled and the solid was removed by filtration and washed with ethanol and then hexanes to afford 0.27 g (61%) of 5-chloro-3-(3-(3-pyridyl)-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 0.67 (s, 1H, NH), 8.6 (m, 3H, 2xAr-H and Ar-CH=C), 7.8 (m, 4H, Ar-H), 7.4 (m, 1H, Ar-H), 7.35 (m, 2H, Ar-H), 6.85 (m, 1H, Ar-H), 3.88 (s, 3H, OCH<sub>3</sub>).

MS m/z 363.0 and 365.1, [M]<sup>+</sup> and [M+2]<sup>+</sup>.

**Example 61: 5-Chloro-3-(4,5-3'-Trimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one**

Tetrakis(triphenylphosphine)palladium(0) (0.36 g) was added to a solution of 4,5-dimethoxy-2-bromobenzaldehyde (2.5 g) in toluene (20 ml) and ethanol (20 ml), followed by addition of 2M aqueous sodium carbonate (20 ml). To this mixture was added 3-methoxyphenylboronic acid (1.7 g), and the mixture was heated to reflux. After 1 hour, the reaction was partitioned between water (150 ml) and ethyl acetate (300 ml). The organic layer was washed with saturated aqueous sodium bicarbonate (150 ml) and brine (150 ml), dried over magnesium sulfate and concentrated. Chromatography (silica, 20-30-40% ethyl acetate / hexanes) afforded 2.7 g (98%) of 2-(3-methoxyphenyl)-4,5-dimethoxybenzaldehyde as a yellow oil which

WO 00/08202

169

PCT/US99/17845 ,

solidified upon standing.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 9.71 (s, 1H, CHO), 7.38 (m, 2H, 2xAr-H), 7.0 (m, 4H, 4xAr-H), 3.9 (s, 3H, OCH<sub>3</sub>), 3.84 (s, 3H, OCH<sub>3</sub>), 3.8 (s, 3H, OCH<sub>3</sub>).

5 MS m/z 273.5 [M+1]<sup>+</sup>.

A mixture of 2-(3-methoxyphenyl)-4,5-dimethoxy-benzaldehyde (0.25 g), 5-chloro-2-oxindole (0.15g) and pyrrolidine (0.4 ml) in ethanol (1 ml) was placed in a sealed tube and heated to 100°C. A precipitate formed within 10  
10 minutes so the reaction was diluted with another 10 ml of ethanol and heating resumed. After 2.5 hours, the reaction was cooled, the solid was removed by filtration and washed with ethanol and then hexanes to afford 0.36 g (92%) of 3-(4,5-3'-trimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one as a  
15 yellow solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.67 (s, 1H, NH), 7.6 (d, J = 2.5Hz, 1H, Ar-H), 7.3 (m, 4H, Ar-H), 7.12 (s, 1H, Ar-CH=C), 6.9 (m, 4H, Ar-H), 3.9 (s, 3H, OCH<sub>3</sub>), 3.79 (s, 3H, OCH<sub>3</sub>), 3.75 (s, 3H, OCH<sub>3</sub>).

20 MS m/z 422.0 and 424.1 [M]<sup>+</sup> and [M+2]<sup>+</sup>.

**Example 62:** 3-(4,5-Dimethoxy-2-naphthalen-2-ylbenzylidene)-1,3-dihydroindol-2-one

Tetrakis(triphenylphosphine)palladium(0) (0.02 g) was added to a solution of 3-(3-bromo-4,5-dimethoxybenzylidene)-  
25 1,3-dihydroindol-2-one (0.20 g) in toluene (1 ml) and ethanol (1 ml) followed by addition of 2M aqueous sodium carbonate (1.1 ml). To this mixture was added naphthalene-2-boronic acid (0.11 g), and the mixture was held at 100°C in a sealed tube for 12 hours. The precipitate which formed was filtered,  
30 washed with ethanol and then dissolved in 20% methanol / toluene, filtered and concentrated. The resulting solid was triturated with hexanes and dried to afford 0.07 g (29%) of 3-(4,5-dimethoxy-2-naphthalen-2-ylbenzylidene)-1,3-dihydroindol-



WO 00/08202

170

PCT/US99/17845

2-one as an orange solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.5 (m, 1H, NH), 7.9 (m, 4H, Ar-H), 7.7 (m, 1H, Ar-H), 7.5 (m, 3H, Ar-H), 7.4 (s, 1H, Ar-CH=C), 7.2 (m, 3H, Ar-H), 6.9 (m, 2H, Ar-H), 3.9 (two s, 3H, COCH<sub>3</sub>), 3.8 (two s, 3H, OCH<sub>3</sub>).

MS m/z 408.0 [M+1]<sup>+</sup>.

**Example 63:**     N-[3-(3'-Acetylamino-6-methoxybiphenyl-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide

3-Bromo-4-methoxybenzaldehyde was condensed with 6-acetamido-2-oxindole to give N-[3-(3-bromo-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide.

Tetrakis(triphenylphosphine)palladium(0) (0.01 g) was added to a solution of N-[3-(3-bromo-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide (0.14 g) in toluene (1 ml) and ethanol (1 ml) followed by addition of 2M aqueous sodium carbonate (0.72 ml). To this mixture was 3-acetylamino-phenylboronic acid (0.07 g), and the mixture was held at 100° C in a sealed tube for 12 hours. The reaction mixture was added to water (50 ml) and extracted with ethyl acetate (2x100 ml). The combined organic layers were washed with saturated aqueous sodium bicarbonate (50 ml) and brine (50 ml), dried over magnesium sulfate and concentrated. Chromatography (silica, dichloromethane then 5-10% methanol/ dichloromethane) afforded 0.07 g (43%) of N-[3-(3'-acetylamino-6-methoxybiphenyl-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide as a yellow solid. About 26% isomer by <sup>1</sup>HNMR.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.49 (s, 1H, NH), 10.42 (s, 1H, NHAc), 9.95 (s, 1H, NHAc), 7.76 (m, 2H, 2xAr-H), 7.58 (m, 3H, 3xAr-H), 7.4 (m, 2H, Ar-H and Ar-CH=C), 7.3 (m, 2H, Ar-H), 7.18 (m, 1H, Ar-H), 6.9 (dd, 2H, J = 2 and 8.5Hz, Ar-H), 3.84 (s, 3H, OCH<sub>3</sub>), 2.05 (s, 3H, NHCOCH<sub>3</sub>), 2.03 (s, 3H, NHCOCH<sub>3</sub>).

MS m/z 442.2 [M+1]<sup>+</sup>.

**Example 64:**     6-Methoxy-3-(4-methoxy-3-thiophen-3-

WO 00/08202

171

PCT/US99/17845

ylbenzylidene)-1,3-dihydroindol-2-one

A mixture of 3(3-thiophene)-4-methoxybenzaldehyde (0.53 g, 2.45 mmol), 6-methoxy-2-oxindole (0.4g, 2.45 mmol) and piperidine (1.2 ml, 12.25 mmol) in 5 ml of ethanol was held in a sealed tube at 100°C for 12 hours. The reaction was cooled and added to diethyl ether (150 ml) and hexanes (150 ml). The precipitate which formed was removed by filtration, washed with diethyl ether and then hexanes and dried to afford 0.57 g (64%) of 6-methoxy-3-(4-methoxy-3-thiophen-3-ylbenzylidene)-1,3-dihydroindol-2-one as a mustard yellow solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.45 (s, 1H, NH), 8.78 (d, J = 2Hz, 1H, Ar-H), 8.30 (dd, J = 2 and 9Hz, 1H, Ar-H), 7.8 (dd, 1H, J = 1.5 and 3Hz, Ar-H), 7.5 (m, 3H, 2xAr-H and Ar-CH=C), 7.18 (d, J = 9Hz, 1H, Ar-H), 6.55 (dd, 1H, J = 2 and 8.5Hz, Ar-H), 6.39 (d, 1H, J = 2Hz, Ar-H), 3.91 (s, 3H, OCH<sub>3</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 2.04 (s, 3H, NHCOCH<sub>3</sub>).

MS m/z 364.1 [M+1]<sup>+</sup>.

**Example 65:** 3-(6-Methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one

4-Methoxy-3-phenylbenzaldehyde was condensed with 2-oxindole to give 0.3 g of 3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.5(s, 1H, CONH), 8.5, 8.5, 7.8, 7.7, 7.5, 7.5, 7.4, 7.4, 7.3, 7.2, 7.2, 7.0, 6.8(m, 13H, aromatic, =CH-), 3.9(s, 3H, OCH<sub>3</sub>).

**Example 66:** 3-(2,3-Dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one

2,3-Dihydro-5-formylbenzofuran (commercially available) was condensed with 2-oxindole to give 0.25 g of 3-(2,3-dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.4 (s, 1H, CONH), 8.6, 8.2, 7.7, 7.6, 7.2, 7.0, 6.9, 6.8 (m, 8H, aromatic, =CH-), 4.6(t, 2H, OCH<sub>2</sub>),

WO 00/08202

172

PCT/US99/17845

3.2 (t, 2H, OCH<sub>2</sub>).

**Example 67:**     5-Chloro-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one

4-Methoxy-3-phenylbenzaldehyde was condensed with 5-chloro-2-oxindole to give 0.3 g of 5-chloro-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 11.5 (s, 1H, CONH), 7.8-6.8 (m, 12H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>).

**Example 68:**     3-(3-Cyclohexyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

3-Cyclohexyl-4-methoxybenzaldehyde was condensed with 4-methyl-2-oxindole to give 0.3 g of 3-(3-cyclohexyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.6 (s, 1H, CONH), 7.7-6.8 (m, 7H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 3.0 (m, 1H, CH), 2.5 (s, 3H, CH<sub>3</sub>), 1.8-1.4 (m, 10H, 5xCH<sub>2</sub>).

**Example 69:**     3-(2,3-Dihydrobenzofuran-5-ylmethylene)-4-methyl-1,3-dihydroindol-2-one

2,3-Dihydro-5-formylbenzofuran (commercially available) was condensed with 4-methyl-2-oxindole to give 0.25 g of 3-(2,3-dihydrobenzofuran-5-ylmethylene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.5 (s, 1H, CONH), 8.4-6.6 (m, 7H, aromatic, =CH-), 4.6 (t, 2H, OCH<sub>2</sub>), 3.2 (t, 2H, OCH<sub>2</sub>), 2.5 (s, 3H, CH<sub>3</sub>).

**Example 70:**     3-(3-Isopropyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one

3-Isopropyl-4-methoxybenzaldehyde was condensed with 7-aza-2-oxindole to give 0.25 g of 3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one as a yellow-orange solid.

WO 00/08202

173

PCT/US99/17845 ,

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 11.5 (s, 1H, CONH), 8.0-6.8 (m, 7H, aromatic, =CH-), 3.7 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 1H, CH), 1.2 (d, 6H, 2xCH<sub>3</sub>).

**Example 71:** 3-(6-Methoxybiphenyl-3-ylmethylene)-1,3-

5 dihydropyrrolo[2,3-b]pyridin-2-one

4-Methoxy-3-phenylbenzaldehyde was condensed with 7-aza-2-oxindole to give 0.25 g of 3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one as a yellow-orange solid.

10 <sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.5 (s, 1H, CONH), 8.5-6.8 (m, 12H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>).

**Example 72:** 3-(3-Cyclohexyl-4-methoxybenzylidene)-1,3-

dihydropyrrolo[2,3-b]pyridin-2-one

15 3-Cyclohexyl-4-methoxybenzaldehyde was condensed with 7-aza-2-oxindole to give 0.25 g of 3-(3-Cyclohexyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one as a yellow-orange solid.

**Example 73:** 3-(2,3-Dihydrobenzofuran-5-ylmethylene)-1,3-

dihydropyrrolo[2,3-b]pyridin-2-one

20 2,3-Dihydro-5-formylbenzofuran (commercially available) was condensed with 7-aza-2-oxindole to give 0.2 g of 3-(2,3-dihydrobenzofuran-5-ylmethylene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one as a yellow-orange solid.

25 <sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.4 (s, 1H, CONH), 8.3-6.8 (m, 7H, aromatic, =CH-), 4.6 (t, 2H, OCH<sub>2</sub>), 3.2 (t, 2H, OCH<sub>2</sub>).

**Example 74:** 3-(3,5-Diisopropyl-4-methoxybenzylidene)-1,3-

dihydropyrrolo[2,3-b]pyridin-2-one

30 3,5-Diisopropyl-4-methoxybenzaldehyde was methylated and then formylated. The aldehyde was condensed with 7-aza-2-oxindole to give 0.2 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 11.5 (s, 1H, CONH), 8.0-6.8 (m, 7H,

WO 00/08202

174

PCT/US99/17845 ,

aromatic, =CH-), 3.7 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 2H, 2xCH), 1.2 (m, 12H, 4xCH<sub>3</sub>).

**Example 75: 5-Bromo-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one**

5        3-Isopropyl-4-methoxybenzaldehyde was condensed with 5-bromo-2-oxindole to give 0.3 g of 5-bromo-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

10        <sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.6 (s, 1H, CONH), 8.5, 8.4, 7.9, 7.9, 7.3, 7.1, 6.8 (m, 7H, aromatic, =CH-), 3.9 (m, 3H, OCH<sub>3</sub>), 3.3 (m, 1H, CH), 1.2 (d, 6H, 2xCH<sub>3</sub>).

**Example 76: 5-Bromo-3-(3-cyclopentyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one**

15        3-Cyclopentyl-4-methoxybenzaldehyde was condensed with 5-bromo-2-oxindole to give 0.3 g of 5-bromo-3-(3-cyclopentyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

20        <sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.6 (s, 1H, CONH), 8.6-6.7 (m, 7H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 1H, CH), 1.8 (m, 8H, 4xCH<sub>2</sub>).

**Example 77: 5-Chloro-3-(3-cyclopentyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one**

25        A suspension of 3.0 g of 4-methyl-2-oxindole was stirred in 50 ml of acetonitrile at room temperature while 3.3 g of N-chlorosuccinimide was added in portions. Trifluoroacetic acid (1 ml) was then added. The suspension was stirred at room temperature for 3 days during which time solids were always present. The solids were collected by vacuum filtration, washed with a small amount of cold acetone and dried overnight  
30        in a vacuum oven at 40°C to give 2.5 g (68 %) of 5-chloro-4-methyl-2-oxindole.

3-Cyclopentyl-4-methoxybenzaldehyde was condensed with 4-methyl-5-chloro-2-oxindole to give 0.3 g of 5-Chloro-3-(3-

WO 00/08202

175

PCT/US99/17845

cyclopentyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 78:** 5-Chloro-3-(6-methoxybiphenyl-3-ylmethylene)-4-methyl-1,3-dihydroindol-2-one

5 4-Methoxy-3-phenylbenzaldehyde was condensed with 4-methyl-5-chloro-2-oxindole to give 0.3 g of 5-chloro-3-(6-methoxybiphenyl-3-ylmethylene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 79:** 5-Chloro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-4-methyl-1,3-dihydroindol-2-one

5-Isopropyl-4-methoxy-2-methylbenzaldehyde was condensed with 4-methyl-5-chloro-2-oxindole to give 0.3 g of 5-chloro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

15 **Example 80:** 5-Chloro-3-(4-methoxy-3,5-dimethylbenzylidene)-4-methyl-1,3-dihydroindol-2-one

2,6-Dimethylphenol was methylated and then formylated. The aldehyde was condensed with 4-methyl-5-chloro-2-oxindole to give 0.3 g of 5-chloro-3-(4-methoxy-3,5-dimethylbenzylidene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 81:** 3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-trifluoromethyl-1,3-dihydroindol-2-one

25 3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 6-trifluoromethyl-2-oxindole to give 0.35 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-6-trifluoromethyl-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 82:** 6-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

30 3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 6-chloro-2-oxindole to give 0.3 g of 6-chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 83:** 3-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-

WO 00/08202

176

PCT/US99/17845

oxo-2,3-dihydro-1H-indol-5-yl]-propionic acid

Potassium cyanide (2.0 g) was added to 15 ml of dimethylsulfoxide and heated to 90°C. 5-chloroethyl-2-oxindole (3.0 g) dissolved in 5 ml of dimethylsulfoxide was added slowly with stirring, and the reaction was heated to 150°C and stirred for 2 hours. The mixture was cooled, poured into ice water and the precipitate which formed was collected by vacuum filtration, washed with water, and dried to give crude product. The crude material was chromatographed on silica gel (5% methanol in chloroform) to give 1.2 g (42%) of 5-cyanoethyl-2-oxindole.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.28 (s, br, NH), 7.12 (s, 1H, H-4), 7.07 (d, J = 8Hz, 1H), 6.74 (d, J = 8Hz, 1H), 3.43 (s, 2H, CH<sub>2</sub>), 2.71-2.82 (m, 4H, 2xCH<sub>2</sub>).

5-Cyanoethyl-2-oxindole (4.02 g) in 10 ml of water containing 25 ml of concentrated hydrochloric acid was refluxed for 4 hours. The mixture was cooled, water added and the resulting solid collected by vacuum filtration, washed with water and dried to give 1.9 g (44 % yield) of 5-carboxyethyl-2-oxindole as a yellow solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 12.00 (s, br, 1H, 5-CH<sub>2</sub>CH<sub>2</sub>COOH), 10.21 (s, 1H, NH-1), 7.05 (s, 1H, H-4), 6.99 (d, J = 9Hz, 1H, H-6), 6.69 (d, J = 9Hz, 1H, H-7), 3.40 (s, 2H, CH<sub>2</sub>-3), 2.74 (t, J = 7Hz, 2H, 5-CH<sub>2</sub>CH<sub>2</sub>COOH) and 2.46 (t, J = 7Hz, 2H, 5-CH<sub>2</sub>CH<sub>2</sub>COOH).

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-(2-carboxyethyl)-2-oxindole to give 0.35 g of 3-[3-(3,5-diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-5-yl]-propionic acid as a yellow-orange solid.

**Example 84:** 3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-methoxy-1,3-dihydroindol-2-one

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 6-methoxy-2-oxindole to give 0.3 g of 3-(3,5-diisopropyl-4-

WO 00/08202

177

PCT/US99/17845

methoxybenzylidene)-6-methoxy-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 85:     5-Butyl-3-(3,5-diisopropyl-4-methoxy-benzylidene)-1,3-dihydroindol-2-one**

5       To 15 g of aluminum chloride suspended in 30 ml of 1,2-dichloroethane in an ice bath was added 7.5 g of 2-oxindole and then 12 g of butyryl chloride. The resulting suspension was stirred at 50 °C overnight. The mixture was poured into ice water and extracted 3 times with ethyl acetate. The  
10 combined ethyl acetate layers were washed with brine, dried over sodium sulfate, and concentrated to dryness to give a brown solid. The solid was chromatographed on silica gel (50 % ethyl acetate in hexane) to give 4.3 g (38%) of 5-butanoyl-2-oxindole as a yellow solid.

15       <sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.67 (s, br, NH), 7.84 (d, J = 8.39Hz, 1H), s, 1H, H-4), 6.88 (d, J = 8.39Hz, 1H), 3.53 (s, 2H, CH<sub>2</sub>), 2.88 (t, J = 7.16Hz, 2H, CH<sub>2</sub>), 1.58-1.64 (m, 2H), 0.9 (t, J = 7.58Hz, 3H, CH<sub>3</sub>).

      Triethylsilane (2.3 g) was added to 2 g of 5-butanoyl-2-oxindole in 20 ml of trifluoroacetic acid at room temperature  
20 and the solution stirred for 3 hours. The reaction was poured into ice water to give a red oil which solidified after standing. The solid was collected by vacuum filtration, washed with water and hexane and dried to give 1.7 g (91%  
25 yield) of 5-butyl-2-oxindole as an off-white solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.19 (s, br, NH), 7.01 (s, 1H, H-4), 6.95 (d, J = 8Hz, 1H), 6.69 (d, J = 8Hz, 1H), 3.4 (s, 2H, CH<sub>2</sub>), 2.49 (t, J = 8Hz, 2H, CH<sub>2</sub>), 1.48-1.52 (m, 2H), 1.25-1.31 (m, 2H), 0.88 (t, J = 7Hz, 3H).

30       3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-butyl-2-oxindole to give 0.3 g of 5-butyl-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.



WO 00/08202

178

PCT/US99/17845 ,

**Example 86:     3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid**

Trimethylsilyldiazomethane in hexane (2 M) was added dropwise to the solution of 2-chloro-3-carboxynitrobenzene  
5 (2.01 g) in methanol (20 ml) at room temperature until no further gas evolution occurred. The excess trimethylsilyldiazomethane was quenched with acetic acid. The reaction mixture was dried by rotary pump and the residue was further  
10 dried in oven for overnight. The product (2-chloro-3-methoxycarbonyl-nitrobenzene) was pure enough for the following reaction.

Dimethyl malonate (6.0 ml) was added to an ice-cold suspension of sodium hydride (2.1 g) in dimethylsulfoxide (15 ml). The reaction mixture was heated to 100°C, stirred for 1  
15 hour and then cooled to room temperature. 2-Chloro-3-methoxycarbonylnitrobenzene (2.15 g) was added in one portion and the mixture was stirred at 100°C for 1.5 hour. The reaction mixture was then cooled to room temperature and  
20 poured into ice water, acidified to pH 5, and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous sodium sulfate and concentrated to give 3.0 g of the dimethyl 2-methoxycarbonyl-6-nitrophenylmalonate.

Dimethyl 2-methoxycarbonyl-6-nitrophenylmalonate (3.0 g) was refluxed in 50 ml of 6 N hydrochloric acid overnight. The  
25 mixture was concentrated to dryness and refluxed for 2 hours with 1.1 g of tin(II) chloride in 20 ml of ethanol. The mixture was filtered through Celite, concentrated and chromatographed on silica gel (ethyl acetate:hexane:acetic acid) to give 0.65 g (37 % yield) of 4-carboxy-2-oxindole as a  
30 white solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 12.96 (s, br, 1H, COOH), 10.74 (s, br, 1H, NH), 7.53 (d, J = 8Hz, 1H), 7.39 (t, J = 8Hz, 1H, H-6), 7.12 (d, J = 8Hz, 1H, H-7), 3.67 (s, 2H, H-3).

WO 00/08202

179

PCT/US99/17845 ,

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 4-carboxy-2-oxindole to give 0.3 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid as a yellow-orange solid.

5 **Example 87:** 3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-(3-methoxyphenyl)-1,3-dihydroindol-2-one

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 6-(3-methoxyphenyl)-2-oxindole to give 0.4 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-6-(3-methoxyphenyl)-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 88:** 7-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-5-methyl-1,3-dihydroindol-2-one

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-methyl-7-chloro-2-oxindole to give 0.35 g of 7-chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-5-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 89:** [3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-5-yl]-carbamic acid tert-butyl ester

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-tert-butyloxy-carbonylamino-2-oxindole to give 0.4 g of [3-(3,5-diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-5-yl]-carbamic acid tert-butyl ester as a yellow-orange solid.

**Example 90:** 5-Chloro-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

3-Isopropyl-4-methoxybenzaldehyde was condensed with 5-chloro-2-oxindole to give 0.3 g of 5-chloro-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

30 <sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.6 (s, 1H, CONH), 8.5, 8.4, 7.9, 7.8, 7.2, 7.1, 6.8 (m, 7H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 3.3 (m, 1H, CH), 1.2 (d, 6H, 2xCH<sub>3</sub>).

**Example 91:** 5-Chloro-3-(3-cyclopentyl-4-methoxy-

WO 00/08202

180

PCT/US99/17845

**benzylidene)-1,3-dihydroindol-2-one**

3-Cyclopentyl-4-methoxybenzaldehyde was condensed with 5-chloro-2-oxindole to give 0.3 g of 5-chloro-3-(3-cyclopentyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.6 (s, 1H, CONH), 8.5, 8.3, 7.9, 7.8, 7.2, 7.1, 6.8 (m, 7H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 3.3 (m, 1H, CH), 1.8 (m, 8H, 4xCH<sub>2</sub>).

**Example 92: 3-(6-Methoxybiphenyl-3-ylmethylene)-4-methyl-1,3-dihydroindol-2-one**

4-Methoxy-3-phenylbenzaldehyde was condensed with 4-methyl-2-oxindole to give 0.3 g of 3-(6-methoxybiphenyl-3-ylmethylene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.5 (s, 1H, CONH), 8.3-6.7 (m, 12H, aromatic, =CH-), 3.9 (s, 3H, OCH<sub>3</sub>), 2.6 (s, 3H, CH<sub>3</sub>).

**Example 93: 3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-4-methyl-1,3-dihydroindol-2-one**

5-Isopropyl-4-methoxy-2-methylbenzaldehyde was condensed with 4-methyl-2-oxindole to give 0.3 g of 3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.3 (s, 1H, CONH), 8.2, 7.8, 7.1, 6.8, 6.8, 6.7 (m, 6H, aromatic, =CH-), 3.8 (s, 3H, OCH<sub>3</sub>), 3.2 (m, 1H, CH), 2.5 (s, 3H, CH<sub>3</sub>), 2.4 (s, 3H, CH<sub>3</sub>), 1.2 (d, 6H, 2xCH<sub>3</sub>).

**Example 94: 5-Bromo-3-(2,3-dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one**

2,3-Dihydro-5-formylbenzofuran (commercially available) was condensed with 5-bromo-2-oxindole to give 0.3 g of 5-bromo-3-(2,3-dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one as a yellow-orange solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ: 10.64 (s, 1H, CONH), 7.71 (d, 1H),

WO 00/08202

181

PCT/US99/17845

7.62 (vinyl, 1H), 7.61 (m, 1H), 7.52 (m, 1H), 7.37 (dd, 1H), 6.93 (d, 1H), 6.984 (d, 1H), 4.64 (t, 2H, CH<sub>2</sub>), 3.22 (t, 2H, CH<sub>2</sub>).

**Example 95:** 5-Chloro-3-(3-isopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

3-Isopropyl-4-methoxybenzaldehyde was condensed with 4-methyl-5-chloro-2-oxindole to give 0.3 g of 5-chloro-3-(3-isopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 96:** 5-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 4-methyl-5-chloro-2-oxindole to give 0.4 g of 5-chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 97:** 5-Chloro-3-(2,2-dimethylchroman-6-ylmethylene)-4-methyl-1,3-dihydroindol-2-one

2,2-Dimethyl-6-formylchromane (commercially available) was condensed with 4-methyl-5-chloro-2-oxindole to give 0.3 g of 5-chloro-3-(2,2-dimethyl-chroman-6-ylmethylene)-4-methyl-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 98:** 3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid

2-Oxindole (6.7 g) was added to a suspension of aluminum chloride (23 g) in dichloroethane (30 ml) cooled in an ice bath. Chloroacetyl chloride (11.3 g) was slowly added and hydrogen chloride gas was evolved. After ten minutes of stirring, the reaction was warmed to 40 - 50°C for 1.5 hours. The mixture was cooled to room temperature and poured into ice water. The precipitate was collected by vacuum filtration, washed with water and dried under vacuum to give 10.3 g (98%) of 5-chloroacetyl-2-oxindole as an off-white solid.

A suspension of 9.3 g of 5-chloroacetyl-2-oxindole was

WO 00/08202

182

PCT/US99/17845 ,

stirred in 90 ml pyridine at 80 - 90°C for 3 hours then cooled to room temperature. The precipitate was collected by vacuum filtration and washed with 20 ml of ethanol. The solid was dissolved in 90 ml of 2.5 N sodium hydroxide and stirred at 70  
5 to 80°C for 3 hours. The mixture was cooled to room temperature and acidified to pH 2 with 0.5 N hydrochloric acid. The precipitate was collected by vacuum filtration and washed thoroughly with water to give crude 5-carboxy-2-oxindole as a dark brown solid. After standing overnight the  
10 filtrate yielded 2 g of 5-carboxy-2-oxindole as a yellow solid. The crude dark brown product was dissolved in hot methanol, the insoluble material removed by filtration and the filtrate concentrated to give 5.6 g of 5-carboxy-2-oxindole as a brown solid. The combined yield was 97%.

15 <sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 2.56 (s, br, 1H, COOH-5), 10.70 (s, 1H, NH-1), 7.82 (dd, J = 2, 8Hz, 1H, H-6), 7.74 (s, br, 1H, H-4), 6.87 (d, J = 8Hz, 1H, H-7), and 3.53 (s, 2H, CH<sub>2</sub>-3).

MS m/z 178 [M+1]<sup>+</sup>.

20 3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-carboxy-2-oxindole to give 0.4 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid as a yellow-orange solid.

**Example 99:** 3-(3,5-Diisopropyl-4-methoxybenzylidene)-5,6-dimethoxy-1,3-dihydroindol-2-one  
25

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5,6-dimethoxy-2-oxindole to give 0.4 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-5,6-dimethoxy-1,3-dihydroindol-2-one as a yellow-orange solid.

30 **Example 100:** N-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-methanesulfonamide

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 6-methylsulfonylamino-2-oxindole to give 0.4 g of N-[3-(3,5-

WO 00/08202

183

PCT/US99/17845

diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-methanesulfonamide as a yellow-orange solid.

**Example 101:** N-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-benzamide

5        3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 6-benzamido-2-oxindole to give 0.4 g of N-[3-(3,5-diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-benzamide as a yellow-orange solid.

**Example 102:** 3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-(3-ethoxyphenyl)-1,3-dihydroindol-2-one

10        Tetrakis(triphenylphosphine)palladium (0.8 g) was added to a mixture of 4.2 g of 3-ethoxyphenylboronic acid, 5.0 g of 5-bromo-2-fluoronitrobenzene and 22 ml of 2 M sodium carbonate solution in 50 ml of toluene and 50 ml of ethanol. The  
15        mixture was heated to reflux for 2 hours. The cooled mixture was concentrated, water was added and the mixture was extracted twice with ethyl acetate. The ethyl acetate layer was washed with water, brine, dried, and concentrated. The residue was chromatographed on silica gel (5 % ethyl acetate  
20        in hexane) to give 5.3 g (90 % yield) of crude 4-fluoro-3'-ethoxy-3-nitrobiphenyl as a yellow oil.

      Dimethyl malonate (11.4 ml) was added dropwise to 4.0 g of sodium hydride suspended in 20 ml of dimethylsulfoxide. The mixture was heated to 100°C for 10 minutes and cooled to  
25        room temperature. Crude 4-fluoro-3'-ethoxy-3-nitrobiphenyl (5.3 g) in 25 ml of dimethylsulfoxide was added and the mixture was stirred at 100°C for 2 hours. The reaction mixture was cooled and quenched with 300 ml of saturated ammonium chloride solution and extracted three times with  
30        ethyl acetate. The extracts were combined, washed with water and brine, dried over anhydrous sodium sulfate and concentrated to give crude dimethyl 3'-ethoxy-3-nitrobiphenyl-4-malonate as a yellow oil.

WO 00/08202

184

PCT/US99/17845 ,

Crude dimethyl 3'-ethoxy-3-nitrobiphenyl-4-malonate in 60 ml of 6 N hydrochloric acid was stirred at 100°C for 4 days and then cooled. The precipitate which formed was collected by filtration, washed with water and hexane, and dried to give  
5 4.7 g (77% based on 5-bromo-2-fluoronitrobenzene) of crude 3'-ethoxy-3-nitrobiphenyl-4-acetic acid as a light tan solid.

Iron chips (2.4 g) were added in one portion to 4.6 g of 3'-ethoxy-3-nitrobiphenyl-4-acetic acid in glacial acetic acid (40 ml). The mixture was refluxed for 2 hours. The reaction  
10 mixture was cooled and concentrated to dryness. The residue was treated repeatedly with ethyl acetate and filtered to remove insolubles. The filtrate was washed twice with 1 N hydrochloric acid and brine, dried over anhydrous sodium sulfate and concentrated to give 3.5 g (91%) of 6-(3-  
15 ethoxyphenyl)-2-oxindole as a light brown solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.4 (s, br, 1H, NH), 7.33 (t, J = 8Hz, 1H, H-3'), 7.35 (d, J = 8Hz, 1H), 7.19 (dd, J = 1, 8Hz, 1H), 7.13 (d, J = 8Hz, 1H), 7.07-7.08 (m, 1H), 7.0 (s, br, 1H), 6.9 (dd, J = 3, 8Hz, 1H), 4.08 (q, J = 7Hz, 2H, O  
20 CH<sub>2</sub>CH<sub>3</sub>), 3.49 (s, 2H, CH<sub>2</sub>), 1.34 (t, J = 7Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS m/z 254.2 [M+1]<sup>+</sup>.

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 6-(3-ethoxyphenyl)-2-oxindole to give 0.4 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-6-(3-ethoxyphenyl)-1,3-  
25 dihydroindol-2-one as a yellow-orange solid.

**Example 103: 3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-phenyl-1,3-dihydroindol-2-one**

Tetrakis(triphenylphosphine)palladium (0.8 g) was added  
30 to a mixture of 3.1 g of benzeneboronic acid, 5 g of 5-bromo-2-fluoronitrobenzene and 22 ml of 2M aqueous sodium carbonate in 50 ml of toluene and 50 ml of ethanol. The mixture was refluxed for 2 hours, concentrated, and the residue extracted

WO 00/08202

185

PCT/US99/17845

twice with ethyl acetate. The ethyl acetate layer was washed with water and brine, dried, and concentrated to give a yellow oil. The oil was chromatographed on silica gel (5% ethyl acetate in hexane) to give 4.75 g (96%) of 4-fluoro-3-nitrobiphenyl as a yellow oil.

Dimethyl malonate (10 ml) in 25 ml of dimethylsulfoxide was added dropwise to 3.5 g of sodium hydride suspended in 25 ml of dimethylsulfoxide and the mixture stirred at 100°C for 10 minutes. The mixture was cooled to room temperature and 4.7 g of 4-fluoro-3-nitrobiphenyl in 25 ml of dimethylsulfoxide was added. The mixture was stirred at 100°C for 2 hours, cooled and quenched with 300 ml of saturated ammonium chloride solution. The mixture was extracted three times with ethyl acetate and the combined organic layers washed with water and brine and evaporated to give a yellow oil, crude dimethyl-3-nitrobiphenyl-4-malonate.

Crude dimethyl-3-nitrobiphenyl-4-malonate was refluxed in 30 ml of 6 N hydrochloric acid for 24 hours. The precipitate was collected by filtration, washed with water and dried to give 4.5 g (80% based on 4-fluoro-3-nitrobiphenyl) of 3-nitrobiphenyl-4-acetic acid as a cream colored solid.

Iron chips (2.6 g) were added all at once to 4.5 g of 3-nitrobiphenyl-4-acetic acid in 40 ml of acetic acid. The mixture was refluxed for 2 hours, concentrated to dryness and the residue taken up in ethyl acetate. The solids were removed by filtration and the filtrate washed twice with 1 N hydrochloric acid and brine and dried over anhydrous sodium sulfate. The filtrate was concentrated to give 3.4 g (93 % yield) of 6-phenyl-2-oxindole as a light brown solid.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ: 10.4 (s, br, 1H, NH-1), 7.57-7.6 (m, 2H), 7.42-7.46 (m, 2H), 7.32-7.37 (m, 1H), 7.27 (d, J = 8, 1H, H-4), 7.19 (dd, J = 2 and 8Hz, 1H, H-5), 7.01 (d, J = 2Hz, 1H, H-7), 3.49 (s, 2H, CH<sub>2</sub>).



WO 00/08202

186

PCT/US99/17845

MS m/z 210 [M+1]<sup>+</sup>.

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 6-phenyl-2-oxindole to give 0.4 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-6-phenyl-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 104:** 3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one

3,5-Diisopropyl-4-methoxybenzaldehyde was condensed with 5-fluoro-2-oxindole to give 0.3 g of 3-(3,5-diisopropyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 105:** 5-Fluoro-3-(4-methoxy-3,5-dimethylbenzylidene)-1,3-dihydroindol-2-one

3,5-Dimethyl-4-methoxybenzaldehyde was condensed with 5-fluoro-2-oxindole to give 0.3 g of 5-fluoro-3-(4-methoxy-3,5-dimethylbenzylidene)-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 106:** 3-(2,2-Dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one

2,2-Dimethyl-6-formylchromane (commercially available) was condensed with 2-oxindole to give 0.3 g of 3-(2,2-dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one as a yellow-orange solid.

**Example 107:** 3-{2-[6-(4-Fluorophenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-5-methyl-1H-pyrrol-3-yl}propionic acid

6-(4-Fluorophenyl)-1,3-dihydroindol-2-one (70 mg, 0.31 mmol) was condensed with 3-(2-formyl-5-methyl-1H-pyrrol-3-yl)-propionic acid (56 mg) to give the title compound.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 13.38 (s, br, 1H, COOH), 12.08 (s, br, 1H, NH), 10.87 (s, br, 1H, NH), 7.79 (d, J = 7.9 Hz, 1H), 7.63-7.69 (m, 3H), 7.23-7.29 (m, 3H), 7.07 (d, J = 1.4 Hz, 1H), 6.06 (d, J = 2.2 Hz, 1H), 2.96 (t, J = 7.2 Hz, 2H, CH<sub>2</sub>), 2.52 (t, J = 7.2 Hz, 2H, CH<sub>2</sub>), 2.32 (s, 3H, CH<sub>3</sub>).

WO 00/08202

187

PCT/US99/17845 ,

MS-EI  $m/z$  390  $[M^+]$ .

Example 108: 4-(2-Carboxyethyl)-5-[6-(4-fluorophenyl)-2-oxo-1,2-dihydro-indol-3-ylidenemethyl]-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester

- 5 6-(4-Fluorophenyl)-1,3-dihydroindol-2-one (60 mg, 0.26 mmol) was condensed with 4-(2-carboxyethyl)-5-formyl-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester (65 mg), prepared by formylation of 4-(2-carboxyethyl)-3-ethoxycarbonyl-2-methylpyrrole (Bulter, A.R., and George, S.D. (1993) 10 Tetrahedron 49: 7017-7026), to give the title compound.

$^1\text{H}$ NMR (360 MHz, DMSO- $d_6$ )  $\delta$  13.85 (s, br, 1H, NH), 12.03 (s, br, 1H, COOH), 11.05 (s, br, 1H, NH), 7.83 (d,  $J$  = 7.9 Hz, 1H), 7.76 (s, 1H, H-vinyl), 7.66-7.70 (m, 2H), 7.25-7.30 (m, 3H), 7.08 (d,  $J$  = 1.4 Hz, 1H), 4.22 (q,  $J$  = 7.0 Hz, 2H, 15  $\text{OCH}_2\text{CH}_3$ ), 3.20 (t,  $J$  = 7.6 Hz, 2H,  $\text{CH}_2$ ), 2.54 (s, 3H,  $\text{CH}_3$ ), 2.45 (m, 2H,  $\text{CH}_2$ ), 4.22 (t,  $J$  = 7.0 Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ),

MS-EI  $m/z$  462  $[M^+]$ .

Example 109: 3-{2-[6-(2-Methoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-5-methyl-1H-pyrrol-3-yl}-propionic acid

- 20 6-(2-Methoxyphenyl)-1,3-dihydroindol-2-one (71 mg, 0.3 mmol) was condensed with 3-(2-formyl-5-methyl-1H-pyrrol-3-yl)-propionic acid (54 mg) to give the title compound.

$^1\text{H}$ NMR (300 MHz, DMSO- $d_6$ )  $\delta$  13.33 (s, br, 1H, NH), 12.07 (s, br, 1H, COOH), 10.74 (s, br, 1H, NH), 7.68 (d,  $J$  = 7.8 Hz, 1H), 7.56 (s, 1H, H-vinyl), 7.25-7.28 (m, 2H), 6.94-7.05 (m, 4H), 5.99 (br s, 1H), 3.70 (s, 3H,  $\text{OCH}_3$ ), 2.89 (m, 2H,  $\text{CH}_2$ ), 2.45 (m, 2H,  $\text{CH}_2$ ), 2.26 (s, 3H,  $\text{CH}_3$ ).

MS-EI  $m/z$  402  $[M^+]$ .

- 30 Example 110: 4-(2-Carboxyethyl)-5-[6-(2-methoxyphenyl)-2-oxo-1,2-dihydro-indol-3-ylidenemethyl]-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester

$^1\text{H}$ NMR (300 MHz, DMSO- $d_6$ )  $\delta$  13.88 (s, br, 1H, NH), 12.07

WO 00/08202

188

PCT/US99/17845 ,

(s, br, 1H, COOH), 10.99 (s, br, 1H, NH), 7.77 (d,  $J = 7.8$  Hz, 1H), 7.74 (s, 1H, H-vinyl), 7.28-7.36 (m, 2H), 7.07-7.1 (m, 2H), 6.99-7.04 (m, 2H), 4.21 (q,  $J = 7.0$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.76 (s, 3H,  $\text{OCH}_3$ ), 3.19 (m, 2H,  $\text{CH}_2$ ), 2.54 (s, 3H,  $\text{CH}_3$ ), 2.50 (m, 2H,  $\text{CH}_2$ ), 1.04 (t,  $J = 7.0$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  474 [ $M^+$ ].

**Example 111:** 3-[2-(5-Chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-5-methyl-1H-pyrrol-3-yl]propionic acid

5-Chloro-1,3-dihydroindol-2-one (61 mg, 0.36 mmol) was condensed with 3-(2-formyl-5-methyl-1H-pyrrol-3-yl)-propionic acid (66 mg) to give the title compound.

MS-EI  $m/z$  300 and 302 [ $M-1$  and  $M+1$ ].

**Example 112:** 4-(2-Carboxyethyl)-5-(5-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester

$^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.88 (s, br, 1H, NH), 12.04 (s, br, 1H, COOH), 11.08 (s, br, 1H, NH), 7.93 (d,  $J = 2.0$  Hz, 1H), 7.83 (s, 1H, H-vinyl), 7.15 (dd,  $J = 2.0, 8.3$  Hz, 1H), 6.87 (d,  $J = 8.3$  Hz, 1H), 4.20 (q,  $J = 7.1$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.21 (t,  $J = 7.6$  Hz, 2H,  $\text{CH}_2$ ), 2.53 (s, 3H,  $\text{CH}_3$ ), 2.45 (t,  $J = 7.6$  Hz, 2H,  $\text{CH}_2$ ), 1.28 (t,  $J = 7.1$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  402 and 404 [ $M-1$  and  $M+1$ ].

**Example 117:** 4-(2-Carboxyethyl)-3-methyl-5-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester

Oxindole (0.25g, 1.88 mmol) was condensed with 4-(2-carboxyethyl)-5-formyl-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester (0.5 g, 1.97 mmol) to give 0.45 g (65%) of the title compound.

$^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.85 (s, br, 1H, NH), 12.12 (s, br, 1H, COOH), 10.99 (s, br, 1H, NH), 7.78 (d,  $J = 7.6$  Hz, 1H), 7.70 (s, 1H, H-vinyl), 7.17 (m 1H), 6.99 (m, 1H), 4.25 (q,  $J = 7.1$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 2.93 (t,  $J = 7.4$  Hz, 2H,  $\text{CH}_2$ ),

WO 00/08202

189

PCT/US99/17845

2.38 (t,  $J = 7.4$  Hz, 2H,  $\text{CH}_2$ ), 2.24 (s, 3H,  $\text{CH}_3$ ), 1.29 (t,  $J = 7.1$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  368 [ $\text{M}^+$ ].

Example 118: 2-Methyl-5-(5-methylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-(3-morpholin-4-ylpropyl)-1H-pyrrole-3-carboxylic acid ethyl ester

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide was stirred with 5-formyl-2-methyl-4-(3-morpholin-4-ylpropyl)-1H-pyrrole-3-carboxylic acid ethyl ester overnight at room temperature to give the title compound.

$^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.86 (s, br, 1H, NH), 11.39 (s, br, 1H, NH), 8.16 (d,  $J = 1.6$  Hz, 1H, H-4), 7.82 (s, 1H, H-vinyl), 7.57 (dd,  $J = 1.6$  & 6.9 Hz, 1H, H-6), 7.21 (m, 1H,  $\text{CH}_3\text{NHSO}_2$ ), 7.07 (d,  $J = 6.9$  Hz, 1H, H-7), 4.21 (q,  $J = 7.0$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.52 (m, 4H,  $2\times\text{CH}_2$ ), 3.03 (m, 2H,  $\text{CH}_2$ ), 2.55 (s, 3H,  $\text{CH}_3$ ), 2.40 (d,  $J = 5.0$  Hz, 3H,  $\text{CH}_3\text{NHSO}_2$ ), 2.25 (m, 6H,  $3\times\text{CH}_2$ ), 1.69 (m, 2H,  $\text{CH}_2$ ), 1.29 (t,  $J = 7.0$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  516 [ $\text{M}^+$ ].

Example 119: 2-Methyl-4-[3-(4-methylpiperazin-1-yl)-propyl]-5-(5-methylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid ethyl ester

$^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.88 (s, br, 1H, NH), 11.38 (s, br, 1H, NH), 8.16 (d,  $J = 1.6$  Hz, 1H, H-4), 7.79 (s, 1H, H-vinyl), 7.57 (dd,  $J = 1.6$  & 6.6 Hz, 1H, H-6), 7.21 (m, 1H,  $\text{CH}_3\text{NHSO}_2$ ), 7.05 (d,  $J = 6.6$  Hz, 1H, H-7), 4.21 (q,  $J = 7.0$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.01 (m, 2H,  $\text{CH}_2$ ), 2.55 (s, 3H,  $\text{CH}_3$ ), 2.40 (d,  $J = 5.4$  Hz, 3H,  $\text{CH}_3\text{NHSO}_2$ ), 2.57 (m, 10H,  $5\times\text{CH}_2$ ), 2.09 (s, 3H,  $\text{CH}_3\text{N}$ ), 1.67 (m, 2H,  $\text{CH}_2$ ), 1.29 (t,  $J = 7.0$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  529 [ $\text{M}^+$ ].

Example 120: 4-(3-Dimethylaminopropyl)-5-(5-methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester

$^1\text{H}$ NMR (360 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.95 (s, br, 1H, NH), 10.76

WO 00/08202

190

PCT/US99/17845

(s, br, 1H, NH), 7.70 (s, 1H, H-vinyl), 7.31 (d,  $J = 2.2$  Hz, 1H, H-4), 6.78 (d,  $J = 8.3$  Hz, 1H, H-7), 6.73 (dd,  $J = 2.2$  & 8.3 Hz, 1H, H-6), 4.21 (q,  $J = 7.0$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.76 (s, 3H,  $\text{OCH}_3$ ), 2.98 (m, 2H,  $\text{CH}_2$ ), 2.53 (s, 3H,  $\text{CH}_3$ ), 2.19 (m, 2H, 5  $\text{CH}_2$ ), 2.11 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 1.64 (m, 2H,  $\text{CH}_2$ ), 1.30 (t,  $J = 7.0$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  411 [ $\text{M}^+$ ].

**Example 121:** 5-(5-Methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-2-methyl-4-(3-morpholin-4-ylpropyl)-1H-pyrrole-3-carboxylic acid ethyl ester

$^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  14.0 (s, br, 1H, NH), 10.77 (s, br, 1H, NH), 7.65 (s, 1H, H-vinyl), 7.40 (m, 1H), 6.76 (m, 2H), 4.2 (q,  $J = 7.0$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.76 (s, 3H,  $\text{OCH}_3$ ), 3.54 (m, 4H,  $2 \times \text{CH}_2$ ), 2.99 (m, 2H,  $\text{CH}_2$ ), 2.52 (s, 3H,  $\text{CH}_3$ ), 2.25 (m, 15 6H,  $3 \times \text{CH}_2$ ), 1.67 (m, 2H,  $\text{CH}_2$ ), 1.29 (t,  $J = 7.0$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  453 [ $\text{M}^+$ ].

**Example 122:** 5-(5-Methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-2-methyl-4-[3-(4-methylpiperazin-1-yl)-propyl]-1H-pyrrole-3-carboxylic acid ethyl ester

$^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.99 (s, br, 1H, NH), 10.78 (s, br, 1H, NH), 7.63 (s, 1H, H-vinyl), 7.39 (d,  $J = 1.8$  Hz, 1H, H-4), 6.76 (m, 2H), 4.2 (q,  $J = 7.1$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.77 (s, 3H,  $\text{OCH}_3$ ), 2.97 (m, 2H,  $\text{CH}_2$ ), 2.52 (s, 3H,  $\text{CH}_3$ ), 2.25 (m, 10H, 25  $5 \times \text{CH}_2$ ), 2.08 (s, 3H,  $\text{NCH}_3$ ), 1.66 (m, 2H,  $\text{CH}_2$ ), 1.29 (t,  $J = 7.1$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  466 [ $\text{M}^+$ ].

**Example 123:** 2-Methyl-4-(3-morpholin-4-ylpropyl)-5-(2-oxo-5-sulfamoyl-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid ethyl ester

$^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.85 (s, br, 1H, NH), 11.33 (s, br, 1H, NH), 8.19 (d,  $J = 1.8$  Hz, 1H, H-4), 7.75 (s, 1H, H-vinyl), 7.63 (dd,  $J = 1.8$  & 8.3 Hz, 1H, H-6), 7.16 (s, 2H,

WO 00/08202

191

PCT/US99/17845 ,

H<sub>2</sub>NSO<sub>2</sub>), 7.02 (d, *J* = 8.3 Hz, 1H, H-7), 4.21 (q, *J* = 7.3 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.53 (m, 4H, 2xCH<sub>2</sub>), 3.01 (m, 2H, CH<sub>2</sub>), 2.55 (s, 3H, CH<sub>3</sub>), 2.28 (m, 6H, 3xCH<sub>2</sub>), 1.68 (m, 2H, CH<sub>2</sub>), 1.29 (t, *J* = 7.3 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

5 MS-EI *m/z* 502 [M<sup>+</sup>].

**Example 124: 2-Methyl-4-[3-(4-methylpiperazin-1-yl)-propyl]-5-(2-oxo-5-sulfamoyl-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid ethyl ester**

<sup>1</sup>HNMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 13.85 (s, br, 1H, NH), 11.31 (s, br, 1H, NH), 8.18 (d, *J* = 1.5 Hz, 1H, H-4), 7.74 (s, 1H, H-vinyl), 7.63 (dd, *J* = 1.5 & 8.1 Hz, 1H, H-6), 7.15 (s, 2H, H<sub>2</sub>NSO<sub>2</sub>), 7.01 (d, *J* = 8.1 Hz, 1H, H-7), 4.21 (q, *J* = 7.1 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.0 (m, 2H, CH<sub>2</sub>), 2.54 (s, 3H, CH<sub>3</sub>), 2.26 (m, 10H, 5xCH<sub>2</sub>), 2.09 (s, 3H, NCH<sub>3</sub>), 1.67 (m, 2H, CH<sub>2</sub>), 1.29 (t, *J* = 7.1 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

15 MS-EI *m/z* 515 [M<sup>+</sup>].

**Example 125: 3-[4-Ethoxycarbonyl-5-methyl-3-(3-morpholin-4-ylpropyl)-1H-pyrrol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid**

<sup>1</sup>HNMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 13.88 (s, br, 1H, NH), 11.03 (s, br, 1H, COOH), 8.17 (s, 1H, H-4), 7.76 (d, *J* = 8.4 Hz, 1H, H-6), 7.59 (s, 1H, H-vinyl), 6.80 (d, *J* = 8.4 Hz, 1H, H-7), 4.21 (q, *J* = 7.2 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.54 (m, 4H, 2xCH<sub>2</sub>), 2.98 (m, 2H, CH<sub>2</sub>), 2.49 (s, 3H, CH<sub>3</sub>), 2.29 (m, 6H, 3xCH<sub>2</sub>), 1.64 (m, 2H, CH<sub>2</sub>), 1.29 (t, *J* = 7.2 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

25 MS-EI *m/z* 467 [M<sup>+</sup>].

**Example 126: 2-Methyl-5-(5-methylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-(3-pyrrolidin-1-ylpropyl)-1H-pyrrole-3-carboxylic acid ethyl ester**

<sup>1</sup>HNMR (360 MHz, DMSO-*d*<sub>6</sub>) δ 13.84 (s, 1H, NH), 11.28 (br s, 1H, NH), 8.06 (d, *J* = 2.1 Hz, 1H), 7.83 (s, 1H, H-vinyl), 7.57 (dd, *J* = 2.1 & 8.3 Hz, 1H), 7.20 (m, 1H, HNCH<sub>3</sub>), 7.06 (d, *J* = 8.3 Hz, 1H), 4.22 (q, *J* = 7.0 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.04 (m, 2H,

WO 00/08202

192

PCT/US99/17845

$\text{CH}_2$ ), 2.55 (s, 3H,  $\text{CH}_3$ ), 2.40 (d,  $J = 4.8$  Hz, 3H,  $\text{HNCH}_3$ ), 2.3-2.36 (m, 6H,  $3\times\text{CH}_2$ ), 1.66-1.71 (m, 6H,  $3\times\text{CH}_2$ ), 1.30 (t,  $J = 7.0$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  500 [ $\text{M}^+$ ].

- 5 **Example 127:** 5-(5-Methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-2-methyl-4-(3-pyrrolidin-1-ylpropyl)-1H-pyrrole-3-carboxylic acid ethyl ester

$^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.96 (s, 1H, NH), 10.75 (br s, 1H, NH), 7.65 (s, 1H, H-vinyl), 7.28 (d,  $J = 2.2$  Hz, 1H), 6.71-6.79 (m, 2H), 4.20 (q,  $J = 7.1$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.75 (s, 3H,  $\text{OCH}_3$ ), 3.0 (m, 2H,  $\text{CH}_2$ ), 2.52 (s, 3H,  $\text{CH}_3$ ), 2.26-2.35 (m, 6H,  $3\times\text{CH}_2$ ), 1.65-1.71 (m, 6H,  $3\times\text{CH}_2$ ), 1.29 (t,  $J = 7.1$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  437 [ $\text{M}^+$ ].

- 15 **Example 128:** 3-[4-Ethoxycarbonyl-5-methyl-3-(3-pyrrolidin-1-ylpropyl)-1H-pyrrol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid

$^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.80 (s, 1H, NH), 11.26 (br s, 1H, NH), 8.2 (s, 1H), 7.77 (m, 2H), 6.92 (d,  $J = 8.1$  Hz, 1H), 4.23 (q,  $J = 6.9$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.03 (m, 2H,  $\text{CH}_2$ ), 2.54 (s, 3H,  $\text{CH}_3$ ), 2.40 (m, 6H,  $3\times\text{CH}_2$ ), 1.68 (m, 6H,  $3\times\text{CH}_2$ ), 1.29 (t,  $J = 6.9$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  451 [ $\text{M}^+$ ].

- 25 **Example 130:** 2-Methyl-5-(2-oxo-5-sulfamoyl-1,2-dihydroindol-3-ylidenemethyl)-4-(3-pyrrolidin-1-ylpropyl)-1H-pyrrole-3-carboxylic acid ethyl ester

$^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.98 (s, 1H, NH), 11.30 (br s, 1H, NH), 8.10 (d,  $J = 1.9$  Hz, 1H), 7.79 (s, 1H, H-vinyl), 7.63 (dd,  $J = 1.9$  & 8.2 Hz, 1H), 7.16 (br s, 2H,  $\text{H}_2\text{NSO}_2$ ), 7.02 (d,  $J = 8.2$  Hz, 1H), 4.21 (q,  $J = 6.9$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.03 (m, 2H,  $\text{CH}_2$ ), 2.54 (s, 3H,  $\text{CH}_3$ ), 2.33 (m, 6H,  $3\times\text{CH}_2$ ), 1.68 (m, 6H,  $3\times\text{CH}_2$ ), 1.29 (t,  $J = 6.9$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

MS-EI  $m/z$  486 [ $\text{M}^+$ ].

WO 00/08202

193

PCT/US99/17845

**Example 131: 5-[4-(2-Hydroxyethyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-2-methyl-4-(3-morpholin-4-ylpropyl)-1H-pyrrole-3-carboxylic acid ethyl ester**

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 13.93 (s, 1H, NH), 10.95 (br s, 1H, NH), 7.65 (s, 1H, H-vinyl), 7.06 (t, J = 7.6 Hz, 1H), 6.83 (d, J = 7.6 Hz, 1H), 6.77 (d, J = 7.6 Hz, 1H), 4.85 (br s, 1H, OH), 4.21 (q, J = 7.0 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.73 (m, 2H, CH<sub>2</sub>), 3.53 (m, 4H, 2xCH<sub>2</sub>), 3.09 (m, 2H, CH<sub>2</sub>), 2.92 (m, 2H, CH<sub>2</sub>), 2.53 (s, 3H, CH<sub>3</sub>), 2.31 (m, 6H, 3xCH<sub>2</sub>), 1.68 (m, 2H, CH<sub>2</sub>), 1.29 (t, J = 7.0 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI m/z 467 [M<sup>+</sup>].

**Example 132: 5-[4-(2-Hydroxyethyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-2-methyl-4-(3-pyrrolidin-1-ylpropyl)-1H-pyrrole-3-carboxylic acid ethyl ester**

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 13.93 (s, 1H, NH), 10.94 (br s, 1H, NH), 7.68 (s, 1H, H-vinyl), 7.06 (t, J = 7.3 Hz, 1H), 6.83 (d, J = 7.3 Hz, 1H), 6.77 (d, J = 7.3 Hz, 1H), 4.87 (br s, 1H, OH), 4.21 (q, J = 7.2 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.72 (m, 2H, CH<sub>2</sub>), 3.09 (m, 2H, CH<sub>2</sub>), 2.94 (m, 2H, CH<sub>2</sub>), 2.53 (s, 3H, CH<sub>3</sub>), 2.38 (m, 6H, 3xCH<sub>2</sub>), 1.67 (m, 6H, 3xCH<sub>2</sub>), 1.29 (t, J = 7.2 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI m/z 451 [M<sup>+</sup>].

**Example 133: 4-(3-Dimethylaminopropyl)-2-methyl-5-(2-oxo-5-sulfamoyl-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid ethyl ester**

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 13.73 (s, 1H, NH), 11.25 (br s, 1H, NH), 8.09 (d, J = 1.6 Hz, 1H), 7.90 (s, 1H, H-vinyl), 7.62 (dd, J = 1.6 & 8.1 Hz, 1H), 7.20 (br s, 2H, H<sub>2</sub>NSO<sub>2</sub>), 7.02 (d, J = 8.1 Hz, 1H), 4.21 (q, J = 7.1 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.01 (m, 2H, CH<sub>2</sub>), 2.54 (s, 3H, CH<sub>3</sub>), 2.16 (m, 2H, CH<sub>2</sub>), 2.11 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.66 (m, 2H, CH<sub>2</sub>), 1.29 (t, J = 7.1 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI m/z 460 [M<sup>+</sup>].



WO 00/08202

194

PCT/US99/17845

**Example 134: 3-[3-(3-Dimethylaminopropyl)-4-ethoxycarbonyl-5-methyl-1H-pyrrol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid**

<sup>1</sup>HNMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 13.77 (s, 1H, NH), 11.25 (br s, 1H, NH), 8.24 (s, 1H), 7.85 (s, 1H, H-vinyl), 7.76 (d, 1H), 6.82 (d, 1H), 4.21 (q, *J* = 6.9 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.0 (m, 2H, CH<sub>2</sub>), 2.54 (s, 3H, CH<sub>3</sub>), 2.20 (m, 2H, CH<sub>2</sub>), 2.14 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.64 (m, 2H, CH<sub>2</sub>), 1.29 (t, *J* = 6.9 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS *m/z* 426.2 [M+H].

**Example 135: 4-(3-Dimethylaminopropyl)-5-[4-(2-hydroxyethyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester**

<sup>1</sup>HNMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 13.95 (s, 1H, NH), 10.97 (br s, 1H, NH), 7.69 (s, 1H, H-vinyl), 7.05 (t, *J* = 7.7 Hz, 1H), 6.82 (d, *J* = 7.7 Hz, 1H), 6.75 (d, *J* = 7.7 Hz, 1H), 4.90 (br s, 1H, OH), 4.20 (q, *J* = 7.2 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.72 (m, 2H, CH<sub>2</sub>), 3.09 (m, 2H, CH<sub>2</sub>), 2.90 (m, 2H, CH<sub>2</sub>), 2.52 (s, 3H, CH<sub>3</sub>), 2.20 (m, 2H, CH<sub>2</sub>), 2.07 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.63 (m, 2H, CH<sub>2</sub>), 1.28 (t, *J* = 7.2 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI *m/z* 425 [M<sup>+</sup>].

**Example 136: 5-[4-(2-Hydroxyethyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-2-methyl-4-[3-(4-methylpiperazin-1-yl)-propyl]-1H-pyrrole-3-carboxylic acid ethyl ester**

<sup>1</sup>HNMR (360 MHz, DMSO-*d*<sub>6</sub>) δ 14.0 (s, 1H, NH), 10.94 (br s, 1H, NH), 7.70 (s, 1H, H-vinyl), 7.06 (t, *J* = 7.4 Hz, 1H), 6.83 (d, *J* = 7.4 Hz, 1H), 6.76 (d, *J* = 7.4 Hz, 1H), 4.77 (br s, 1H, OH), 4.22 (q, *J* = 7.21 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.72 (m, 2H, CH<sub>2</sub>), 3.36 (m, 2H, CH<sub>2</sub>), 3.13 (m, 4H, 2xCH<sub>2</sub>), 2.56 (m, 2H, CH<sub>2</sub>), 2.53 (s, 3H, CH<sub>3</sub>), 2.09 (m, 8H, 4xCH<sub>2</sub>), 1.28 (t, *J* = 7.1 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

WO 00/08202

195

PCT/US99/17845

**Example 137: 4-(3-Dimethylaminopropyl)-2-methyl-5-(5-methylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid ethyl ester**

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide  
5 (113 mg, 0.5 mmol) was condensed with 4-(3-dimethylamino-propyl)-5-formyl-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester (133 mg) to give 96 mg (40%) of the title compound.

<sup>1</sup>HNMR (360 MHz, DMSO-*d*<sub>6</sub>) δ 13.73 (s, br, 1H, NH), 11.32 (s, br, 1H, NH), 8.05 (d, *J* = 1.8 Hz, 1H, H-4), 7.92 (s, 1H, H-vinyl), 7.57 (dd, *J* = 1.8 & 8.1 Hz, 1H, H-6), 7.22 (m, 1H, CH<sub>3</sub>NHSO<sub>2</sub>), 7.05 (d, *J* = 8.1 Hz, 1H, H-7), 4.21 (q, *J* = 7.1 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.01 (m, 2H, CH<sub>2</sub>), 2.54 (s, 3H, CH<sub>3</sub>), 2.41 (d, *J* = 4.0 Hz, 3H, CH<sub>3</sub>NHSO<sub>2</sub>), 2.14 (s, 8H, CH<sub>2</sub> & N(CH<sub>3</sub>)<sub>2</sub>), 1.66 (m, 2H, CH<sub>2</sub>), 1.29 (t, *J* = 7.1 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

15 MS-EI *m/z* 474 [M<sup>+</sup>].

**Example 138: 4-(3-Dimethylaminopropyl)-2-methyl-5-(5-methylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid**

A mixture of 4-(3-dimethylaminopropyl)-5-formyl-2-methyl-1H-pyrrole-3-carboxylic acid ethyl ester (300 mg, 1.13 mmol)  
20 and sodium hydroxide in MeOH (4 mL)/water (2 mL) was stirred at 80° C for 19 hr. The reaction was then diluted with EtOAc. The aqueous layer was separated, neutralized and concentrated. The residue was dissolved in MeOH, filtered and the filtrate  
25 was concentrated to give 251 mg (94%) of 4-(3-dimethylaminopropyl)-5-formyl-2-methyl-1H-pyrrole-3-carboxylic acid.

<sup>1</sup>HNMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 11.88 (br s, 1H, NH), 9.50 (s, 1H, CHO), 2.96 (m, 2H, CH<sub>2</sub>), 2.39 (s, 3H, CH<sub>3</sub>), 2.22 (m, 2H, CH<sub>2</sub>), 2.14 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.68 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>).

30 MS-EI *m/z* 238 [M<sup>+</sup>].

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide (113 mg) was condensed with 4-(3-dimethylaminopropyl)-5-

WO 00/08202

196

PCT/US99/17845

formyl-2-methyl-1H-pyrrole-3-carboxylic acid (119 mg) to give 50 mg 4-(3-dimethylaminopropyl)-2-methyl-5-(5-methylsulfamoyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrole-3-carboxylic acid.

- 5 <sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 13.52 (s, br, 1H, NH), 11.25 (s, br, 1H, COOH), 8.01 (d, J = 1.7 Hz, 1H, H-4), 7.84 (s, 1H, H-vinyl), 7.52 (dd, J = 1.7 & 8.3 Hz, 1H, H-6), 7.22 (m, 1H, CH<sub>3</sub>NHSO<sub>2</sub>), 7.05 (d, J = 8.3 Hz, 1H, H-7), 3.08 (m, 2H, CH<sub>2</sub>), 2.56 (s, 3H, CH<sub>3</sub>), 2.40 (m, 3H, CH<sub>3</sub>NHSO<sub>2</sub>), 2.17 (m, 8H, CH<sub>2</sub> & N(CH<sub>3</sub>)<sub>2</sub>), 1.72 (m, 2H, CH<sub>2</sub>).

MS-EI m/z 446 [M<sup>+</sup>].

**Example 139: 4-(2-Carboxyethyl)-3-methyl-5-(4-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester**

- 15 <sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 13.97 (s, br, 1H, NH), 12.15 (br s, 1H, COOH), 11.03 (s, 1H, NH), 7.69 (s, 1H, H-vinyl), 7.10 (t, J = 7.5 Hz, 1H), 6.83 (d, J = 7.5 Hz, 1H), 6.76 (d, J = 7.5 Hz, 1H), 4.28 (q, J = 7.0 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 2.88 (t, 2H, CH<sub>2</sub>), 2.62 (s, 3H, CH<sub>3</sub>), 2.43 (t, 2H, CH<sub>2</sub>), 2.26 (s, 3H, CH<sub>3</sub>),  
20 1.31 (t, J = 7.0 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI m/z 382 [M<sup>+</sup>].

**Example 140: 4-(2-Carboxyethyl)-5-(5-methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester**

- 25 <sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 13.97 (s, br, 1H, NH), 12.14 (br s, 1H, COOH), 10.81 (s, 1H, NH), 7.71 (s, 1H, H-vinyl), 7.46 (m, 1H), 6.78 (m, 2H), 4.28 (q, J = 7.0 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 2.97 (t, 2H, CH<sub>2</sub>), 2.40 (t, 2H, CH<sub>2</sub>), 2.26 (s, 3H, CH<sub>3</sub>), 1.31 (t, J = 7.0 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

- 30 MS-EI m/z 398 [M<sup>+</sup>].

WO 00/08202

197

PCT/US99/17845 ,

**Example 141: 4-(2-Carboxyethyl)-5-(6-methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester**

<sup>1</sup>HNMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 13.70 (s, br, 1H, NH), 12.11 (br s, 1H, COOH), 10.95 (s, 1H, NH), 7.0 (d, *J* = 8.4 Hz, 1H, H-4), 7.55 (s, 1H, H-vinyl), 6.59 (dd, *J* = 2.7 & 8.4 Hz, 1H, H-5), 6.45 (d, *J* = 2.7 Hz, 1H, H-7), 4.27 (q, *J* = 7.0 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 2.92 (t, 2H, CH<sub>2</sub>), 2.39 (t, 2H, CH<sub>2</sub>), 2.25 (s, 3H, CH<sub>3</sub>), 1.31 (t, *J* = 7.0 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI *m/z* 398 [M<sup>+</sup>].

**Example 142: 5-(6-Bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-(2-carboxy-ethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester**

<sup>1</sup>HNMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 13.77 (s, br, 1H, NH), 12.13 (br s, 1H, COOH), 11.14 (s, 1H, NH), 7.77 (br s, 2H), .12 (m, 2H), 4.28 (m, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 2.95 (m, 2H, CH<sub>2</sub>), 2.40 (m, 2H, CH<sub>2</sub>), 2.26 (s, 3H, CH<sub>3</sub>), 1.29 (m, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI *m/z* 446 & 448 [M-1 & M+1].

**Example 143: 5-(5-Bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-(2-carboxy-ethyl)-3-methyl-1H-pyrrole-2-**

**carboxylic acid ethyl ester**

<sup>1</sup>HNMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 13.86 (s, br, 1H, NH), 12.2 (br s, 1H, COOH), 11.12 (s, 1H, NH), 8.11 (d, *J* = 2.0 Hz, 1H, H-4), 7.83 (s, 1H, H-vinyl), 7.33 (dd, *J* = 2.0 & 8.3 Hz, 1H, H-6), 6.83 (d, *J* = 8.3 Hz, 1H, H-7), 4.28 (q, *J* = 7.0 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 2.98 (t, 2H, CH<sub>2</sub>), 2.39 (m, 2H, CH<sub>2</sub>), 2.25 (s, 3H, CH<sub>3</sub>), 1.31 (t, *J* = 7.0 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI *m/z* 446 & 448 [M-1 & M+1].

**Example 144: 4-(2-Carboxyethyl)-3-methyl-5-(2-oxo-6-phenyl-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester**

<sup>1</sup>HNMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 13.86 (s, br, 1H, NH), 12.2 (br s, 1H, COOH), 11.11 (s, 1H, NH), 7.89 (d, *J* = 7.7 Hz, 1H),

WO 00/08202

198

PCT/US99/17845

7.77 (s, 1H, H-vinyl), 7.65 (m, 2H), 7.46 (t, 2H), 7.32-7.39 (m, 2H), 7.12 (d,  $J = 1.6$  Hz, 1H), 4.29 (q,  $J = 7.0$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 2.97 (m, 2H,  $\text{CH}_2$ ), 2.42 (m, 2H,  $\text{CH}_2$ ), 2.27 (s, 3H,  $\text{CH}_3$ ), 1.31 (t,  $J = 7.0$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

5 MS-EI  $m/z$  444  $[\text{M}^*]$ .

**Example 145: 4-(2-Carboxyethyl)-3-methyl-5-(2-oxo-5-sulfamoyl-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester**

$^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.84 (s, br, 1H, NH), 11.0 (br s, 2H), 8.26 (d,  $J = 1.6$  Hz, 1H, H-4), 7.88 (s, 1H, H-vinyl), 7.67 (dd,  $J = 1.6$  & 8.4 Hz, 1H, H-6), 7.22 (br m, 2H,  $\text{H}_2\text{NSO}_2$ ), 7.03 (d,  $J = 8.4$  Hz, 1H, H-7), 4.29 (q,  $J = 7.1$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 2.97 (t, 2H,  $\text{CH}_2$ ), 2.37 (m, 2H,  $\text{CH}_2$ ), 2.26 (s, 3H,  $\text{CH}_3$ ), 1.31 (t,  $J = 7.1$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

15 MS-EI  $m/z$  447  $[\text{M}^*]$ .

**Example 146: 4-(2-Carboxyethyl)-3-methyl-5-(5-methylsulfamoyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester**

$^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.89 (s, br, 1H, NH), 12.2 (br s, 1H, COOH), 11.43 (s, 1H, NH), 8.25 (d,  $J = 1.8$  Hz, 1H, H-4), 7.94 (s, 1H, H-vinyl), 7.62 (dd,  $J = 1.8$  & 8.4 Hz, 1H, H-6), 7.27 (m, 1H,  $\text{HNSO}_2$ ), 7.07 (d,  $J = 8.4$  Hz, 1H, H-7), 4.29 (q,  $J = 7.1$  Hz, 2H,  $\text{OCH}_2\text{CH}_3$ ), 3.01 (t, 2H,  $\text{CH}_2$ ), 2.45 (m, 2H,  $\text{CH}_2$ ), 2.42 (d,  $J = 5.3$  Hz, 3H,  $\text{NCH}_3$ ), 2.27 (s, 3H,  $\text{CH}_3$ ), 1.32 (t,  $J = 7.1$  Hz, 3H,  $\text{OCH}_2\text{CH}_3$ ).

25 MS-EI  $m/z$  461  $[\text{M}^*]$ .

**Example 147: 4-(2-Carboxyethyl)-5-(5-dimethylsulfamoyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester**

30  $^1\text{H}$ NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.88 (s, br, 1H, NH), 12.14 (br s, 1H, COOH), 11.47 (s, 1H, NH), 8.26 (d,  $J = 1.8$  Hz, 1H, H-4), 8.0 (s, 1H, H-vinyl), 7.58 (dd,  $J = 1.8$  & 7.9 Hz, 1H, H-6), 7.10 (d,  $J = 7.9$  Hz, 1H, H-7), 4.29 (q,  $J = 7.2$  Hz, 2H,

WO 00/08202

199

PCT/US99/17845 ,

OCH<sub>2</sub>CH<sub>3</sub>), 3.03 (t, 2H, CH<sub>2</sub>), 2.61 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 2.42 (t, 2H, CH<sub>2</sub>), 2.26 (s, 3H, CH<sub>3</sub>), 1.32 (t, J = 7.2 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>).

MS-EI m/z 489 [M<sup>+</sup>].

5 **Example 148: 4-(2-Carboxyethyl)-5-(5-isopropylsulfamoyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester**

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 13.84 (s, br, 1H, NH), 11.0 (br s, 2H, COOH & NH), 8.23 (d, J = 1.5 Hz, 1H, H-4), 7.92 (s, 1H, H-vinyl), 7.64 (dd, J = 1.5 & 8.4 Hz, 1H, H-6), 7.43 (br m, 1H, HNC(CH<sub>3</sub>)<sub>2</sub>), 7.05 (d, J = 8.4 Hz, 1H, H-7), 4.29 (q, J = 7.1 Hz, 2H, OCH<sub>2</sub>CH<sub>3</sub>), 2.96 (t, 2H, CH<sub>2</sub>), 2.34 (t, 2H, CH<sub>2</sub>), 2.25 (s, 3H, CH<sub>3</sub>), 1.31 (t, J = 7.1 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>), 0.95 (d, J = 6.6 Hz, 6H, HNC(CH<sub>3</sub>)<sub>2</sub>).

MS-EI m/z 489 [M<sup>+</sup>].

15 **Example 149: 3-[3-(3-Dimethylaminopropyl)-1H-indol-2-ylmethylenel]-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide**

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide (98 mg, 0.43 mmol) was condensed with 3-(3-dimethylamino-propyl)-1H-indole-2-carbaldehyde (100 mg) to give the title compound as an orange solid.

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.90 (s, 1H, NH), 11.40 (br s, 1H, NH), 8.17 (d, J = 1.5 Hz, 1H), 8.15 (s, 1H, H-vinyl), 7.68 (d, J = 8.4 Hz, 1H), 7.63 (dd, J = 1.5 & 8.4 Hz, 1H), 7.54 (d, J = 8.7 Hz, 1H), 7.30 (m, 2H), 7.08 (m, 2H), 3.15 (t, J = 6.6 Hz, 2H, CH<sub>2</sub>), 2.43 (d, J = 4.2 Hz, 3H, NCH<sub>3</sub>), 2.17 (t, J = 6.6 Hz, 2H, CH<sub>2</sub>), 2.14 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.78 (m, 2H, CH<sub>2</sub>).

MS-EI m/z 438 [M<sup>+</sup>].

30 **Example 150: 3-[3-(3-Dimethylaminopropyl)-1H-indol-2-ylmethylenel]-4-(2-hydroxyethyl)-1,3-dihydroindol-2-one**

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 13.12 (s, 1H, NH), 11.01 (br s, 1H, NH), 7.95 (s, 1H, H-vinyl), 7.66 (d, J = 7.9 Hz, 1H), 7.52 (d, J = 7.9 Hz, 1H), 7.26 (t, J = 7.54 Hz, 1H), 7.11 (t, J =

WO 00/08202

200

PCT/US99/17845 ,

7.9 Hz, 1H), 7.06 (t,  $J = 7.4$  Hz, 1H), 6.87 (d,  $J = 7.4$  Hz, 1H), 6.78 (d,  $J = 7.4$  Hz, 1H), 4.91 (br s, 1H, OH), 3.79 (m, 2H, CH<sub>2</sub>), 3.16 (m, 2H, CH<sub>2</sub>), 3.04 (m, 2H, CH<sub>2</sub>), 2.21 (m, 2H, CH<sub>2</sub>), 2.09 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.77 (m, 2H, CH<sub>2</sub>).

5 MS  $m/z$  389 [M<sup>+</sup>].

**Example 151: 3-[3-(3-Dimethylaminopropyl)-1H-indol-2-ylmethylene]-5-methoxy-1,3-dihydroindol-2-one**

<sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  13.11 (s, 1H, NH), 10.84 (br s, 1H, NH), 7.92 (s, 1H, H-vinyl), 7.65 (d,  $J = 8.1$  Hz, 1H), 7.50  
10 (d,  $J = 8.1$  Hz, 1H), 7.44 (s, 1H), 7.26 (t,  $J = 7.5$  Hz, 1H), 7.05 (t,  $J = 7.5$  Hz, 1H), 6.79 (s, 2H), 3.79 (s, 3H, OCH<sub>3</sub>), 3.11 (m, 2H, CH<sub>2</sub>), 2.20 (m, 2H, CH<sub>2</sub>), 2.12 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.76 (m, 2H, CH<sub>2</sub>).

MS  $m/z$  375 [M<sup>+</sup>].

15 **Example 152: 5-Methyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one**

3-Methylindole-2-carbaldehyde was prepared as described in the literature: 1) David W.M. Benzies, Pilar Martinez Fresneda and R. Alan Jones, *Synthetic Communications*, 16 (14),  
20 1799-1807 (1986) and 2) Chatterjee, A. and Biswas, K.M., *J. Org. Chem.*, 1973, 38, 4002.

A mixture of 5-methyl-2-oxindole (59 mg), 3-methylindole-2-carbaldehyde (56 mg) and piperidine (30 mg) in ethanol (1 mL) was held in a sealed tube at 90°C overnight.  
25 The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 72 mg (74%) of 5-methyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one.

<sup>1</sup>H NMR (360 MHz, DMSO-*d*<sub>6</sub>) 13.03 (s, br, 1H, NH), 10.88 (s, br, 1H, NH), 7.84 (s, 1H, vinyl), 7.71 (s, br, 1H), 6.62 (d,  $J = 8$  Hz, 1H), 7.48 (d,  $J = 8$  Hz, 1H), 7.26 (t,  $J = 7$  Hz, 1H), 7.06 (t,  $J = 7$  Hz, 1H), 7.0 (d,  $J = 8$  Hz, 1H), 6.78 (d,  $J = 8$  Hz, 1H), 2.59 (s, 3H, CH<sub>3</sub>), 2.32 (s, 3H, CH<sub>3</sub>).

WO 00/08202

201

PCT/US99/17845

MS m/z 289 [M+1]<sup>+</sup>.

**Example 153: 3-(3-Methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide**

A mixture of 5-aminosulfonyl-2-oxindole (84 mg),  
5 3-methylindole-2-carbaldehyde (56 mg) (prepared according to  
*Synthetic Communications*, 1986, 16, 1799) and piperidine (30  
mg) in ethanol (1 mL) was held in a sealed tube at 90°C  
overnight. The mixture was cooled to room temperature. The  
solid was collected by vacuum filtration, washed with cold  
10 ethanol and dried in a vacuum oven to give 88 mg (71% yield)  
of 3-(3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-  
dihydro-1H-indole-5-sulfonic acid amide.

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) 12.86 (s, br, 1H, NH), 11.3 (s,  
br, 1H, NH), 8.3 (d, J = 2Hz, 1H), 7.97 (s, 1H, vinyl), 7.63  
15 (dd, J = 2 and 8Hz, 1H), 7.6 (d, J = 8Hz, 1H), 7.47 (d, J =  
8Hz, 1H), 7.24 (t, J = 7.5Hz, 1H), 7.12 (s, br, 2H, NH<sub>2</sub>), 7.02  
(t, J = 7.5Hz, 1H), 6.98 (d, J = 8Hz, 1H), 2.57 (s, 3H, CH<sub>3</sub>).

MS m/z 354.1 [M+1]<sup>+</sup>

**Example 154: 3-(3-Methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide**

A suspension of 5-chlorosulfonyl-2-oxindole (3.38 g) in  
2M methylamine in tetrahydrofuran (10 mL) was stirred at room  
temperature for 4 hours at which time a white solid had  
formed. The solid was collected by vacuum filtration, washed  
25 twice with 5 mL of water and dried under vacuum at 40°C to  
give 3.0 g (88% yield) of 5-methylaminosulfonyl-2-oxindole.

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 10.87 (s, br, 1H, NH-1), 7.86  
(s, br, 1H, 5-SO<sub>2</sub>NHCH<sub>3</sub>), 7.61 (d, J = 8Hz 1H, H-6), 7.32 (d, J =  
5Hz, 1H, H-4), 6.97 (d, J = 8Hz, 1H, H-7), 2.53 (s, 2H, CH<sub>2</sub>-3),  
30 and 2.36 (s, 3H, 5-SO<sub>2</sub>NHCH<sub>3</sub>).

MS m/z 226.

A mixture of 5-methylaminosulfonyl-2-oxindole (90 mg),  
3-methylindole-2-carbaldehyde (56 mg) (prepared according to



WO 00/08202

202

PCT/US99/17845 ,

*Synthetic Communications*, 1986, 16, 1799) and piperidine (30 mg) in ethanol (1 mL) was held in a sealed tube at 90°C overnight. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 90 mg (70% yield) of 3-(3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide (mixture of isomers).

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) 11.13 (s, br, 1H, NH), 10.94 (s, br, 1H, NH), 8.84 (d, J = 16Hz, 1H), 7.98 (s, 1H), 6.97-7.66 (m, 6H, Ar-H), 2.6 (s, 3H, CH<sub>3</sub>), 2.4 (d, J = 5Hz, 3H, CH<sub>3</sub>N).

**Example 155: 3-(3-Methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide**

A mixture of 5-dimethylaminosulfonyl-2-oxindole (96 mg), 3-methylindole-2-carbaldehyde (56 mg) (prepared according to *Synthetic Communications*, 1986, 16, 1799) and piperidine (30 mg) in ethanol (1 mL) was held in a sealed tube at 90°C overnight. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 76 mg (57% yield) of 3-(3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide (mixture of isomers).

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) 11.09 (s, br, 1H, NH), 10.95 (s, br, 1H, NH), 8.78 (d, J = 16Hz, 1H), 7.8 (s, 1H, vinyl), 7.45-7.54 (m, 2H), 7.32 (d, J = 8Hz, 1H), 7.1 (t, 1H), 6.98 (d, J = 8Hz, 1H), 6.93 (t, 1H), 2.56 (s, 6H, 2 x CH<sub>3</sub>), 2.39 (s, 3H, CH<sub>3</sub>)

**Example 156: 3-(3-Methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid (piperidine salt)**

A mixture of 5-carboxy-2-oxindole (113 mg), 3-methylindole-2-carbaldehyde (56 mg) (prepared according to *Synthetic Communications*, 1986, 16, 1799) and piperidine (30 mg) in ethanol (1 mL) was held in a sealed tube at 90°C overnight. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold

WO 00/08202

203

PCT/US99/17845 ,

ethanol and dried in a vacuum oven to give 75 mg (58% yield) of 3-(3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid (piperidine salt).

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) 13.0 (s, br, 1H, NH), 8.35 (d, J = 1.5Hz, 1H, H-4), 7.91 (s, 1H, vinyl), 7.82 (dd, J = 1.5 and 8Hz, 1H, H-6), 7.63 (d, J = 8Hz, 1H), 7.49 (d, J = 8Hz, 1H), 7.26 (t, J = 7.5Hz, 1H), 7.06 (t, J = 7.5Hz, 1H), 6.87 (d, J = 8Hz, 1H), 2.91 (t, 4H, piperidine), 2.6 (s, 3H, CH<sub>3</sub>), 1.6 (m, 4H, piperidine), 1.54 (m, 2H, piperidine).

10 **Example 157: 5-Acetyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one**

2-Oxindole (3 g) was suspended in 1,2-dichloroethane and 3.2 mL of acetyl chloride slowly added. The resulting suspension was stirred at 50°C for 5 hours, cooled, and poured  
15 into water. The resulting precipitate was collected by vacuum filtration, washed copiously with water and dried under vacuum to give 2.9 g (73% yield) of 5-acetyl-2-oxindole as a brown solid.

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) δ 10.75 (s, br, NH), 7.83 (d, J = 8Hz, 1H), 7.78 (s, 1H, H-4), 6.88 (d, J = 8Hz, 1H), 3.53 (s, 2H, CH<sub>2</sub>), 2.49 (s, 3H, CH<sub>3</sub>).

A mixture of 5-acetyl-2-oxindole (70 mg), 3-methylindole-2-carbaldehyde (56 mg) (prepared according to *Synthetic Communications*, 1986, 16, 1799) and piperidine (30  
25 mg) in ethanol (1 mL) was held in a sealed tube at 90°C overnight. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum to give 83 mg (75% yield) of 5-acetyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-  
30 dihydro-indol-2-one.

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) 12.86 (s, br, 1H, NH), 11.29 (s, br, 1H, NH), 8.43 (d, J = 1.5Hz, 1H, H-4), 8.01 (s, 1H, vinyl), 7.78 (dd, J = 1.5 and 8Hz, 1H, H-6), 7.58 (d, J = 8Hz,

WO 00/08202

204

PCT/US99/17845

1H), 7.45 (d, J = 8Hz, 1H), 7.23 (t, 1H), 7.01 (t, 1H), 6.92 (d, J = 8Hz, 1H), 2.58 (s, 3H, CH<sub>3</sub>), 2.54 (s, 3H, CH<sub>3</sub>).

MS m/z 317.2 [M+1]<sup>+</sup>.

**Example 158: 5-Acetyl-3-(1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one**

Indole-2-carbaldehyde was prepared as described in the literature: Michel Barbier, Michel Devys, Christiane Tempête, Albert Kollmann and Jean-François Bousquet, *Synthetic Communications*, 23(22), 3109-3117 (1993).

A mixture of 5-acetyl-2-oxindole (88 mg), indole-2-carbaldehyde (87 mg) and piperidine (4 mg) in ethanol (2 mL) was held in a sealed tube at 90°C for 3 hours. The mixture was cooled to room temperature. The solid which formed was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 133 mg (88% yield) of 5-acetyl-3-(1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one as an orange solid.

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) 12.81 (s, br, 1H, NH), 11.38 (s, br, 1H, NH), 8.38 (d, J = 1Hz, H-4), 8.17 (s, 1H, vinyl), 7.88 (dd, J = 1 and 8Hz, 1H, H-6), 7.69 (d, J = 8Hz, 1H), 7.58 (d, J = 8Hz, 1H), 7.29 (t, 1H), 7.21 (s, 1H), 7.09 (t, 1H), 7.0 (d, J = 8Hz, 1H), 2.59 (s, 3H, CH<sub>3</sub>).

MS m/z 303.1 [M+1]<sup>+</sup>.

**Example 159: 3-(1H-Indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide**

A mixture of 5-aminosulfonyl-2-oxindole (106 mg), indole-2-carbaldehyde (87 mg) (prepared according to *Synthetic Communications*, 1993, 23, 3109) and piperidine (4 mg) in ethanol (2 mL) was held in a sealed tube at 90°C for 3 hours.

The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 140 mg (83% yield) of 3-(1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide as an orange solid.

WO 00/08202

205

PCT/US99/17845 ,

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) 12.81 (s, br, 1H, NH), 11.38 (s, br, 1H, NH), 8.18 (d, J = 1Hz, H-4), 8.12 (s, 1H, vinyl), 7.7 (m, 2H), 7.6 (d, J = 8Hz, 1H), 7.26-7.29 (m, 2H), 7.2 (s, 2H, NH<sub>2</sub>), 7.05-7.11 (m, 2H).

5 MS m/z 340 [M+1]<sup>+</sup>.

**Example 160: 5-Amino-3-(1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one**

A mixture of 5-tert-butoxycarbonylamino-2-oxindole (124 mg), indole-2-carbaldehyde (87 mg) (prepared according to  
10 *Synthetic Communications*, 1993, 23, 3109) and piperidine (4 mg) in ethanol (2 mL was held in a sealed tube at 90°C for 3 hours. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 196 mg of  
15 5-tert-butoxycarbonylamino-3-(1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one as a bright orange solid. The product was then dissolved in trifluoroacetic acid/dichloromethane (2 mL each) and stirred at room temperature for 1 hour. The reaction mixture was then  
20 concentrated. The residue was dissolved in water and basified with saturated sodium bicarbonate solution. The solid was collected by vacuum filtration, washed with water and dried in a vacuum oven to give 129 mg of 5-amino-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one as a brown solid.

25 <sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) 13.1 (s, br, 1H, NH), 10.57 (s, br, 1H, NH), 7.65 (s, 1H, vinyl), 7.64 (d, J = 8Hz, 1H), 7.54 (d, J = 8Hz, 1H), 7.24 (t, 1H), 7.1 (s, 1H), 7.06 (t, 1H), 6.93 (d, J = 2Hz, 1H), 6.61 (d, J = 8Hz, 1H), 6.5 (dd, J = 2 and 8Hz, 1H), 4.8 (s, br, 2H, NH<sub>2</sub>).

30 MS m/z 276.1 [M+1]<sup>+</sup>.

**Example 161: 3-(1H-Indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid**

A mixture of 5-carboxy-2-oxindole (88.5 mg), indole-2-carbaldehyde (87 mg) (prepared according to *Synthetic*

WO 00/08202

206

PCT/US99/17845 ,

Communications, 1993, 23, 3109) and piperidine (1 drop) in ethanol (3 mL) was held in a sealed tube at 95°C for 4 hours. The mixture was cooled to room temperature. The solid was filtered and acidified with 2N hydrochloric acid. The  
5 resulting solid was collected by vacuum filtration, washed with cold ethanol and dried in vacuum oven to give 60 mg (40% yield) of 3-(1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid as a mustard-colored solid.

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) 12.8 (s, br, 1H, NH), 12.69 (s,  
10 1H, COOH), 11.35 (s, br, 1H, NH), 8.33 (s, 1H), 8.16 (s, 1H), 7.86 (dd, J = 1 and 8Hz, 1H), 7.68 (d, J = 8Hz, 1H), 7.59 (d, J = 8Hz, 1H), 7.28 (t, 1H), 7.21 (s, 1H), 7.08 (t, 1H), 6.99 (d, J = 8Hz, 1H).

MS m/z 304 [M]<sup>+</sup>.

15 **Example 162: 6-Chloro-3-(1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one**

A mixture of 6-chloro-2-oxindole (41 mg) (commercially available), indole-2-carbaldehyde (42 mg) (prepared according to *Synthetic Communications*, 1993, 23, 3109) and piperidine  
20 (1 drop) in ethanol (3 mL) was held in a sealed tube at 95°C for 3 hours. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in vacuum oven to give 62 mg (87% yield) of 6-chloro-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one as  
25 a shiny orange solid.

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) 12.83 (s, br, 1H, NH), 11.17 (s, br, 1H, NH), 7.98 (s, 1H), 7.74 (d, J = 8Hz, 1H), 7.67 (d, J = 8Hz, 1H), 7.58 (d, J = 8Hz, 1H), 7.28 (m, 1H), 7.15 (s, br, 1H), 7.08 (m, 2H), 6.91 (d, J = 2Hz, 1H).

30 MS m/z 293 and 295.

**Example 163: 3-(1H-Indol-2-ylmethylene)-1,3-dihydro-indol-2-one**

A mixture of 2-oxindole (133 mg), indole-2-carbaldehyde (174 mg) (prepared according to *Synthetic Communications*,

WO 00/08202

207

PCT/US99/17845

1993, 23, 3109) and piperidine (10 mg) in ethanol was held in a sealed tube at 95°C for 4 hours. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 3-(1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one.

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) δ 12.97 (s, br, 1H, NH), 11.01 (s, br, 1H, NH), 7.92 (s, 1H, H-vinyl), 7.72 (d, J = 7Hz, 1H), 7.66 (d, J = 7Hz, 1H), 7.57 (dd, J = 1, 8Hz, 1H), 7.27 (dt, J = 1, 7Hz, 1H), 7.21 (dt, J = 1, 8Hz, 1H), 7.13 (s, 1H), 7.01-7.1 (m, 2H), 6.91 (d, J = 8Hz, 1H).

MS EI 260.

**Example 164: 5-Chloro-3-(1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one**

A mixture of 5-chloro-2-oxindole (167 mg), indole-2-carbaldehyde (174 mg) (prepared according to *Synthetic Communications*, 1993, 23, 3109) and piperidine (10 mg) in ethanol was held in a sealed tube at 95°C for 4 hours. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 5-chloro-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one.

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) δ 12.88 (s, br, 1H, NH), 11.12 (s, br, 1H, NH), 8.07 (s, 1H, H-vinyl), 7.86 (d, J = 2Hz, 1H), 7.68 (d, J = 8Hz, 1H), 7.59 (dd, J = 1, 8Hz, 1H), 7.28 (m, 1H), 7.24 (dd, J = 2, 8Hz, 1H), 7.14 (s, 1H), 7.09 (dt, J = 1, 8Hz, 1H), 6.90 (d, J = 8Hz, 1H).

MS EI 294 and 296.

**Example 165: 5-Bromo-3-(1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one**

A mixture of 5-bromo-2-oxindole (212 mg), indole-2-carbaldehyde (174 mg) (prepared according to *Synthetic Communications*, 1993, 23, 3109) and piperidine

WO 00/08202

208

PCT/US99/17845 ,

(10 mg) in ethanol was held in a sealed tube at 95°C for 4 hours. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 5-bromo-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one.

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) δ 12.87 (s, br, 1H, NH), 11.12 (s, br, 1H, NH), 8.07 (s, 1H, H-vinyl), 7.98 (d, J = 2Hz, 1H), 7.68 (d, J = 8Hz, 1H), 7.59 (d, J = 8Hz, 1H), 7.36 (dd, J = 2, 8Hz, 1H), 7.28 (dt, J = 1, 8Hz, 1H), 7.14 (s, 1H), 7.08 (dt, J = 1, 8Hz, 1H), 6.86 (d, J = 8Hz, 1H).

MS EI 338 and 340.

**Example 166: 3-(1H-indol-2-ylmethylene)-4-methyl-1,3-dihydro-indol-2-one**

A mixture of 4-methyl-2-oxindole (147 mg), indole-2-carbaldehyde (174 mg) (prepared according to *Synthetic Communications*, 1993, 23, 3109) and piperidine (10 mg) in ethanol was held in a sealed tube at 95°C for 4 hours. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 3-(1H-indol-2-ylmethylene)-4-methyl-1,3-dihydro-indol-2-one-

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) δ 13.12 (s, br, 1H, NH), 11.02 (s, br, 1H, NH), 7.79 (s, 1H, H-vinyl), 7.64 (d, J = 8Hz, 1H), 7.57 (d, J = 8Hz, 1H), 7.27 (d, J = 7Hz, 1H), 7.23 (s, 1H), 7.05 - 7.14 (m, 2H), 6.84 (d, J = 8Hz, 1H), 6.78 (d, J = 8Hz, 1H), 2.61 (s, 3H, CH<sub>3</sub>).

MS EI 274.

**Example 167: 3-(3-Methyl-1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one**

A mixture of 2-oxindole (133 mg), 3-methylindole-2-carbaldehyde (190 mg) (prepared according to *Synthetic Communications*, 1986, 16, 1799) and piperidine (10 mg) in ethanol was held in a sealed tube at 95°C for 4 hours. The

WO 00/08202

209

PCT/US99/17845

mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindole-2-one.

5 <sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) δ 13.01 (s, br, 1H, NH), 10.99 (s, br, 1H, NH), 7.89 (s, 1H, H-vinyl), 7.88 (d, J = 7Hz, 1H), 7.63 (d, J = 8Hz, 1H), 7.49 (d, J = 8Hz, 1H), 7.27 (dt, J = 1, 8Hz, 1H), 7.2 (dt, J = 1, 8Hz, 1H), 7.01-7.08 (m, 2H), 6.9 (d, J = 8Hz, 1H), 2.59 (s, 3H, CH<sub>3</sub>).

10 MS EI 274.

**Example 168: 5-Chloro-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one**

A mixture of 5-chloro-2-oxindole (167 mg), 3-methylindole-2-carbaldehyde (190 mg) (prepared according to  
15 *Synthetic Communications*, 1986, 16, 1799) and piperidine (10 mg) in ethanol was held in a sealed tube at 95°C for 4 hours. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 5-chloro-3-(3-methyl-1H-indol-  
20 2-ylmethylene)-1,3-dihydroindol-2-one.

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) δ 12.96 (s, br, 1H, NH), 11.09 (s, br, 1H, NH), 8.09 (d, J = 2Hz, 1H), 8.02 (s, 1H, H-vinyl), 7.63 (d, J = 8Hz, 1H), 7.5 (d, J = 8Hz, 1H), 7.29 (dt, J = 1, 7Hz, 1H), 7.21 (dd, J = 2, 8Hz, 1H), 7.07 (dt, J = 1, 7Hz,  
25 1H), 6.89 (d, J = 8Hz, 1H), 2.61 (s, 3H, CH<sub>3</sub>).

MS EI 308.

**Example 169: 5-Bromo-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one**

A mixture of 5-bromo-2-oxindole (212 mg), 3-methyl-  
30 indole-2-carbaldehyde (190 mg) (prepared according to *Synthetic Communications*, 1986, 16, 1799) and piperidine (10 mg) in ethanol was held in a sealed tube at 95°C for 4 hours. The mixture was cooled to room temperature. The solid



WO 00/08202

210

PCT/US99/17845

was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 5-bromo-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one.

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) δ 12.96 (s, br, 1H, NH), 11.09 (s, br, 1H, NH), 8.21 (d, J = 2Hz, 1H), 8.02 (s, 1H, H-vinyl), 7.63 (d, J = 8Hz, 1H), 7.5 (d, J = 8Hz, 1H), 7.33 (dd, J = 2, 8Hz, 1H), 7.29 (dt, J = 1, 7Hz, 1H), 7.07 (dt, J = 1, 7Hz, 1H), 6.84 (d, J = 8Hz, 1H), 2.62 (s, 3H, CH<sub>3</sub>).

MS EI 352 and 354.

10 **Example 170: 4-Methyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one**

A mixture of 4-methyl-2-oxindole (147 mg), 3-methyl-indole-2-carbaldehyde (190 mg) (prepared according to *Synthetic Communications*, 1986, 16, 1799) and piperidine (10 mg) in ethanol was held in a sealed tube at 95°C for 4 hours. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and dried in a vacuum oven to give 4-methyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydro-indol-2-one.

20 <sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) δ 13.07 (s, br, 1H, NH), 11.0 (s, br, 1H, NH), 7.78 (s, 1H, H-vinyl), 7.64 (d, J = 8Hz, 1H), 7.51 (d, J = 8Hz, 1H), 7.27 (dt, J = 1, 8Hz, 1H), 7.04 - 7.12 (m, 2H), 6.85 (d, J = 8Hz, 1H), 6.78 (d, J = 8Hz, 1H), 2.64 (s, 3H, CH<sub>3</sub>), 2.52 (s, 3H, CH<sub>3</sub>).

25 MS EI 288.

**Example 171: 3-(1H-indol-2-ylmethylene)-5-[(1H-indol-2-ylmethylene)-aminol]-1,3-dihydroindol-2-one**

A mixture of 5-amino-2-oxindole (74 mg), indole-2-carbaldehyde (87 mg) (prepared according to *Synthetic Communications*, 1983, 23, 3109) and piperidine (4 mg) in ethanol (2 mL) was held in a sealed tube at 95 °C for 3 hours. The mixture was cooled to room temperature. The solid was collected by vacuum filtration, washed with cold ethanol and

WO 00/08202

211

PCT/US99/17845

dried in a vacuum oven. The crude solid was then chromatographed on silica gel to give 50 mg (25% yield) of 3-(1H-indol-2-ylmethylene)-5-[(1H-indol-2-ylmethylene)-amino]-1,3-dihydro-indol-2-one

5 **Example 172: 3-(1H-Indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide**

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.03 (br, s, 1H, NH), 10.80 (br, s, 1H, NH), 9.37 (s, 1H), 8.09-8.24 (m, 3H), 7.51 (m, 1H), 7.41 (m, 1H), 7.13 (m, 2H), 7.03 (s, 2H, NH<sub>2</sub>), 6.85 (d, J = 7.8 Hz, 1H).

MS-EI m/z 339 [M]<sup>+</sup>.

**Example 173: 3-(1H-Indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide**

15 <sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.75 (br, s, 1H, NH), 11.40 (br, s, 1H, NH), 8.15 (s, 1H, H-vinyl), 8.10 (d, J = 1.5 Hz, 1H), 7.54-7.65 (m, 3H), 7.20-7.27 (m, 3H), 7.02-7.06 (m, 2H), 2.38 (d, J = 4.8 Hz, 3H, CH<sub>3</sub>).

MS-EI m/z 353 [M]<sup>+</sup>.

20 **Example 174: 3-(1H-Indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid**

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.13 (br, s, 1H, NH), 10.88 (br, s, 1H, NH), 8.22-8.50 (m, 2H), 7.66-7.94 (m, 2H), 7.52 (m, 1H), 7.23 (m, 2H), 6.94 (m, 1H).

MS-EI m/z 304 [M]<sup>+</sup>.

25 **Example 175: 3-(3-Methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide**

30 <sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 12.92 (br, s, 1H, NH), 11.35 (br, s, 1H, NH), 8.36 (d, J = 2.2 Hz, 1H), 8.03 (s, 1H, H-vinyl), 7.67 (m, 2H), 7.53 (d, J = 7.6 Hz, 1H), 7.30 (m, 1H), 7.16 (s, 2H, NH<sub>2</sub>), 7.08 (m, 1H), 7.03 (d, J = 8.6 Hz, 1H), 2.63 (s, 3H, CH<sub>3</sub>).

MS-EI m/z 353 [M]<sup>+</sup>.

**Example 176: 3-(1H-Indol-5-ylmethylene)-2-oxo-2,3-dihydro-1H-**

WO 00/08202

212

PCT/US99/17845

indole-5-sulfonic acid methylamide

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 11.38 (br, s, 1H, NH), 10.85 (br, s, 1H, NH), 8.11 (m, 1H), 8.01 (m, 1H), 7.90 (m, 1H), 7.78 (m, 1H), 7.37-7.53 (m, 3H), 7.10 (m, 1H), 6.95 (m, 1H),  
5 6.4 (m, 1H), 2.28 (s, 3H, CH<sub>3</sub>).

MS-EI m/z 353 [M]<sup>+</sup>.Example 177: 3-(1H-Indol-5-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.54 (br, s, 1H, NH), 10.86 (br, s, 1H, NH), 8.56, 7.76, 7.46, 6.91 (m, 9H).

Example 178: 3-(1H-Indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.03 (br, s, 1H, NH), 10.84 (br, s, 1H, NH), 9.39 (s, 1H), 8.25 (s, 1H, H-vinyl), 8.22 (d, *J* = 1.5 Hz, 1H), 8.16 (m, 1H), 7.40-7.46 (m, 2H), 7.06-7.15 (m, 2H), 7.04 (m, 1H, CH<sub>3</sub>NH), 6.88 (d, *J* = 7.8 Hz, 1H), 2.32 (d, *J* = 5.1 Hz, 3H, CH<sub>3</sub>).

MS-EI m/z 355 [M]<sup>+</sup>.

20 Example 179: 5-Amino-3-(1H-indol-5-ylmethylene)-1,3-dihydro-indol-2-one

MS-EI m/z 275 [M]<sup>+</sup>.Example 180: 5-Amino-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one

25 <sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 13.13 (br, s, 1H, NH), 10.57 (br, s, 1H, NH), 7.63 (s, 1H, H-vinyl), 7.61 (d, *J* = 7.6 Hz, 1H), 7.47 (d, *J* = 7.6 Hz, 1H), 7.25 (t, *J* = 7.6 Hz, 1H), 7.03-7.06 (m, 2H), 6.61 (d, *J* = 8.3 Hz, 1H), 6.49 (dd, *J* = 1.4 & 8.3 Hz, 1H), 4.70 (br s, 2H, NH<sub>2</sub>), 2.55 (s, 3H, CH<sub>3</sub>).

MS-EI m/z 289 [M]<sup>+</sup>.

30 Example 182: 3-(2-Methyl-1H-indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.0 (br, s, 1H, NH), 10.85 (br, s, 1H, NH), 7.90 (s, 1H, H-vinyl), 7.79 (dd, 1H), 7.41-

WO 00/08202

213

PCT/US99/17845 ,

7.44 (m, 2H), 7.02-7.18 (m, 3H), 6.91 (d, 1H), 2.45 (s, 3H, CH<sub>3</sub>).

MS-EI m/z 318 [M]<sup>+</sup>.

Example 183: 3-(1H-Indol-5-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide

MS-EI m/z 367 [M]<sup>+</sup>.

Example 184: 3-(2-Methyl-1H-indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide

<sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 11.95 (br, s, 1H, NH), 10.86 (br, s, 1H, NH), 7.84 (s, 1H, H-vinyl), 7.41 (dd, J = 1.8 & 8.4 Hz, 1H), 7.33 (d, J = 8.4 Hz, 1H), 7.04 (m, 1H), 6.85-6.95 (m, 4H).

MS-EI m/z 381 [M]<sup>+</sup>.

Example 185: 3-(1H-Indol-5-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide

MS-EI m/z 339 [M]<sup>+</sup>.

Example 186: 3-(1H-Indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 12.80 (br, s, 1H, NH), 11.44 (br, s, 1H, NH), 8.30 (s, 1H, H-vinyl), 8.15 (d, J = 2.2 Hz, 1H), 7.70 (d, J = 7.2 Hz, 1H), 7.60 (m, 2H), 7.30 (m, 1H), 7.23 (s, 1H), 7.07-7.12 (m, 2H), 2.63 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 367 [M]<sup>+</sup>.

Example 187: 5-Amino-3-(2-methyl-1H-indol-3-ylmethylene)-1,3-dihydroindol-2-one

MS-EI m/z 289 [M]<sup>+</sup>.

Example 188: 3-(1H-Indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 12.11 (br, s, 1H, NH), 10.94 (br, s, 1H, NH), 9.50 (s, 1H), 8.44 (s, 1H), 8.33 (m, 2H), 7.51-7.53 (m, 2H), 7.25 (m, 2H), 7.03 (d, J = 7.9 Hz, 1H), 2.63 (s, 6H, 2xCH<sub>3</sub>).

MS-EI m/z 367 [M]<sup>+</sup>.

WO 00/08202

214

PCT/US99/17845

**Example 189:** 3-(2-Methyl-1H-indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide

MS-EI  $m/z$  353 [M]<sup>+</sup>.

**Example 190:** 3-[2-(2-Oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid

1-(Morpholin-4-yl)cyclohexene (300 g), 214 g of triethylamine and 1400 mL of dichloromethane were refluxed for 15 minutes and then cooled in a water bath to 15-20 °C. Ethyl succinyl chloride (266 g) dissolved in 500 mL of dichloromethane was added over 30 minutes. The mixture was refluxed for 30 minutes and cooled to ambient temperature in a water bath. The solid was collected by vacuum filtration, washed with 100 mL of dichloromethane and discarded. The filtrate was returned to the flask and the solvent removed by distillation to give 454 g of crude 4-(2-morpholin-4-yl-cyclohex-1-enyl)-4-oxo-butyric acid ethyl ester as an oil.

Crude 4-(2-morpholin-4-ylcyclohex-1-enyl)-4-oxo-butyric acid ethyl ester (454 g), 398 g of diethyl aminomalonate hydrochloride, 162 g of sodium acetate and 350 mL of glacial acetic acid were heated to 108°C over 30 minutes. The mixture was held at 100-108 °C for 2 hours and then cooled to about 50°C in a water bath. Water (2500 mL) and 700 mL of ethyl acetate were added. The ethyl acetate layer was separated and washed three times with brine, twice with saturated sodium bicarbonate solution and once with brine. The solution was dried over anhydrous sodium sulfate, and the ethyl acetate was removed using a rotary evaporator to give 494 g (105% yield) of crude 3-(2-ethoxycarbonyl-ethyl)-4,5,6,7-tetrahydro-1H-indole-2-carboxylic acid ethyl ester as an oil. The crude mixture was chromatographed on a silica gel column using a 1:10 mixture of ethyl acetate:hexane as the eluent to give 122.1 g of pure 3-(2-ethoxycarbonyl-ethyl)-4,5,6,7-tetrahydro-1H-indole-2-carboxylic acid ethyl ester as a low melting

WO 00/08202

215

PCT/US99/17845 ,

solid.

$^1\text{H}$  NMR ( $d_6$ -DMSO):  $\delta$  11.0 (s, 1H, pyrrole NH), 4.2, 4.0 (t, each 4H,  $\text{COCH}_2$ ), 2.8, 2.4 (t, each 4H,  $-\text{CH}_2\text{CH}_2\text{CO}-$ ), 2.4 (m, 4H,  $-\text{CH}_2-$ ,  $-\text{CH}_2-$ ), 1.7 (m, 4H,  $-\text{CH}_2\text{CH}_2-$ ).

5 Purified 3-(2-ethoxycarbonyl-ethyl)-4,5,6,7-tetrahydro-1H-indole-2-carboxylic acid ethyl ester (122.1 g) and 328 mL of 5N sodium hydroxide were refluxed for 80 minutes. The heat was turned off and 165 mL of 10 N hydrochloric acid was carefully added to the vigorously stirred mixture using an addition funnel over the reflux condenser. Hydrochloric acid was added until the pH was 2-3. The mixture was then cooled in an ice bath at which time the oil in the mixture solidified. The solid was collected by vacuum filtration, washed 3 times with water and dried under vacuum at 50-60°C to give 54.9 g (71% yield) of 3-(4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid as a dark brown solid.

$^1\text{H}$  NMR ( $d_6$ -DMSO):  $\delta$  13.1 (s, 1H, pyrrole NH), 11.8 (br s, 1H, COOH), 9.8 (s, 1H, CH), 2.5, 2.3 (t, 4H,  $-\text{CH}_2\text{CH}_2\text{CO}-$ ), 2.4 (m, 4H,  $-\text{CH}_2-$ ,  $-\text{CH}_2-$ ), 1.7 (m, 4H,  $-\text{CH}_2\text{CH}_2-$ ).

20 A mixture of 24 g of dimethylformamide and 300 mL of dichloromethane was cooled to -9°C. Phosphorus oxychloride (50 g) was rapidly added via an addition funnel. 3-(4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid (54.9 g) was added in portions with vigorous stirring. The mixture was warmed to room temperature and then refluxed for 10 minutes. It was then cooled to 5 °C and diluted with 300 mL of water. The pH was adjusted to 10 using 10N sodium hydroxide. The layers were separated. The aqueous layer was cooled to 10°C and the pH adjusted to 2-3 with about 130 mL of 10 N hydrochloric acid. The oil which formed solidified and was collected by vacuum filtration, washed three times with water and dried under vacuum at 50°C to give 52.8 g (93% yield) of 3-(2-formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid as a

WO 00/08202

216

PCT/US99/17845

dark brown solid.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.1 (s, 1H, pyrrole NH), 11.7 (br s, 1H, COOH), 9.4 (s, 1H, CHO), 2.8, 2.5 (t, 4H, -CH<sub>2</sub>CH<sub>2</sub>CO-), 2.4 (m, 4H, -CH<sub>2</sub>-, -CH<sub>2</sub>-), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-).

5        3-(2-Formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid (5.4 g), 3.6 g of 2-oxindole and 2.7 g of piperidine (or 2.2 g of pyrrolidine) in 25 mL of ethanol were refluxed for 4 hours. Upon slow addition of acetic acid (8 mL), a precipitate formed. The mixture was refluxed for 5 minutes and cooled to  
10        ambient temperature. The precipitate was collected by vacuum filtration and washed with 20 mL of ethanol. The solids were slurry-washed refluxing in 30 mL of ethanol, cooled, collected by vacuum filtration, washed with 30 mL of ethanol and dried under vacuum to give 5.5 g (68% yield) of 3-[2-(2-oxo-1,2-  
15        dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid as an orange solid: mp 263-265 °C.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.1 (s, 1H, pyrrole NH), 12.0 (br s, 1H, COOH), 10.7 (s, 1H, CONH), 7.7, 7.1, 6.9, 6.8 (m, each 4H, aromatic), 7.6 (s, 1H, -CH=), 2.9, 2.7 (t, each 4H, -CH<sub>2</sub>CH<sub>2</sub>CO-),  
20        2.4 (m, 4H, -CH<sub>2</sub>-, -CH<sub>2</sub>-), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-).

**Example 191: 3-[2-(5-Chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid**

3-(2-Formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic  
25        acid (5.4 g), 3.7 g of 5-chloro-2-oxindole and 2.7 g of piperidine (or 2.2 g of pyrrolidine) in 25 mL of ethanol was refluxed for 4 hours. Slow addition of acetic acid (8 mL) resulted in a precipitate. The mixture was refluxed for 5 minutes and cooled to ambient temperature. The precipitate  
30        was collected by vacuum filtration and washed with 20 mL of ethanol. The solids were slurry-washed by refluxing in 30 mL of ethanol, cooled, collected by vacuum filtration, washed with 30 mL of ethanol and dried under high vacuum to give 6.5

WO 00/08202

217

PCT/US99/17845

g (80% yield) 3-[2-(5-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid as an orange solid: mp 370-384 °C.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.3 (s, 1H, pyrrole NH), 12.0 (br s, 1H, COOH), 10.7 (s, 1H, CONH), 7.8, 7.1, 6.8 (m, each 3H, aromatic), 7.7 (s, 1H, -CH=), 2.9, 2.7 (t, each 4H, CH<sub>2</sub>CH<sub>2</sub>CO), 2.5 (m, 4H, -CH<sub>2</sub>-, -CH<sub>2</sub>-), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-).

**Example 192:** 3-[2-(5-Bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid

A mixture of 3-(2-Formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid (3.4 g), 2.7 g of 5-bromo-2-oxindole and 1.4 g of pyrrolidine in 25 mL of ethanol was refluxed for 4 hours. Upon slow addition of acetic acid (8 mL), a precipitate formed. The mixture was refluxed for 5 minutes and cooled to ambient temperature. The precipitate was collected by vacuum filtration and washed with 20 mL of ethanol. The solids were slurry-washed by refluxing in 30 mL of ethanol, cooled, collected by vacuum filtration, washed with 30 mL of ethanol and dried under vacuum to give 5.2 g (98% yield) of 3-[2-(5-bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid as a red-orange solid, mp 286-289 °C.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.3 (s, 1H, pyrrole NH), 12.0 (br s, 1H, COOH), 10.7 (s, 1H, CONH), 7.9, 7.1, 6.8 (m, each 3H, aromatic), 7.7 (s, 1H, -CH=), 2.9, 2.7 (t, each 4H, -CH<sub>2</sub>CH<sub>2</sub>CO-), 2.5 (m, 4H, -CH<sub>2</sub>-, -CH<sub>2</sub>-), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-).

**Example 193:** 3-[2-(4-Methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid

Diethyl oxalate (30 mL) in 20 mL of dry ether was added with stirring to 19 g of potassium ethoxide suspended in 50 mL of dry ether. The mixture was cooled in an ice bath and 20 mL



WO 00/08202

218

PCT/US99/17845

of 3-nitro-o-xylene in 20 mL of dry ether was slowly added. The thick dark red mixture was refluxed for 0.5 hr, concentrated, and treated with 10% sodium hydroxide until almost all of the solid dissolved. Hydrogen peroxide (30%)  
5 was then added to the dark red mixture until the red color changed to yellow. The mixture was treated alternately with 10% sodium hydroxide and 30% hydrogen peroxide until the dark red color was no longer present. The solid was filtered off and the filtrate acidified with 6 N hydrochloric acid. The  
10 resulting precipitate was collected by vacuum filtration, washed with water, and dried under vacuum to give 9.8 g (45% yield) of 1-methyl-6-nitrophenylacetic acid as an off-white solid. The solid was hydrogenated in methanol over 10% palladium on carbon to give 9.04 g of 4-methyl-2-oxindole as a  
15 white solid.

$^1\text{H}$  NMR ( $d_6$ -DMSO, 360 MHz)  $\delta$  10.27 (s, br, 1H, NH-1), 7.06 (t, 7.71 Hz, 1H, H-6), 6.74 (d, 7.73 Hz, H-5), 6.63 (d, 7.73 Hz, 1H, H-7), 3.36 (s, 2H,  $\text{CH}_2$ ), 2.18 (s, 3H,  $\text{CH}_3$ ).

A mixture of 3-(2-formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid (5.4 g), 3.2 g of 4-methyl-2-oxindole and  
20 2.7 g of piperidine in 25 mL of ethanol was refluxed for 4 hours. Upon slow addition of acetic acid (8 mL), a precipitate formed. The mixture was refluxed for 5 minutes and cooled to ambient temperature. The precipitate was  
25 collected by vacuum filtration and washed with 20 mL of ethanol. The solids were slurry-washed in 30 mL of refluxing ethanol, cooled, collected by vacuum filtration, washed with 30 mL of ethanol and dried under vacuum to give 6.2 g (80% yield) of 3-[2-(4-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic  
30 acid as an orange solid.

$^1\text{H}$  NMR ( $d_6$ -DMSO):  $\delta$  13.3 (s, 1H, pyrrole NH), 12.0 (br s, 1H, COOH), 10.7 (s, 1H, CONH), 7.7 (s, 1H, -CH=), 7.0, 6.8 (m,

WO 00/08202

219

PCT/US99/17845

each 2H, aromatic), 2.8, 2.7 (t, each 4H,  $-\text{CH}_2\text{CH}_2\text{CO}-$ ), 2.6 (s, 1H,  $\text{CH}_3$ ), 2.5 (m, 4H,  $-\text{CH}_2-$ ,  $-\text{CH}_2-$ ), 1.7 (m, 4H,  $-\text{CH}_2\text{CH}_2-$ ).

**Example 194:** 3-[2-(5-Methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid

A mixture of 3-(2-formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid (5.4 g), 3.2 g of 5-methyl-2-oxindole and 2.7 g of piperidine in 25 mL of ethanol was refluxed for 4 hours. Upon addition of acetic acid (8 mL), a precipitate formed. The mixture was refluxed for 5 minutes and cooled to ambient temperature. The precipitate was collected by vacuum filtration and washed with 20 mL of ethanol. The solids were slurry-washed with 30 mL of refluxing ethanol, cooled, collected by vacuum filtration, washed with 30 mL of ethanol and vacuum dried to give 6.2 g (80% yield) of 3-[2-(5-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid as an orange solid.

$^1\text{H}$  NMR ( $d_6$ -DMSO):  $\delta$  13.3 (s, 1H, pyrrole NH), 12.0 (br s, 1H, COOH), 10.7 (s, 1H, CONH), 7.7 (s, 1H,  $-\text{CH}=\text{}$ ), 7.0, 6.8 (m, each 3H aromatic), 2.8, 2.7 (t, each 4H,  $-\text{CH}_2\text{CH}_2\text{CO}-$ ), 2.6 (s, 1H,  $\text{CH}_3$ ), 2.5 (m, 4H,  $-\text{CH}_2-$ ,  $-\text{CH}_2-$ ), 1.7 (m, 4H,  $-\text{CH}_2\text{CH}_2-$ ).

MS: m/z 349.

**Example 195:** 3-[2-(6-Chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid

A mixture of 3-(2-formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid (5.4 g), 3.7 g of 6-chloro-2-oxindole and 2.7 g of piperidine in 25 mL of ethanol was refluxed for 4 hours. Upon slow addition of acetic acid (8 mL), a precipitate formed. The mixture was refluxed for 5 minutes and cooled to ambient temperature. The precipitate was collected by vacuum filtration and washed with 20 mL of ethanol. The solids were slurry-washed in 30 mL of refluxing

WO 00/08202

220

PCT/US99/17845 ;

ethanol, cooled, collected by vacuum filtration, washed with 30 mL of ethanol and vacuum dried to give 6.5 g (80%) of 3-[2-(6-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid as an orange solid.

- 5 <sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.3 (s, 1H, pyrrole NH), 12.0 (br s, 1H, COOH), 10.7 (s, 1H, CONH), 7.7, 7.0, 6.9 (m, each 3H, aromatic), 7.6 (s, 1H, -CH=), 2.9, 2.7 (t, each 4H, -CH<sub>2</sub>CH<sub>2</sub>CO-), 2.4 (m, 4H, -CH<sub>2</sub>-, -CH<sub>2</sub>-), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-).

MS: m/z 371.

- 10 **Example 196:** 3-[2-(6-Methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid

- A mixture of 3-(2-formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid (5.4 g), 3.6 g of 6-methoxy-2-oxindole and  
15 2.7 g of piperidine in 25 mL of ethanol was refluxed for 4 hours. Upon slow addition of acetic acid (8 mL), a precipitate formed. The mixture was refluxed for 5 minutes and cooled to ambient temperature. The precipitate was collected by vacuum filtration and washed with 20 mL of  
20 ethanol. The solids were slurry-washed in 30 mL of refluxing ethanol, cooled, collected by vacuum filtration, washed with 30 mL of ethanol and vacuum dried to give 6.4 g (80% yield) of 3-[2-(6-methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid as an orange  
25 solid, mp 263-266 °C.

- <sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.0 (s, 1H, pyrrole NH), 12.0 (s, 1H, COOH), 10.7 (s, 1H, CONH), 7.6, 6.5, 6.4 (m, each 3H, aromatic), 7.4 (s, 1H, -CH=), 3.7 (s, 3H, CH<sub>3</sub>), 2.9, 2.7 (t, each 4H, -CH<sub>2</sub>CH<sub>2</sub>CO-), 2.5 (m, 4H -CH<sub>2</sub>-, -CH<sub>2</sub>-), 1.7 (m, 4H,  
30 -CH<sub>2</sub>CH<sub>2</sub>-).

MS: m/z 365.

**Example 197:** N,N-Dimethyl-3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide

WO 00/08202

221

PCT/US99/17845 ,

3-[2-(2-Oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid (10 g) was dissolved in 100 mL of dimethylformamide. Carbonyldiimidazole (6.3 g) was added and the mixture stirred at ambient temperature for 1  
5 hour. Dimethylamine (2.7 g) and 30 mL of dimethylformamide were added and the stirring continued overnight at room temperature. Fifty mL of water was added to the mixture and stirring was continued for 10 minutes. The precipitate was collected by vacuum filtration, washed with 20 mL of water and  
10 then 20 mL of ethanol. The solid was slurry-washed in 30 mL of refluxing ethanol for 5 minutes and cooled to room temperature. The solid was collected by vacuum filtration, washed with 20 mL of ethanol and vacuum dried to give 8.9 g (83% yield) of N,N-dimethyl-3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide.  
15

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.1 (s, 1H, pyrrole NH), 10.7 (s, 1H, CONH), 7.7, 7.1, 6.9, 6.8 (m, each 4H, aromatic), 7.6 (s, 1H, -CH=), 3.3 (s, 6H, CH<sub>3</sub>), 2.9, 2.7 (t, each 4H, -CH<sub>2</sub>CH<sub>2</sub>CO-), 2.5 (m, 4H -CH<sub>2</sub>-, -CH<sub>2</sub>-), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-).

20 MS: m/z 364.

**Example 198: 3-[3-(3-Dimethylaminopropyl)-4,5,6,7-tetrahydro-1H-indol-2-ylmethylene]-1,3-dihydroindol-2-one**

3-(4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid (9.7 g) in 100 mL of tetrahydrofuran was stirred with 8.1 g of  
25 carbonyl diimidazole for 1 hour. Dimethylamine (2.5 g) was added and the mixture stirred for 2 hours. The solvent was evaporated and the residue taken up in ethyl acetate, washed with water, 0.1 N hydrochloric acid, water, and brine, dried over sodium sulfate and evaporated to give 7.7 g (70% yield)  
30 of 3-(4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid dimethyl amide.

3-(4,5,6,7-Tetrahydro-1H-indol-3-yl)-propionic acid dimethyl amide (7.7 g) and 13 g of borane-tetrahydrofuran

WO 00/08202

222

PCT/US99/17845 ,

complex in 50 mL of tetrahydrofuran was refluxed for 3 hours. The reaction was quenched with acetone and then water and evaporated to dryness. The residue was chromatographed on silica gel to give 2 g (20% yield) of 3-(3-dimethylamino-propyl)-4,5,6,7-tetrahydro-1H-indole as a yellow oil.

Dimethylformamide (0.8 g) and 13 mL of dichloromethane was cooled to -9 °C. Phosphorus oxychloride (1.7 g) was rapidly added via a dropping funnel. 3-(3-dimethylamino-propyl)-4,5,6,7-tetrahydro-1H-indole (2 g) was added in portions with vigorous stirring. The mixture was warmed to room temperature and then refluxed for 10 minutes. It was then cooled to 5°C, and diluted with 20 mL of water. The pH was adjusted to 10 with 10N sodium hydroxide. The layers were separated. The organic layer was washed with water and brine, dried over anhydrous sodium sulfate and evaporated to give 1.8 g (80%) of 3-(3-dimethylaminopropyl)-2-formyl-4,5,6,7-tetrahydro-1H-indole as a dark oil which solidified.

3-(3-Dimethylaminopropyl)-2-formyl-4,5,6,7-tetrahydro-1H-indole (1.8 g), 1 g of 2-oxindole and 0.1 g of piperidine in 10 mL of ethanol was refluxed for 4 hours and then cooled to room temperature. The precipitate which formed was collected by vacuum filtration and washed with 4 mL of ethanol. The solids were slurry-washed with 8 mL of refluxing ethanol, cooled, collected by vacuum filtration, washed with 3 mL of ethanol and vacuum dried to give 1.8 g (70% yield) of 3-[3-(3-dimethylaminopropyl)-4,5,6,7-tetrahydro-1H-indol-2-ylmethylene]-1,3-dihydroindol-2-one as an orange solid.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.1 (s, 1H, pyrrole NH), 10.7 (s, 1H, CONH), 7.6, 7.1, 7.0, 6.8 (m, each 4H, aromatic), 7.6 (s, 1H, -CH=), 3.3 (s, 6H, CH<sub>3</sub>), 2.7 (m, 4H, -CH<sub>2</sub>-, -CH<sub>2</sub>-), 2.5, 2.3, 1.4 (t, each 6H, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N-), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-).

MS: m/z 350.

WO 00/08202

223

PCT/US99/17845

**Example 199: 3-[2-(2-Oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide**

3-[2-(2-Oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid (10 g) was dissolved  
5 in 100 mL of dimethylformamide. Carbonyldiimidazole (6.3 g) was added and the mixture stirred at ambient temperature for 1 hour. Ammonia (1 g) in 30 mL of dimethylformamide was added and the stirring continued overnight. Fifty mL of water was added to the mixture and stirring was continued for 10  
10 minutes. The precipitate that formed was collected by vacuum filtration, washed with 20 mL of water and then 20 mL of ethanol. The solid was slurry-washed with 30 mL of refluxing ethanol for 5 minutes and cooled to room temperature. The solid was collected by vacuum filtration, washed with 20 mL of  
15 ethanol and vacuum dried to give 11 g (83% yield) of 3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.3 (s, 1H, pyrrole NH), 12.0 (br s, 1H, COOH), 10.7 (s, 1H, CONH), 7.6, 7.1, 6.9, 6.8 (m, each 4H, aromatic), 7.6 (s, 1H, -CH=), 7.2, 6.7 (s, each 2H, NH<sub>2</sub>), 2.9, 2.7 (t, 4H, -CH<sub>2</sub>CH<sub>2</sub>CO-), 2.4 (m, 4H, -CH<sub>2</sub>-, -CH<sub>2</sub>-), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-).

MS: m/z 336.

**Example 200: 3-[3-(3-Morpholin-4-yl-3-oxopropyl)-4,5,6,7-tetrahydro-1H-indol-2-ylmethylenel]-1,3-dihydroindol-2-one**

3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid (10 g) was dissolved  
in 100 mL of dimethylformamide. Carbonyldiimidazole (6.3 g) was added and the mixture stirred at ambient temperature for 1  
30 hour. Morpholine (5.2 g) in 30 mL of dimethylformamide was added and the stirring continued overnight. Fifty mL of water was added to the mixture and stirring was continued for 10 minutes. The precipitate was collected by vacuum filtration,

WO 00/08202

224

PCT/US99/17845

washed with 20 mL of water and then 20 mL of ethanol. The solid that formed was slurry-washed with 30 mL of refluxing ethanol for 5 minutes and cooled to room temperature. The solid was collected by vacuum filtration, washed with 20 mL of ethanol and vacuum dried to give 9.6 g (80%) of 3-[3-(3-morpholin-4-yl-3-oxopropyl)-4,5,6,7-tetrahydro-1H-indol-2-ylmethylene]-1,3-dihydroindol-2-one as an orange solid.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.3 (s, 1H, pyrrole NH), 12.0 (br s, 1H, COOH), 10.7 (s, 1H, CONH), 7.6, 7.1, 6.9, 6.8 (m, each 4H, aromatic), 7.6 (s, 1H, -CH=), 3.3 (multiplets, 8H, morpholine CH<sub>2</sub>), 2.9, 2.7 (t, each 4H, -CH<sub>2</sub>CH<sub>2</sub>CO-), 2.5 (m, 4H, CH<sub>2</sub>, CH<sub>2</sub>), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-).

MS: m/z 406.

**Example 201: N-Methyl-3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide**

3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid (10 g) was dissolved in 100 mL of dimethylformamide. Carbonyldiimidazole (6.3 g) was added and the mixture stirred at ambient temperature for 1 hour. Methyl amine (1.8 g) in 30 mL of dimethylformamide was added and the stirring continued overnight. Fifty mL of water was added to the mixture and stirring was continued for 10 minutes. The precipitate was collected by vacuum filtration, washed with 20 mL of water and then 20 mL of ethanol. The solid was slurry-washed in 30 mL of refluxing ethanol for 5 minutes and cooled to room temperature. The solid was collected by vacuum filtration, washed with 20 mL of ethanol and vacuum dried to give 8.3 g (80% yield) of N-methyl-3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide as an orange solid.

MS: m/z 350.

**Example 202: N-(2-Morpholin-4-ylethyl)-3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-**

WO 00/08202

225

PCT/US99/17845

**Example 202: N-(2-Morpholin-4-ylethyl)-3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide**

5           3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid (10 g) was dissolved in 100 mL of dimethylformamide. Carbonyldiimidazole (6.3 g) was added and the mixture stirred at ambient temperature for 1 hour. 4-(2-Aminoethyl)morpholine (7.7 g) and 30 mL of  
10 dimethylformamide were added and the stirring continued overnight. Fifty mL of water was added to the mixture and stirring was continued for 10 minutes. The precipitate was collected by vacuum filtration, washed with 20 mL of water and then 20 mL of ethanol. The solid was slurry-washed in 30 mL  
15 of refluxing ethanol for 5 minutes and cooled to room temperature. The solid was collected by vacuum filtration, washed with 20 mL of ethanol and vacuum dried to give 1 g (83%) of N-(2-morpholin-4-ylethyl)-3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-  
20 propionamide, mp 256-258 °C.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.1 (s, 1H, pyrrole NH), 10.7 (s, 1H, -CONH-), 7.7 (t, 1H, -CONHCH<sub>2</sub>-), 7.6, 7.1, 6.9, 6.8 (m, each 4H, aromatic), 7.5 (s, 1H, -CH=), 3.5 (t, each 4H, -CH<sub>2</sub>CH<sub>2</sub>-), 3.1, 2.8 (m, each 2H, -CH<sub>2</sub>NCH<sub>2</sub>-), 2.7, 2.5 (t, each 4H, -CH<sub>2</sub>CH<sub>2</sub>CO-), 2.3, 2.2 (m, each 4H, -NHCH<sub>2</sub>CH<sub>2</sub>N-), 2.2 (m, 4H, -CH<sub>2</sub>-, -CH<sub>2</sub>-), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-).

**Example 203: 3-[2-(2-Oxo-1,2-dihydropyrrolo[2,3-b]pyridin-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid**

30           A mixture of 7-azaoxindole (99 mg), 110 mg of 3-(2-formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid and 2 drops of piperidine in 2 mL of ethanol was refluxed for 5 h. The reaction mixture was cooled and concentrated. The residue



WO 00/08202

226

PCT/US99/17845

was acidified with acetic acid to pH 6. The resulting precipitate was collected by vacuum filtration, washed with water and dried in a vacuum oven at 40 °C to give 25.4 mg (13% yield) of 3-[2-(2-oxo-1,2-dihydropyrrolo[2,3-b]pyridin-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid.

MS: m/z 338.

**Example 204:** 3-{2-[6-(3-Methoxyphenyl)-2-oxo-1,2-dihydro-indol-3-ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl}-propionic acid

A mixture of 103 mg 6-(3-methoxyphenyl)-2-oxindole, 95 mg 3-(2-formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid and piperidine (3 drops) in ethanol (2 mL) was heated in sealed tube to 90°C and held there overnight. The reaction mixture was concentrated and acidified with 6 N hydrochloric acid. The precipitate that formed was collected by filtration and washed with water and hexane to give 156 mg of 3-{2-[6-(3-methoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl}-propionic acid as a brown solid (82%).

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO, 360 MHz): δ 13.26 (s, br, 1H, NH-1), 1.78 (s, br, 1H, NH-1), 7.72 (d, 8.1 Hz, 1H, H-4), 7.65 (s, 1H, H-vinyl), 7.35 (d, 7.9 Hz, 1H), 7.26 (dd, 1.3 Hz, 8.1 Hz, 1H, H-5), 7.18 (d, 7.9 Hz, 1H), 7.13 (t, 2.0 Hz, 1H), 7.09 (d, 1.3 Hz, 1H, H-7), 6.90 (dd, 2.0 Hz, 1H), 3.82 (s, 3H, OCH<sub>3</sub>), 2.91 (t, 7.4 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>COOH), 2.66 (t, 5.9 Hz, 2H, H-7'), 2.38-2.46 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>COOH and H-4'), 1.69-1.76 (m, 4H, H-5', 6').

MS: m/z 443.2.

**Example 205:** 3-{2-[6-(4-Methoxyphenyl)-2-oxo-1,2-dihydro-indol-3-ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl}-propionic acid

A mixture of 103 mg 6-(4-methoxyphenyl)-2-oxindole, 95 mg 3-(2-formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid

WO 00/08202

227

PCT/US99/17845

and piperidine (3 drops) in ethanol (2 mL) was heated to 90°C in a sealed tube and held there for 4 hrs. The reaction mixture was concentrated and acidified with 6 N hydrochloric acid. Ethyl acetate was added upon which a solid precipitated from the aqueous layer. The precipitate was collected by filtration, and washed with water and hexane to give 57 mg of 3-{2-[6-(4-Methoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl}-propionic acid as a brown solid (30%).

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO, 360 MHz): δ 13.24 (s, br, 1H, NH-1), 11.61 (s, br, 1H, COOH), 10.76 (s, br, 1H, NH-1), 7. (d, 8.1 Hz, 1H, H-4), 7.61 (s, 1H, H-vinyl), 7.56 (d, 8.8 Hz, 2H, H-3, 5), 7.21 (dd, 1.5 Hz, 8.1 Hz, 1H, H-5), 7.04 (d, J = 1.5 Hz, 1H, H-7), 7.01 (d, 8.8 Hz, 2H, H-2", 6" ), 3.79 (s, 3H, OCH<sub>3</sub>), 2.91 (t, 7.4 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>COOH ), 2.67 (t, 5.9 Hz, 2H, H-7'), 2.40-2.46 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>COOH and H-4'), 1.72-1.78 (m, 4H, H-5', 6').

MS: m/z 441.2.

**Example 206: 3-[2-(2-Oxo-6-phenyl-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid**

A mixture of 90 mg 6-phenyl-2-oxindole, 95 mg 3-(2-formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid and piperidine ( 3 drops) in ethanol (2 mL) was held in a sealed tube at 90 °C for 4 hrs. The reaction mixture was concentrated and acidified with 6 N hydrochloric acid. The precipitate was collected by filtration and washed with water and hexane to give 59 mg of 3-[2-(2-oxo-6-phenyl-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid as a brown solid (31% yield).

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO, 360 MHz): δ 13.27 (s, br, 1H, NH-1'), 12.06 (s, v br, 1H, COOH), 10.80 (s, br, 1H, NH-1), 7.74 (d, 7.9 Hz, 1H, H-4), 7.64 (s, 1H, H-vinyl), 7.62 (d, 7.7 Hz, 2H),

WO 00/08202

228

PCT/US99/17845

7.44 (t, 7.7 Hz, 2H), 7.32 (dd, 7.7 Hz, 1H), 7.27 (dd, 1.1, 7.9 Hz 1H, H-5), 7.10 (d, 1.1 Hz, 1H, H-7), 2.92 (t, 7.3 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>COOH), 2.67 (t, 5.5 Hz, 2H, H-7'), 2.41-2.46 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>COOH and H-4'), 1.73-1.76 (m, 4H, H-5', 6').

5 MS: m/z 411.2.

**Example 207: 3-{2-[6-(2-Methoxyphenyl)-2-oxo-1,2-dihydro-indol-3-ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl}-propionic acid**

Tetrakis(triphenylphosphine)palladium (1 g) was added to  
10 a mixture of 5 g of 2-methoxyphenylboronic acid, 6.6 g of 5-bromo-2-fluoronitrobenzene and 30 mL of 2 M sodium carbonate solution in 50 mL of toluene and 50 mL of ethanol. The mixture was refluxed for 2 hours, concentrated, and the residue extracted twice with ethyl acetate. The ethyl acetate  
15 layer was washed with water and brine, dried, and concentrated to give a dark green oil which solidified on standing to give crude 4-fluoro-2'-methoxy-3-nitrobiphenyl.

Dimethyl malonate (14 mL) was added dropwise to 2.9 g of sodium hydride suspended in 50 mL of dimethylsulfoxide. The  
20 mixture was stirred at 100°C for 15 minutes and cooled to room temperature. Crude 4-fluoro-2'-methoxy-3-nitrobiphenyl in 60 mL of dimethylsulfoxide was added and the mixture stirred at 100 °C for 2 hours. The reaction mixture was cooled, quenched with 300 mL of saturated ammonium chloride solution and  
25 extracted twice with ethyl acetate. The extracts were combined, washed with saturated ammonium chloride, water, and brine, dried over anhydrous sodium sulfate and concentrated to give crude dimethyl 2'-methoxy-3-nitrobiphenyl-4-malonate as a yellow oil.

30 Crude 2'-methoxy-3-nitrobiphenyl-4-malonate was stirred at 100 °C in 50 mL of 6 N hydrochloric acid for 24 hours and cooled. The precipitate that formed was collected by filtration, washed with water and hexane, and dried to give 9.8 of 2'-methoxy-2-nitrobiphenyl-4-acetic acid as a light tan

WO 00/08202

229

PCT/US99/17845

solid.

Iron chips (5 g) were added in one portion to 9.8 g of 2'-methoxy-3-nitrobiphenyl-4-acetic acid in 50 mL of glacial acetic acid and the mixture was stirred at 100 °C for 3 hours.

5 The reaction mixture was concentrated to dryness, sonicated in ethyl acetate and filtered to remove insolubles. The filtrate was washed twice with 1 N hydrochloric acid, then with water and brine, dried over anhydrous sodium sulfate and concentrated. The residue was chromatographed on silica gel  
10 using ethyl acetate:hexane 1:2 as the eluent to give 5.4 g (69% based on 5-bromo-2-fluoronitrobenzene) of 6-(2-methoxyphenyl)-2-oxindole as a rose colored solid.

A mixture of 103 mg of 6-(2-methoxyphenyl)-2-oxindole, 95 mg of 3-(2-formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic  
15 acid and piperidine (3 drops) in ethanol (2 mL) was held in a sealed tube at 90 °C for 4 hrs. The reaction mixture was concentrated and acidified with 6 N hydrochloric acid. The precipitate was collected by filtration and washed with water and hexane to give 67 mg of 3-{2-[6-(2-methoxyphenyl)-2-oxo-  
20 1,2-dihydroindol-3-ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl}-propionic acid as a brown solid (35% yield).

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO, 360 MHz): δ 13.26 (s, br, 1H, H1, 20, vr H OH) 10.71 (s, br, 1H, NH-1), 7.67 (d, 7.7 Hz, 1H, H-4), 7.61 (s, 1H, H-vinyl), 7.27-7.34 (m, 2H, ), 7.01-7.10 (m, 2H), 7.05  
25 (dd, 1.2, 7.7 Hz, 1H, H-5), 6.99 (d, 1.2 Hz, 1H, H-7), 3.75 (s, 3H, OCH<sub>3</sub>), 2.91 (t, 7.5 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>COOH), 2.68 (t, 7 Hz, 2H, H-7'), 2.40-2.46 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>COOH and H-4'), 1.71-1.78 (m, 4H, H-5',6').

MS m/z 441.2.

30 **Example 208:** 3-[2-(5-Isopropylaminosulfonyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid

To a 100 mL flask charged with 27 mL of chlorosulfonic

WO 00/08202

230

PCT/US99/17845

acid was slowly added 13.3 g of 2-oxindole. The reaction temperature was maintained below 30 °C during the addition. After the addition, the reaction mixture was stirred at room temperature for 1.5 hr, heated to 68°C and stirred an  
5 additional 1 hr, cooled, and poured into water. The precipitate was washed with water and dried in a vacuum oven to give 11.0 g of 5-chlorosulfonyl-2-oxindole (50% yield) which was used without further purification.

A suspension of 3 g of 5-chlorosulfonyl-2-oxindole, 1.15  
10 g of isopropylamine and 1.2 mL of pyridine in 50 mL of dichloromethane was stirred at room temperature for 4 hours at which time a white solid was present. The solid was collected by vacuum filtration, slurry-washed with hot ethanol, cooled, collected by vacuum filtration and vacuum dried at 40°C to  
15 give 1.5 g (45%) of 5-isopropylaminosulfonyl-2-oxindole.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO, 300 MHz) δ 10.69 (s, br, 1H, NH), 7.63 (dd, 1.8 Hz, 1H, H-6), 7.59 (d, 1 Hz, 1H, H-4), 7.32 (d, 7 Hz, 1H, NH-SO<sub>2</sub>-), 6.93 (d, 8 Hz, 1H, H-7), 3.57 (s, 2H, H-3), 3.14-3.23 (m, 1H, CH-(CH<sub>3</sub>)<sub>2</sub>), 0.94 (d, 7 Hz, 6H, CH<sub>3</sub>).

A mixture of (2-Formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid (5.4 g), 5.7 g of 5-isopropylaminosulfonyl-2-oxindole and 2.7 g of piperidine in 25 mL of ethanol was refluxed for 4 hours. Upon addition of acetic acid (8 mL), a precipitate formed. The mixture was refluxed for 5 minutes  
25 and cooled to ambient temperature. The precipitate was collected by vacuum filtration and washed with 20 mL of ethanol. The solids were slurry-washed in 30 mL of refluxing ethanol, cooled, collected by vacuum filtration, washed with 30 mL of ethanol and vacuum dried to give 8.1 g (80% yield) of  
30 3-[2-(5-isopropylaminosulfonyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid as an orange solid.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.3 (s, 1H, pyrrole NH), 12.0 (br s,

WO 00/08202

231

PCT/US99/17845,

1H, COOH), 10.7 (s, 1H, CONH), 8.1, 7.5, 7.3, 7.0 (m, each 4H, aromatic), 7.7 (s, 1H, -CH=), 2.3 (m, 1H, -CH-), 2.9, 2.7 (t, each 4H, -CH<sub>2</sub>CH<sub>2</sub>CO-), 2.4 (m, 4H, -CH<sub>2</sub>-, -CH<sub>2</sub>-), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-), 0.9 (d, 6H, CH<sub>3</sub>).

5 MS m/z 458.

**Example 209:** 3-[2-(6-Morpholin-4-yl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid

A mixture of 4 g of 6-(morpholin-4-yl)-2-oxindole, 3.75 g  
10 of 3-(2-formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)propionic acid, and 1.8 mL of piperidine in ethanol (60 mL) was refluxed for 6 hrs. The reaction mixture was concentrated and acidified with 6 N hydrochloric acid to pH 6. The precipitate was collected by filtration, washed once with water, twice  
15 with ethyl acetate and twice with methanol to give 2.6 g of 3-[2-(6-Morpholin-4-yl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]propionic acid as an orange solid (34% yield).

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO, 360 MHz): δ 13.04 (s, br, 1H, NH-1'),  
20 12.05 (s, vbr, 1H, COOH), 10.60 (s, br, 1H, NH-1), 7.50 (d, 8.0 Hz, 1H, H-4), 7.39 (s, 1H, H-vinyl), 6.60 (d, 8.0 Hz, 1H, H-5), 6.43 (s, 1H, H-7), 3.73 (d, 4.7, 4H, H-2'', 6''), 3.09 (s, 4H, H-3'', 5''), 2.86 (t, 7.1 Hz, 2H, CH<sub>2</sub>CH<sub>2</sub>COOH), 2.64 (s, br, 2H, H-7'), 2.37-2.43 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>COOH and H-4'),  
25 1.71-1.75 (m, 4H, H-5', 6').

MS m/z 422.3.

**Example 210:** 3-[2-(5-Chloro-4-methyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid

30 A suspension of 3.0 g of 4-methyl-2-oxindole was stirred in 50 mL of acetonitrile at room temperature while 3.3 g of N-chlorosuccinimide was added in portions. Trifluoroacetic acid (1 mL) was then added. A white precipitate formed. The

WO 00/08202

232

PCT/US99/17845

suspension was stirred at room temperature for 3 days. The white solid was collected by vacuum filtration, washed with a small amount of cold acetone and dried overnight in a vacuum oven at 40°C to give 2.5 g (68%) of 5-chloro-4-methyl-2-oxindole.

A mixture of (2-Formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid (5.4 g), 4.0 g of 5-chloro-4-methyl-2-oxindole and 2.7 g of piperidine in 25 mL of ethanol was refluxed for 4 hours. Addition of acetic acid (8 mL) resulted in the formation of a precipitate. The mixture was refluxed for 5 minutes and cooled to ambient temperature. The precipitate was collected by vacuum filtration and washed with 20 mL of ethanol. The solids were slurry-washed with 30 mL of refluxing ethanol, cooled, collected by vacuum filtration, washed with 30 mL of ethanol and vacuum dried to give 6.8 g (80%) of 3-[2-(5-Chloro-4-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid as an orange solid.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.1 (s, 1H, pyrrole NH), 12.0 (br s, 1H, COOH), 10.7 (s, 1H, CONH), 7.6 (s, 1H, -CH=), 7.3, 6.7 (m, each 2H, aromatic), 2.9, 2.7 (t, each 4H, -CH<sub>2</sub>CH<sub>2</sub>CO-), 2.7 (s, 3H, CH<sub>3</sub>), 2.4 (m, 4H, -CH<sub>2</sub>-, -CH<sub>2</sub>-), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-).

**Example 211:** 3-[2-(5-Bromo-4-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid

A mixture of (2-Formyl-4,5,6,7-tetrahydro-1H-indol-3-yl)-propionic acid (5.4 g), 5.0 g of 5-bromo-4-methyl-2-oxindole and 2.7 g of piperidine in 25 mL of ethanol was refluxed for 4 hours. Acetic acid (8 mL) was slowly added and a precipitate formed. The mixture was refluxed for 5 minutes and cooled to ambient temperature. The precipitate was collected by vacuum filtration and washed with 20 mL of ethanol. The solids were slurry-washed in 30 mL of refluxing ethanol, cooled, collected

WO 00/08202

233

PCT/US99/17845 ,

by vacuum filtration, washed with 30 mL of ethanol and vacuum dried to give 7.6 g (80%) of 3-[2-(5-bromo-4-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid as an red-orange solid.

5       <sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 13.1 (s, 1H, pyrrole NH), 12.0 (br s, 1H, COOH), 10.7 (s, 1H, CONH), 7.8 (s, 1H, -CH=), 7.3, 6.7, (m, each 2H, aromatic), 2.9, 2.7 (t, each 4H, -CH<sub>2</sub>CH<sub>2</sub>CO-), 2.7 (s, 3H, CH<sub>3</sub>), 2.4 (m, 4H -CH<sub>2</sub>-, -CH<sub>2</sub>-), 1.7 (m, 4H, -CH<sub>2</sub>CH<sub>2</sub>-).

MS: m/z 427, 429.

10   **Example 212: 3-[2-(5-Bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-N-(2-morpholin-4-ylethyl)-propionamide**

3-[2-(5-Bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid (12.3 g) was dissolved in 150 mL of dimethylformamide. Carbonyldiimidazole (6.3 g) was added and the mixture stirred at ambient temperature for 1 hour. 4-(2-Aminoethyl)morpholine (7.7 g) and 30 mL of dimethylformamide were added and the stirring continued overnight at room temperature. Fifty mL of water was added to the mixture and stirring was continued for 10 minutes. The precipitate was collected by vacuum filtration, washed with 20 mL of water and then 20 mL of ethanol. The solid was slurry-washed in 30 mL of refluxing ethanol for 5 minutes and cooled to room temperature. The solid was collected by vacuum filtration, washed with 20 mL of ethanol and vacuum dried to give 12.5 g (80%) of 3-[2-(5-bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-N-(2-morpholin-4-ylethyl)-propionamide.

25   **Example 213: 3-[2-(5-Chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-N-(2-morpholin-4-ylethyl)-propionamide**

3-[2-(5-Chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid (11 g) was



WO 00/08202

234

PCT/US99/17845 ;

dissolved in 150 mL of dimethylformamide. Carbonyldiimidazole (6.3 g) was added and the mixture stirred at ambient temperature for 1 hour. 4-(2-Aminoethyl)morpholine (7.7 g) and 30 mL of dimethylformamide were added and the stirring continued overnight at room temperature. Fifty mL of water was added to the mixture and stirring was continued for 10 minutes. The precipitate was collected by vacuum filtration, washed with 20 mL of water and then 20 mL of ethanol. The solid was slurry-washed in 30 mL of refluxing ethanol for 5 minutes and cooled to room temperature. The solid was collected by vacuum filtration, washed with 20 mL of ethanol and vacuum dried to give 11.5 g (80%) of 3-[2-(5-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-N-(2-morpholin-4-ylethyl)-propionamide.

15 **Example 214: 3-[2-(2-Oxo-1,2-dihydroindol-3-ylidenemethyl)-phenyl]-propionic acid**

To a solution of 13.4 g of phthalic dicarboxaldehyde in 100 mL of dichloromethane was added in portions over 5 minutes 35 g of ethyl (triphenyl-phosphoranylidene)acetate. The reaction mixture was stirred at room temperature overnight and concentrated. The residue was stirred in 500 mL of a 6:1 mixture of hexanes:ethyl acetate for 1 hr. The solids were removed by filtration and the filtrates concentrated. The product was chromatographed on a silica gel column to give 10 g of ethyl 3-(2-formylphenyl)propionate as a mixture of *E* and *Z* isomers.

The above mixture was dissolved in 100 mL of ethyl acetate containing 100 mg of 5% palladium on carbon and stirred under hydrogen (balloon pressure) for 10 hrs. The mixture was filtered through a bed of celite which was washed with ethyl acetate. The combined filtrates were concentrated to give 8 g of ethyl 3-(2-formylphenyl)propionate.

A mixture of 1 g of ethyl 3-(2-formylphenyl)propionate, 500 mg oxindole and 0.1 mL piperidine in 5 mL of ethanol was

WO 00/08202

235

PCT/US99/17845 ,

stirred at 90 °C overnight. The mixture was evaporated and purified on a silica gel column to give 450 mg of ethyl (E)-3-[(2-oxo-1,2-dihydroindol-3-ylidenemethyl)phenyl]propionate (M+1 = 322).

5 To 300 mg of ethyl (E)-3-[(2-oxo-1,2-dihydroindol-3-ylidenemethyl)phenyl]-propionate in 3 mL of ethanol was added 2 mL of 2N sodium hydroxide. The mixture was stirred at 90°C for 2 hrs, cooled and acidified with 6 N hydrochloric acid to pH 3. The solid was collected by filtration and washed with  
10 cold ethanol to give 30 mg of 3-[2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-phenyl]-propionic acid as a yellow solid.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO): δ 12.1 (s, 1H, COOH), 10.6 (s, 1H, CONH), 7.7 (s, 1H, =CH) 7.6, 7.4, 7.3, 7.2, 7.0, 6.8, 6.7 (m, 8H, aromatic), 2.9, 2.5 (t, each 4H, CH<sub>2</sub>CH<sub>2</sub>).

15 MS: m/z 294.

**Example 215: 3-[4-Methyl-2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrol-3-yl]-propionic acid**

3,5-Dimethyl-4-(2-methoxycarbonylethyl)-1H-pyrrole-2-carboxylic acid ethyl ester (127 g) was dissolved in acetic  
20 acid (1900 mL), water (1900 mL) and tetrahydrofuran (1900 mL) and cooled to -30 °C. Ceric ammonium nitrate (1097 g) was added in portions with stirring to give a reddish-orange suspension. The suspension was stirred at 0 °C for 2 hours, neutralized to pH 7 with sodium bicarbonate and extracted with  
25 ethyl acetate (2000 mL). The ethyl acetate layer was separated, washed with brine (200 mL) and dried over anhydrous sodium sulfate (20 g). The solvent was removed to give 80.2 g (60%) of 5-formyl-4-(2-methoxycarbonylethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester as an oil.

30 5-Formyl-4-(2-methoxycarbonylethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester (80.2 g), 2-oxindole (37.9 g) and ethanol (300 mL) were warmed to 70 °C in a 500 mL, 3-neck round bottom flask equipped with mechanical stirring and a

WO 00/08202

236

PCT/US99/17845

reflux condenser. Piperidine (1.3 g) was added and the mixture was refluxed for 4 hours. The mixture was cooled to 10°C and the orange precipitate collected by vacuum filtration and washed with 30 mL of ethanol. The solid was slurry-washed in 150 mL of refluxing ethanol, cooled, collected by vacuum filtration, washed with 30 mL of ethanol and vacuum dried to give 81.7 g (75%) of 4-(2-Methoxycarbonylethyl)-3-methyl-5-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester as an orange solid.

4-(2-Methoxycarbonylethyl)-3-methyl-5-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester (81.7 g), 56.5 g of potassium hydroxide, 200 mL of ethanol and 200 mL of water were charged to a 1 L, 3 neck round bottom flask equipped with mechanical stirring and a thermometer. The mixture was stirred at 90°C for 90 minutes, cooled to room temperature, and acidified with acetic acid until a precipitate formed. The precipitate was collected by vacuum filtration, washed with 50 mL of water and vacuum dried to give 69.1 g (85 %) of 4-(2-carboxyethyl)-3-methyl-5-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid as a red solid.

4-(2-carboxyethyl)-3-methyl-5-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid (10 g) suspended in 50 mL of ethylene glycol (b.p. 198 °C) was sealed in a 1 L pressure reactor, the reactor was pressurized to 120 psi with nitrogen and then heated to 150 °C for 3 hours. The reaction mixture was cooled to room temperature and then diluted with 50 mL of water. The resulting precipitate was collected by vacuum filtration and was washed twice with 100 mL of water to give a mixture of 3-[4-methyl-2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrol-3-yl]-propionic acid and 3-[4-methyl-2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrol-3-yl]-propionic acid 2-hydroxyethyl ester as a dark orange solid.

WO 00/08202

237

PCT/US99/17845

The solid was used in the next step without drying.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 13.3 (s, br, 1H, NH), 10.77 (s, 1 H, NH), 7.6 (s, 1 H, H-vinyl), 7.67, 7.08, 6.97, 6.85 (m, 4 H, Ar-H), 4.73 (t, J = 6Hz, 1 H, OH), 3.97-4.0 (m, 2H, CH<sub>2</sub>), 3.5-  
5 3.55 (m, 2 H, CH<sub>2</sub>), 2.98 (t, J = 7.5 Hz, 2 H, CH<sub>2</sub>), 2.51 (t, J = 7.5 Hz, 2 H, CH<sub>2</sub>), 2.04 (s, 3 H, CH<sub>3</sub>).

MS m/z 341 (M+1).

3-[4-methyl-2-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-  
1H-pyrrol-3-yl]-propionic acid and 3-[4-methyl-2-(2-oxo-1,2-  
10 dihydroindol-3-ylidenemethyl)-1H-pyrrol-3-yl]-propionic acid  
2-hydroxyethyl ester, 1.9 g of potassium hydroxide, 50 mL of  
water, and 50 mL of ethanol in a 500 mL 3 neck round bottom  
flask were stirred at 70°C for 1 hour. The mixture was  
cooled to room temperature and acidified with 2 N  
15 hydrochloric acid until a precipitate formed. The  
precipitate was collected by vacuum filtration and washed  
with ethanol:water mixture (1:1, 100mL). The solid was  
slurry-washed with ethyl acetate:ethanol mixture (1:1, 100  
mL) at 70°C for 30 minutes and then cooled to room  
20 temperature. The product was collected by vacuum filtration  
and vacuum dried at 40°C overnight to give 7.8 g (90%  
overall yield) of 3-[4-methyl-2-(2-oxo-1,2-dihydroindol-3-  
ylidenemethyl)-1H-pyrrol-3-yl]-propionic acid as a dark  
orange solid. (R. B. Woodward et al., Tetrahedron, 1990, 46  
25 (22), 7599-7659)

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 13.28 (s, br, 1 H, NH), 12.05 (s, 1 H, COOH), 10.78 (s, 1 H, NH), 7.68 (d, J = 7 Hz, 1 H, Ar-H),  
7.64 (s, 1 H, H-vinyl), 7.11 (t, J = 7 Hz, 1 H, Ar-H), 7.11  
(s, 1 H), 6.97 (t, J = 7 Hz, 1 H, Ar-H), 6.86 (d, J = 7 Hz,  
30 1 H, Ar-H), 2.94 (t, J = 7.5 Hz, 2 H, CH<sub>2</sub>), 2.41 (t, J = 7.5  
Hz, 2 H, CH<sub>2</sub>), 2.04 (s, 3H, CH<sub>3</sub>).

**Example 216:** 3-[2-(5-Chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid

WO 00/08202

238

PCT/US99/17845 ,

5-Formyl-4-(2-ethoxycarbonylethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester (281 mg), 5-chloro-2-oxindole (168 mg), and piperidine (2 drops) in ethanol (2 mL) were refluxed for 2 hours. The reaction mixture was cooled and the precipitate that formed was filtered, washed with ethanol and hexanes, and dried to give 369 mg (86%) of 4-(2-ethoxycarbonylethyl)-3-methyl-5-(5-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester as a light yellow needle crystals.

10 A mixture of 4-(2-ethoxycarbonylethyl)-3-methyl-5-(5-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid ethyl ester (346 mg) and potassium hydroxide (560 mg) in ethanol (5 mL) was heated to 95°C and then cooled, upon which red crystals formed. The crystals were dissolved in water and acidified with 2 N hydrochloric acid to pH 2. The precipitate was filtered, washed with water, and dried in a vacuum oven overnight to give 299 mg (100%) of 4-(2-carboxyethyl)-3-methyl-5-(5-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid as a brown solid.

4-(2-carboxyethyl)-3-methyl-5-(5-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid suspended in ethylene glycol (5 mL) was heated in a sealed tube in a pre-heated oil bath at 200 °C for 2 hours. The reaction mixture was cooled to 90 °C and potassium hydroxide (2 pellets) were added. The mixture was then heated at 90°C for 30 minutes, after which time it was cooled, poured into water, and acidified with 2 N hydrochloric acid to pH 2. The precipitate that formed was filtered, washed with water, and dried in a vacuum oven overnight to give 77 mg (29%) of 3-[2-(5-Chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ 13.31 (s, br. 1 H, NH), 12.05 (s, 1 H,

WO 00/08202

239

PCT/US99/17845

COOH), 10.89 (s, br, 1 H, NH), 7.85 (d,  $J = 2$  Hz, 1 H, H-4),  
7.75 (s, 1 H, H-vinyl), 7.16 (d,  $J = 3$  Hz, 1 H), 7.11 (dd,  $J$   
= 2.8 Hz, 1 H, H-6), 6.84 (d,  $J = 8$  Hz, 1 H, H-7), 2.97 (t,  
 $J = 7.5$  Hz, 2 H, CH<sub>2</sub>), 2.41 (t,  $J = 7.5$  Hz, 2 H, CH<sub>2</sub>), 2.04  
5 (s, 3 H, CH<sub>3</sub>).

MS  $m/z$  331 (M+1).

**Example 217: 3-[2-(6-Methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid**

5-Formyl-4-(2-ethoxycarbonyl-ethyl)-3-methyl-1H-pyrrole-  
10 2-carboxylic acid ethyl ester (591 mg), 6-methoxy-2-oxindole  
(333 mg) and piperidine (0.1 mL) in ethanol (4 mL) were  
stirred at 90°C for 2 hours. Potassium hydroxide (537 mg)  
was added to the mixture and it was then stirred at 95°C for  
and additional hour. The reaction mixture was cooled and  
15 concentrated, dissolved in water, and acidified with 2 N  
hydrochloric acid to pH 2. The precipitate was filtered,  
washed with water, and dried in a vacuum oven overnight to  
give 730 mg (99%) of 4-(2-carboxyethyl)-3-methyl-5-(6-  
methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-  
20 2-carboxylic acid.

4-(2-carboxyethyl)-3-methyl-5-(6-methoxy-2-oxo-1,2-  
dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid  
(501 mg) suspended in ethylene glycol (10 mL) was heated in  
a sealed tube in a pre-heated oil bath at 200°C for 2 hours.  
25 The reaction mixture was cooled to 90 °C and potassium  
hydroxide (2 pellets) was added. The mixture was then  
stirred at 90°C for 30 minutes, after which time it was  
cooled, poured into water, and acidified with 2 N  
hydrochloric acid to pH 2. The precipitate was filtered,  
30 washed with water, and dried in a vacuum oven overnight to  
give 221 mg (48%) of 3-[2-(6-methoxy-2-oxo-1,2-dihydro-  
indol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic  
acid.

WO 00/08202

240

PCT/US99/17845

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ 13.09 (s, 1 H, NH), 12.05 (s, 1 H, COOH), 10.74 (s, 1 H, NH), 7.58 (d, J = 8 Hz, 1 H, H-4), 7.48 (s, 1 H, H-vinyl), 7.04 (d, J = 2 Hz, 1 H), 6.56 (dd, J = 2.8 Hz, 1 H, H-5), 6.43 (d, J = 2 Hz, 1 H), 3.75 (s, 3 H, OCH<sub>3</sub>), 2.91 (t, J = 7.5 Hz, 2 H, CH<sub>2</sub>), 2.4 (t, J = 7.5 Hz, 2 H, CH<sub>2</sub>), 2.03 (s, 3 H, CH<sub>3</sub>).

MS m/z 325 (M-1).

**Example 218: 3-[2-(4-Methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid**

10 A mixture of 5-formyl-4-(2-ethoxycarbonylethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester (281 mg), 4-methyl-2-oxindole (147 mg) and piperidine (2 drops) in ethanol (2 mL) was stirred at 90°C for 2 hours. Potassium hydroxide (213 mg) was added, the temperature was increased  
15 to 95°C and held there for 1 hour. The reaction mixture was cooled and concentrated. The residue was dissolved into water and acidified with 2 N hydrochloric acid to pH 2. The precipitate was filtered, washed with water and dried in a vacuum oven overnight to give 337 mg (95%) of 4-(2-  
20 carboxyethyl)-3-methyl-5-(4-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid.

4-(2-carboxyethyl)-3-methyl-5-(4-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid (300 mg) suspended in ethylene glycol (5 mL) was heated in a  
25 sealed tube in a pre-heated oil bath at 200 °C for 2 hours. The reaction mixture was cooled to 90°C and potassium hydroxide (1 pellet) was added. It was then stirred at 90°C for 30 minutes. The reaction mixture was cooled, poured into water, and acidified with 2 N hydrochloric acid to pH 2. The  
30 precipitate was filtered, washed with water, and purified on a silica gel column to give 115 mg (44% yield) of 3-[2-(4-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid.

WO 00/08202

241

PCT/US99/17845

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ 13.37 (s, 1 H, NH), 12.04 (s, 1 H, COOH), 10.81 (s, 1 H, NH), 7.69 (s, 1 H, H-vinyl), 7.09 (d, J = 2.5 Hz, 1 H), 7.01 (t, J = 7.5 Hz, 1 H, Ar-H), 6.79 (d, J = 7.5 Hz, 1 H, Ar-H), 6.74 (d, J = 7.5 Hz, 1 H, Ar-H), 2.88 (t, J = 7.2 Hz, 2 H, CH<sub>2</sub>), 2.61 (s, 3 H, CH<sub>3</sub>-4), 2.44 (t, J = 7.2 Hz, 2 H, CH<sub>2</sub>), 2.04 (s, 3 H, CH<sub>3</sub>).

MS m/z 309 (M-1).

**Example 219: 3-[2-(6-Chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid**

10 5-Formyl-4-(2-ethoxycarbonyl-ethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester (281 mg), 6-chloro-2-oxindole (168 mg) and piperidine (2 drops) in ethanol (2 mL) were stirred at 90 °C for 2 hours. Potassium hydroxide (537 mg) was added to the mixture and it was then stirred at 95°C for  
15 and additional hour. The reaction mixture was cooled and concentrated. The residue was dissolved into water and acidified with 2 N hydrochloric acid to pH 2. The precipitate was filtered, washed with water, and dried in a vacuum oven overnight to give 270 mg (72%) of 4-(2-carboxy-  
20 ethyl)-3-methyl-5-(6-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid.

4-(2-carboxyethyl)-3-methyl-5-(6-chloro-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid (240 mg) suspended in ethylene glycol (5 mL) was held in a  
25 sealed tube in a pre-heated oil bath at 200 °C for 2 hours. The reaction mixture was cooled to 90°C and potassium hydroxide (2 pellets) was added. It was then stirred at 90°C for 30 minutes. The reaction mixture was cooled, poured into water, and acidified with 2 N hydrochloric acid to pH 2. The  
30 precipitate was filtered, washed with water, and purified on a silica gel column using ethyl acetate:hexanes:glacial acetic acid 50:50:10 as eluent to give 45 mg (21%) of 3-[2-(6-chloro-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-



WO 00/08202

242

PCT/US99/17845 ,

yl]-propionic acid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ 13.2 (s, 1H, NH), 12.04 (s, 1 H, COOH), 10.92 (s, 1 H, NH), 7.72 (d, J = 8 Hz, 1 H, H-4), 7.68 (s, 1 H, H-vinyl), 7.15 (d, J = 2.4 Hz, 1 H), 7.01 (dd, J = 2.8 Hz, 1 H, H-6), 6.86 (d, J = 2 Hz, 1 H, H-7), 2.94 (t, J = 7.5 Hz, 2 H, CH<sub>2</sub>), 2.4 (t, J = 7.5 Hz, 2 H, CH<sub>2</sub>), 2.03 (s, 3 H, CH<sub>3</sub>).

MS m/z 329 (M-1).

**Example 220: 3-[2-(5-Bromo-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid**

10 A mixture of 5-formyl-4-(2-ethoxycarbonylethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester (281 mg), 5-bromo-2-oxindole (220 mg), and piperidine (2 drops) in ethanol (2 mL) was stirred at 90 °C for 2 hours. Potassium hydroxide (537 mg) was added and the temperature was  
15 increased to 95° C for 1 hour. The reaction mixture was cooled and concentrated. The residue was dissolved into water and acidified with 2 N hydrochloric acid to pH 2. The precipitate that formed was filtered, washed with water and dried in a vacuum oven overnight to give 411 mg (98%) of 4-  
20 (2-carboxyethyl)-3-methyl-5-(5-bromo-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid.

4-(2-carboxyethyl)-3-methyl-5-(5-bromo-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid (380 mg) suspended in ethylene glycol (5 mL) was held in a sealed tube  
25 in a pre-heated oil bath at 200°C for 2 hours. The reaction mixture was cooled to 90°C and potassium hydroxide (1 pellet) was added. It was then stirred at 90°C for 30 minutes. The reaction mixture was cooled, poured into water, and acidified with 2 N hydrochloric acid to pH 2. The precipitate was  
30 filtered, washed with water, and purified using silica gel column chromatography to give 168 mg (49% yield) of 3-[2-(5-bromo -2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid.

WO 00/08202

243

PCT/US99/17845

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ 13.32 (s, 1 H, NH), 12.0 (s, 1 H, COOH), 10.9 (s, 1 H, NH), 7.97 (d, J = 2 Hz, 1 H, h-4), 7.75 (s, 1 H, H-vinyl), 7.23 (dd, v = 2.8 Hz, 1 H, H-6), 7.16 (d, v = 2.6 Hz, 1 H), 6.8 (d, J = 8 Hz, 1 H, H-7), 2.97 (t, J = 7.7 Hz, 2 H, CH<sub>2</sub>), 2.41 (t, J = 7.7 Hz, 2 H, CH<sub>2</sub>), 2.04 (s, 3 H, CH<sub>3</sub>).

MS m/z 375/377.

**Example 221: 3-[2-(5-Methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid**

5-Methylisatin (15.0 g) and 60 mL of hydrazine hydrate were heated to 140-160°C for 4 hours. Thin layer chromatography (ethyl acetate:hexane 1:2, silica gel) showed no starting material remaining. The reaction mixture was cooled to room temperature, poured into 300 mL of ice water, and acidified to pH 2 with 6 N hydrochloric acid. After standing at room temperature for 2 days, the precipitate was collected by vacuum filtration, washed with water, and dried under vacuum to give 6.5 g (47%) of 5-methyl-2-oxindole.

<sup>1</sup>HNMR (360 MHz, DMSO-d<sub>6</sub>) δ 10.20 (s, br, 1 H, NH-1), 6.99 (s, 1 H, H-4), 6.94 (d, J = 8.11 Hz, 1 H, H-6), 6.68 (d, J = 8.11 Hz, 1 H, H-7), 3.39 (s, 2 H, CH<sub>2</sub>-3), and 2.22 (s, 3 H, CH<sub>3</sub>-5).

5-Formyl-4-(2-ethoxycarbonyl-ethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester (560 mg), 5-methyl-2-oxindole (300 mg), and piperidine (4 drops) in ethanol (4 mL) were stirred at 90°C for 2 hours. Potassium hydroxide (537 mg) was added to the mixture and it was then stirred at 95°C for an additional hour. The reaction mixture was cooled and concentrated. The residue was dissolved into water and acidified with 2 N hydrochloric acid to pH 2. The precipitate was filtered, washed with water, and dried in a vacuum oven overnight to give 496 mg of 4-(2-carboxyethyl)-3-methyl-5-(5-methyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid.

WO 00/08202

244

PCT/US99/17845

4-(2-carboxyethyl)-3-methyl-5-(5-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid (496 mg) suspended in ethylene glycol (2 mL) was held in a sealed tube in a pre-heated oil bath at 200°C for 2 hours.

5 The reaction mixture was cooled to 90°C and potassium hydroxide (157 mg) was added. It was then stirred at 90°C for 30 minutes. The reaction mixture was cooled, poured into water, and acidified with 2 N hydrochloric acid to pH 2. The precipitate was filtered, washed with water, and dried in a vacuum oven overnight to give 128 mg (29%) of 3-[2-(5-methyl-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ 13.31 (s, 1 H, NH), 12.07 (s, 1 H, COOH), 10.68 (s, 1 H, NH), 7.59 (s, 1 H), 7.51 (br s, 1 H, H-4), 7.09 (d, J = 2.7 Hz, 1H), 6.91 (br d, J = 8 Hz, 1 H, H-6), 6.73 (d, J = 8 Hz, 1 H, H-7), 2.93 (t, J = 7.5 Hz, 2 H, CH<sub>2</sub>), 2.41 (t, J = 7.5 Hz, 2 H, CH<sub>2</sub>), 2.3 (s, CH<sub>3</sub>-5), 2.04 (s, 3 H, CH<sub>3</sub>).

MS m/z 311 (M+1).

20 **Example 222:** 3-[2-(5-Methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid

A mixture of 5-formyl-4-(2-ethoxycarbonyl-ethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester (562 mg), 5-methoxy-2-oxindole (326 mg), and piperidine (2 drops) in ethanol (2 mL) were stirred at 90° C for 2 hours. Potassium hydroxide (537 mg) was added, the temperature was increased to 95°C and held there for 1 hour. The reaction mixture was cooled and concentrated. The residue was dissolved into water and acidified with 2 N hydrochloric acid to pH 2. The precipitate was filtered, washed with water, and dried in a vacuum oven overnight to give 240 mg (65%) of 4-(2-carboxyethyl)-3-methyl-5-(5-methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid.

WO 00/08202

245

PCT/US99/17845 ,

4-(2-carboxyethyl)-3-methyl-5-(5-methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-1H-pyrrole-2-carboxylic acid (240 mg) suspended in ethylene glycol (2 mL) was held in a sealed tube in a pre-heated oil bath at 200°C for 2 hours.

5 The reaction mixture was cooled to 90°C and potassium hydroxide (1 pellet) was added. It was then stirred at 90°C for 30 minutes. The reaction mixture was cooled, poured into water and acidified with 2 N hydrochloric acid to pH 2. The precipitate was filtered, washed with water, and dried in a vacuum oven overnight to give 30.5 mg (14%) of 3-[2-(5-methoxy-2-oxo-1,2-dihydroindol-3-ylidenemethyl)-4-methyl-1H-pyrrol-3-yl]-propionic acid.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 13.38 (s, 1 H, NH), 12.07 (s, 1 H, COOH), 10.59 (s, 1 H, NH), 7.63 (s, 1 H, H-vinyl), 7.1 (d, J = 2.1 Hz, 1 H), 6.75 (d, J = 8 Hz, 1 H, H-7), 6.69 (dd, J = 2.8 Hz, 1 H, H-6), 3.76 (s, 3 H, OCH<sub>3</sub>), 2.96 (t, J = 7.4 Hz, 2 H, CH<sub>2</sub>), 2.41 (t, J = 7.4 Hz, 2 H, CH<sub>2</sub>), 2.04 (s, 3 H, CH<sub>3</sub>).

MS m/z 327 (M+1).

**Example 223:** 3-{2-[6-(3-Methoxyphenyl)-2-oxo-1,2-dihydro-indol-3-ylidenemethyl]-4-methyl-1H-pyrrol-3-yl}-propionic acid

Tetrakis(triphenylphosphine)palladium (0.7 g) was added to a mixture of 5 g of 3-methoxyphenylboronic acid, 3.8 g of 5-bromo-2-fluoronitrobenzene, and 11 mL of 2 M sodium carbonate solution in 100 mL of toluene. The mixture was refluxed for 2 hours, diluted with water and extracted with ethyl acetate. The ethyl acetate was washed with saturated sodium bicarbonate and brine, dried, and concentrated to give an oily solid. The solid was purified on a silica gel column using a 1:6 mixture of ethyl acetate:hexane as eluent, to give 4.3 g (77%) of 4-fluoro-3'-methoxy-3-nitrobiphenyl.

Dimethyl malonate (9.7 mL) was added dropwise to 2.0 g of sodium hydride suspended in 50 mL of dimethylsulfoxide. The mixture was stirred at 100°C for 35 minutes and cooled to room

WO 00/08202

246

PCT/US99/17845 ,

temperature. 4-Fluoro-2'-methoxy-3-nitrobiphenyl (4.2 g) in 50 mL of dimethylsulfoxide was added and the mixture was stirred at 100°C for 1 hour. The reaction mixture was cooled, quenched with 300 mL of saturated ammonium chloride solution and extracted twice with ethyl acetate. The extracts were combined, washed with brine, dried over anhydrous sodium sulfate and concentrated to give crude dimethyl 3'-methoxy-3-nitrobiphenyl-4-malonate as a pale yellow solid.

Crude 3'-methoxy-3-nitro-biphenyl-4-malonate was stirred at 110°C in 45 mL of 6 N hydrochloric acid for 4 days and then cooled. The precipitate which formed was collected by filtration, washed with water and hexane, and dried to give 5.3 g of 3'-methoxy-2-nitrobiphenyl-4-acetic acid as a light tan solid.

3'-Methoxy-3-nitrobiphenyl-4-acetic acid (5.2 g) was dissolved in methanol and hydrogenated over 0.8 g of 10% palladium on carbon for 3 hours at room temperature. The catalyst was removed by filtration, washed with methanol, and the filtrates combined and concentrated to give a brown solid. The solid was purified on a silica gel column, using a 33:66:1 mixture of ethyl acetate:hexane:acetic acid as eluent, to give 3.0 g (75 % based on 4-fluoro-3'-methoxy-3-nitrobiphenyl) of 6-(3-methoxyphenyl)-2-oxindole as a pink solid.

<sup>1</sup>H NMR (360 MHz, DMSO-d<sub>6</sub>) δ 10.39 (s, br, 1 H, NH), 7.35 (t, J = 7.85 Hz, 1 H), 7.26 (d, J = 7.78 Hz, 1 H), 7.19 (dd, J = 1.22, 7.8 Hz, 1 H), 7.13-7.16 (m, 1 H), 7.09-7.1 (m, 1 H), 7.01 (d, J = 1.48 Hz, 1 H), 6.90-6.93 (m, 1 H), 3.8 (s, 3 H, OCH<sub>3</sub>), 3.49 (s, 2H, CH<sub>2</sub>).

MS m/z (relative intensity, %) 240.0 ([M+1]<sup>+</sup>, 100).

5-Formyl-4-(2-ethoxycarbonyl-ethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester (281 mg), 6-(3-methoxyphenyl)-2-oxindole (287 mg), and piperidine (2 drops) in

WO 00/08202

247

PCT/US99/17845 ,

ethanol (5 mL) were stirred at 90°C overnight. The precipitate was filtered and washed with ethanol. The yellow orange solid and potassium hydroxide (4 pellets) were stirred in ethanol (3 mL) at 90°C for 2.5 hours. The reaction mixture was cooled and concentrated. The residue was dissolved into water and acidified with 2 N hydrochloric acid to pH 2. The precipitate was filtered, washed with water, and dried in a vacuum oven overnight to give 413 mg of 4-(2-Carboxyethyl)-5-[6-(3-methoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-3-methyl-1H-pyrrole-2-carboxylic acid.

4-(2-Carboxyethyl)-5-[6-(3-methoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-3-methyl-1H-pyrrole-2-carboxylic acid (413 mg) suspended in ethylene glycol (5 mL) was held in sealed tube in a pre-heated oil bath at 200°C for 2 ½ hours. The reaction mixture was cooled to 90°C and potassium hydroxide (4 pellets) was added. It was then stirred at 100°C for 2 more hours. The reaction mixture was cooled, poured into water and acidified with 2 N hydrochloric acid to pH 2. The precipitate was filtered, washed with water and dried in a vacuum oven overnight. The crude solid was purified on a silica gel column using 33:66:1 ethyl acetate:hexanes:glacial acetic acid as eluent to give 75 mg (20%) of 3-{2-[6-(3-methoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-4-methyl-1H-pyrrol-3-yl}-propionic acid as an orange-red solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ 13.27 (s, 1 H, NH), 12.06 (s, 1 H, COOH), 10.86 (s, 1 H, NH), 7.76 (d, J = 8 Hz, 1 H, H-4), 7.68 (s, 1 H, H-vinyl), 7.36 (t, J = 8 Hz, 1 H), 7.29 (dd, J = 1.5, 8 Hz, 1 H), 6.92, 7.09, 7.13, 7.2 (m, 5 H, Ar-H), 3.82 (s, 3 H, OCH<sub>3</sub>), 2.96 (t, J = 7.4 Hz, 2 H, CH<sub>2</sub>), 2.42 (t, J = 7.4 Hz, 2 H, CH<sub>2</sub>), 1.97 (s, 3H, CH<sub>3</sub>).

WO 00/08202

248

PCT/US99/17845

**Example 224: 3-{2-[6-(3-Ethoxyphenyl)-2-oxo-1,2-dihydro-indol-3-ylidenemethyl]-4-methyl-1H-pyrrol-3-yl}-propionic acid**

A mixture of 5-formyl-4-(2-ethoxycarbonylethyl)-3-methyl-1H-pyrrole-2-carboxylic acid ethyl ester (281 mg), 6-  
5 (3-ethoxyphenyl)-2-oxindole (304 mg), and piperidine (2 drops) in ethanol (5 mL) were held at 90 °C overnight. The precipitate that formed was filtered, washed with ethanol. The precipitate, an orange solid, was stirred with potassium hydroxide (4 pellets) in ethanol (3 mL) at 90°C for 2.5  
10 hours. The reaction mixture was cooled and concentrated. The residue was dissolved into water and acidified with 2N hydrochloric acid to pH 2. The precipitate was filtered, washed with water, and dried in a vacuum oven overnight to give 370 mg of 4-(2-Carboxyethyl)-5-[6-(3-ethoxyphenyl)-2-  
15 oxo-1,2-dihydroindol-3-ylidenemethyl]-3-methyl-1H-pyrrole-2-carboxylic acid.

4-(2-Carboxyethyl)-5-[6-(3-ethoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-3-methyl-1H-pyrrole-2-carboxylic acid (350 mg) suspended in ethylene glycol (5 mL) was held in  
20 a sealed tube in a pre-heated oil bath at 200°C for 2.5 hours. The reaction mixture was cooled to 100°C and potassium hydroxide (4 pellets) was added. It was then stirred at 100°C for 2 hours. The reaction mixture was cooled, poured into water, and acidified with 2 N hydrochloric acid to pH 2. The  
25 precipitate was filtered, washed with water, and dried in a vacuum oven overnight. The crude solid was purified on a silica gel column using 33:66:1 ethyl acetate:hexanes:glacial acetic acid to give 140 mg (44%) of 3-{2-[6-(3-ethoxyphenyl)-2-oxo-1,2-dihydroindol-3-ylidenemethyl]-4-methyl-1H-pyrrol-3-  
30 yl}-propionic acid as a brown solid.

<sup>1</sup>HNMR (d<sub>6</sub>-DMSO) δ 13.28 (s, 1 H, NH), 12.04 (s, 1 H, COOH), 10.86 (s, 1 H, NH), 7.76 (d, J = 8 Hz, 1 H, H-4), 7.68 (s, 1 H, H-vinyl), 7.34 (t, J = 8 Hz, 1 H), 7.28 (dd, J = 2, 8

WO 00/08202

249

PCT/US99/17845 ,

Hz, 1 H, H-5), 7.08 (d,  $J = 2$  Hz, 1 H, H-7), 7.18, 7.13, 6.9 (m, 4 H, Ar-H), 4.1 (q,  $J = 7$  Hz, 2 H,  $\text{OCH}_2\text{CH}_3$ ), 2.96 (t,  $J = 7.5$  Hz, 2 H,  $\text{CH}_2$ ), 2.43 (t,  $J = 7.5$  Hz, 2 H,  $\text{CH}_2$ ), 2.05 (s, 3 H,  $\text{CH}_3$ ), 1.35 (t,  $J = 7$  Hz, 3 H,  $\text{OCH}_2\text{CH}_3$ ).

## 5 5. BIOLOGICAL EVALUATION

It will be appreciated that, in any given series of compounds, a spectrum of biological activity will be afforded. In its presently preferred embodiments, this invention relates to 3-methylidenyl-2-indolinones which demonstrate the ability to modulate RTK, CTK, and STK activity. The assays employed to select those compounds demonstrating of the desired activity are set forth below. Table 2 shows the results of testing of compounds of this invention using the described assays. The results shown are not to be construed as limiting the scope of this invention in any manner whatsoever.



WO 00/08202

250

PCT/US99/17845

Compound	HER-2 kinase IC <sub>50</sub> (μM)	BrdU/HER 2 driven IC <sub>50</sub> (μM)	BrdU/EGF driven IC <sub>50</sub> (μM)	SKOV3 growth (2%FBS) IC <sub>50</sub> (μM)	A431 growth (2%FBS) IC <sub>50</sub> (μM)	bio FLK1 (μM) % if inhibition at 100 (μM)	bio flkGST IC <sub>50</sub> (μM)	bio fgfr1 IC <sub>50</sub> (μM)	bio PDGF IC <sub>50</sub> (μM)	bio EGFr IC <sub>50</sub> (μM)	bio pyk2 IC <sub>50</sub> (μM)	cell EGF IC <sub>50</sub> (μM)	bio src IC <sub>50</sub> (μM)	HUVEC- VEGF IC <sub>50</sub> (μM)	HUVEC- aFGF IC <sub>50</sub> (μM)	PDGF- Induced BrdU Incorp. IC <sub>50</sub> (μM)	FGF- Induced BrdU Incorp. IC <sub>50</sub> (μM)
1	13.8	8.7	6.6	5.5	5.4	>50			>100	>100		45.7					
2	13.7	ND	6.5	4.8	4.5	>50			>100	>100		>100					
3	6.4	9.1	11	2.6	2.2												
4	5.8	4.4	26.6	7	5.1					>100		20.08					
5	6.5	10.5	7.2	5.4	4.3					>100		21.47					
6	15.3	4.7	12	19.4	2.9					>100		40.47					
7	>100	16.5	24.5	2.5	1.2					>100		40.37					
8	8.36	33.4	>50	0.86	0.95					>100		43.66					
9	>100	4.5	11	14.5	17.3					>100		34.59					
10	45.78	14.1	29	11.2	8.5					>100		56.07					
11	57.3	9.3	23.3	15	12.5		24.12		>100	>100		72.94					
12	>100	31.9	30.6	10.1	7.9					>100		71.92					
13	>100	9	26.6	21.3	9				>100	>100		30.78					
14	22.7	ND	ND	ND	ND					>100		>100					
15	19	4.4	7.8	7	4.8				28.52	>100		46.52					
16	9.9	4.2	6.4	5.9	4.1				>100	>100		44.48					
17	7.6	5.5	7.4	6.1	4.3					>100		67.98					
18	8.4	>50	>50	1	0.6				86.93	>100		>100					
19	9.2	3.3	9.3	5.9	4.3					>100		34.80					
20	38.6	15.7	30	35.5	29.5					>100		76.08					
21	11.8	6.6	8	13	15.3					>100		18.45					
22	12.4	ND	ND	ND	ND					>100		>100					
23	33.2	27.1	32.9	17.8	14.5					>100		37.71					
24	13.6	7.1	7.5	6.5	4.5				>100	>100		36.47					
26	15.3	>50	>50	5.4	3					>100		93.99					
27	15.8	23.4	26.8	5.1	6.4				>100	>100		30.58					
28	16.92									>100		>100					

TABLE 2

WO 00/08202

251

PCT/US99/17845

Compound	HER-2 kinase IC <sub>50</sub> (μM)	BrdU HER 2 driven IC <sub>50</sub> (μM)	BrdU EGF driven IC <sub>50</sub> (μM)	SKOV3 growth (2%FBS) IC <sub>50</sub> (μM)	A431 growth (2%FBS) IC <sub>50</sub> (μM)	bio FLK1 (μM) % if inhibition at 100 (μM)	bio fIKGST IC <sub>50</sub> (μM)	bio fgR1 IC <sub>50</sub> (μM)	bio PDGF IC <sub>50</sub> (μM)	bio EGFr IC <sub>50</sub> (μM)	bio pyk2 IC <sub>50</sub> (μM)	cell EGF IC <sub>50</sub> (μM)	HUVEC- VEGF IC <sub>50</sub> (μM)	HUVEC aFGF IC <sub>50</sub> (μM)	PDGF- Induced BrdU incorp. IC <sub>50</sub> (μM)	FGF- Induced BrdU incorp. IC <sub>50</sub> (μM)
29	17.1	8.1	15.2	7.9	5.6				>100	>100		>100				
30	28.02	8.1	30.3	16	9.9				>100	>100		63.44				
31	12.92	28.7	38.6	0.4	0.3				>100	>100		29.57				
32	90.18	ND	ND	ND	ND				>100	>100		>100				
33	20.2	ND	ND	10.3	6.7				>100	>100		53.19				
34	>100	3.5	11	7.6	4.4				56.68	>100		38.19				
35	20.7	13.8	28.1	16.2	13.4				>100	>100		>100				
36	20.7	21.4	31	>25	>25				>100	>100		66.81				
37	22.5	ND	ND	ND	ND				>100	>100		>100				
38	32.3	ND	ND	ND	ND				>100	>100		>100				
39	33.7	5.4	16.2	9.6	9.2				>100	>100		>100				
40	36.6	22.6	28.6	22.5	14.2				>100	>100		78.95				
41	39.8	ND	ND	ND	ND				>100	>100		>100				
42	47.5	ND	ND	ND	ND				43.08	>100		>100				
43	52.9	ND	ND	ND	ND				10.99	>100		>100				
44	53.6	ND	ND	ND	ND				>100	>100		78.93				
45	56.9	>50	>50	ND	3.2				23.98	>100		>100				
46	59.9	25.3	45.7	7.7	5.5				>100	>100		>100				
47	60.6	>50	>50	46.9	40.4					36.82		>100				
48	65	ND	ND	ND	ND				>100	>100		>100				
49	65.5	9.1	24.1	20.6	20.7				>100	>100		>100				
50	67.6	31.1	37.9	ND	ND				>100	>100		>100				
51	71.8	ND	ND	14.5	ND				>100	>100		>100				
52	72.8	ND	ND	ND	ND				>100	>100		>100				
53	80.7	ND	ND	ND	ND				>100	>100		85.04				
54	85.6	ND	ND	ND	ND				>100	>100		79.32				
55	89.3	ND	ND	ND	ND				>100	>100		78.36				
56	96.2	ND	ND	>20	ND				10.1	>100		>100				
57	>100	9.6	12.3	4	5.8				>100	>100		>100				
58	>100	13.5	27.5	22	10.5				>100	>100		>100				
59	>100	14.4	44.5	17.2	6				>100	>100		>100				

WO 00/08202

252

PCT/US99/17845

Compound	HER-2 kinase IC <sub>50</sub> (μM)	BrdU HER 2 driven IC <sub>50</sub> (μM)	BrdU EGF driven IC <sub>50</sub> (μM)	SKOV3 growth (2%FBS) IC <sub>50</sub> (μM)	A431 growth (2%FBS) IC <sub>50</sub> (μM)	bio FLK1 (μM) % if inhibition at 100 (μM)	bio flkGST IC <sub>50</sub> (μM)	bio fgR1 IC <sub>50</sub> (μM)	bio PDGF IC <sub>50</sub> (μM)	bio EGF IC <sub>50</sub> (μM)	bio pyk2 IC <sub>50</sub> (μM)	cell EGF IC <sub>50</sub> (μM)	bio src IC <sub>50</sub> (μM)	HUVEC- VEGF IC <sub>50</sub> (μM)	HUVEC aFGF IC <sub>50</sub> (μM)	PDGF- Induced BrdU Incorp. IC <sub>50</sub> (μM)	FGF- Induced BrdU Incorp. IC <sub>50</sub> (μM)
60	>100	40	>50	4.8	5.9				24.49	>100		>100					
61	>100	>50	>50	>25	>25				>100	>100		>100					
62	>100	28.8	>50	6.1	6				>100	>100		>100					
63	>100	20.4	30.5	>25	11				>100	>100		>100					
64	>100								>100	>100		>100					
65	>100									>100		>100					
66	>100									>100		>100					
67	>100									>100		>100					
68	>100									>100		>100					
69	>100									>100		>100					
70	>100									>100		>100					
71	>100								22.6			>100					
72	>100								29.6			>100					
73	>50								11.8			>100					
74	>100								7.9			>50					
107	>100								13.2			>100					
108	>100								<0.78			>100					
109	>100								2.97			>100					
110	>100								2.79			>100					
111	>100								3.15			>100					
112									0.01			>100					
114	61.73								0.02			>100					
116	6.07								0.05			>100					
117									0.02			>100					
118									>100								
119									>100								
120									9.97								
121									26.66								
122									2.13								
123									15.21								
124									33.89								

WO 00/08202

253

PCT/US99/17845

Compound	HER-2 kinase IC <sub>50</sub> (μM)	BrdU HER 2 driven IC <sub>50</sub> (μM)	SKOV3 growth (2%FBS) IC <sub>50</sub> (μM)	A431 growth (2%FBS) IC <sub>50</sub> (μM)	bio FLK1 (μM) % if inhibition at 100 (μM)	bio fIKGST IC <sub>50</sub> (μM)	bio fgfr1 IC <sub>50</sub> (μM)	bio PDGF IC <sub>50</sub> (μM)	bio EGFr IC <sub>50</sub> (μM)	bio pyk2 IC <sub>50</sub> (μM)	cell EGF IC <sub>50</sub> (μM)	bio src IC <sub>50</sub> (μM)	HUVEC- VEGF IC <sub>50</sub> (μM)	HUVEC aFGF IC <sub>50</sub> (μM)	PDGF- Induced BrdU incorp. IC <sub>50</sub> (μM)	FGF- Induced BrdU incorp. IC <sub>50</sub> (μM)
125								5.73								
126						14.8		63.28								
127								0.86								
128								17.11								
129								31.69								
130						12.88		2.32								
131								23.18		2.18						
132								0.65		0.22						
133						8.18		2.69								
134								4.88								
135								0.5	0.19							
136						0.56		26.95		2.77						
137						7.54		6.17	>100	>10						
138						17.65		26.24	>100	2.18						
139						7.56	12.57	>100	>100							
140						6.65	12.83	>100	>100							
141						11.53	8.38	85.11	>100							
142						21.14	5.9	>100	>100							
143						26.92	16.56	>100	>100							
144						9.03	2.77	>100	>100							
145						11.79	15.57	>100	>100							
146						15.6	>20	>100	>100							
147						15.28	>20	>100	>100							
148						23.94	>20	>100	>100							
149						2.66	0.77	5.26		5.02						
150						0.2		0.34		0.01						
151						0.71		0.65								
152	>100						>20		>100							
153	>100						>20	>100	>100							
154	>100						>20	>100	>100			>100				
155	>100								>100							
156	>100				4.2		3.60	8.32	8.2	0.22		0.15				
157	>100						>20	>100	30.8			>100				
158						0.26	>20	18.03	>100	12.35	>100					
159					1.53	2.87	>20	74.08	>100	9.6	>100					
160						0.12	5.27	2.72	>100	0.68	>100	9				

WO 00/08202

254

PCT/US99/17845

Compound	HER-2 kinase IC <sub>50</sub> (μM)	BrdU HER 2 driven IC <sub>50</sub> (μM)	SKOV3 growth (2%FBS) IC <sub>50</sub> (μM)	A431 growth (2%FBS) IC <sub>50</sub> (μM)	bio FLK1 (μM) % if inhibition at 100 (μM)	bio tkGST IC <sub>50</sub> (μM)	bio fgR1 IC <sub>50</sub> (μM)	bio PDGF IC <sub>50</sub> (μM)	bio EGFr IC <sub>50</sub> (μM)	bio pyk2 IC <sub>50</sub> (μM)	cell EGF IC <sub>50</sub> (μM)	bio src IC <sub>50</sub> (μM)	HUVEC- VEGF IC <sub>50</sub> (μM)	HUVEC aFGF IC <sub>50</sub> (μM)	PDGF- Induced BrdU Incorp. IC <sub>50</sub> (μM)	FGF- Induced BrdU Incorp. IC <sub>50</sub> (μM)
161					2.1	<0.008	>20	25.06	>100	3.4	>100					
162							>20	8.02	>100		>100					
163							>20	>100	>100		>100					
164	>100						>20	>100	>100		>100					
165	>100						>20	48.71	>100		>100					
166	>100						>20	>100	>100		>100					
167							>20	>100	>100		>100					
168							>20	>100	>100							
169					15.8		>20	>100	17.8							
170					10.7		>20	>100	>100							
171					20.35		>20	68.56	>100		>100					
172						10.47	0.28	89.41	>100	14.75	>100					
173						11.05	0.29	69.14	22.04	9.8	>100					
174						1.85	0.41	2.5	>100	2.57	>100					
175						1.51	0.73	55.38	>100	5.53	>100					
176						2.08	0.75	79.22	>100	66.08	>100					
177						1.34	1.6	>100	6.13	>100	>100					
178						>20	3.23	>100	>100	57.09	>100					
179						1.84	4.52	>100	>100	>100	>100					
180						1.11	5.42	1.11	>100	<0.78	>100					
181						12.53	5.74	>100	>100	>100	>100					
182						18.68	5.94	5.4	>100	37.09	>100					
183						1.55	7.68	>100	>100	>100	>100					
184						>20	8.6	>100	>100	>100	>100					
185						3.39	9.43	>100	>100	>100	>100					
186						6.12	10.38	>100	>100	>100	>100					
187						>20	13.82	>100	>100	75.8	>100					
188						>20	17.47	>100	>100	68.41	>100					
189						>20	>20	>100	>100	>100	>100					
190					0.06	0.087	5.85	2.08	>100	<0.78		0.11	0.0019	0.0159	2.9	18.2
191					0.02	0.04	0.04	0.05	>100	68.5	>100	11.96	0.00004	0.25	1.1	23.6
192					0.03	0.03	1.81	0.004	>100	4.87	>100	10.49	0.0003	0.076	1.4	15.7

WO 00/08202

255

PCT/US99/17845

Compound	HER-2 kinase IC <sub>50</sub> (μM)	BrdU HER 2 driven IC <sub>50</sub> (μM)	BrdU EGF driven IC <sub>50</sub> (μM)	SKOV3 growth (2%FBS) IC <sub>50</sub> (μM)	A431 growth (2%FBS) IC <sub>50</sub> (μM)	bio FLK1 IC <sub>50</sub> (μM) % ff inhibition at 100 (μM)	bio fKST IC <sub>50</sub> (μM)	bio fKST IC <sub>50</sub> (μM)	bio EGFR IC <sub>50</sub> (μM)	bio EGFR IC <sub>50</sub> (μM)	bio pyk2 IC <sub>50</sub> (μM)	cell EGF IC <sub>50</sub> (μM)	bio src IC <sub>50</sub> (μM)	HUVEC VEGF IC <sub>50</sub> (μM)	HUVEC aEGF IC <sub>50</sub> (μM)	PDGF- Induced BrdU Incorp. IC <sub>50</sub> (μM)	FGF- Induced BrdU Incorp. IC <sub>50</sub> (μM)
193						0.02	3.60	0.02	44.55	90.72	17.32		48.59	0.0094	0.0025	13.5	26.7
194						0.001	6.67	0.04	81.35		4.09		20.58	0.001	0.0044	14.2	30.1
195						0.03	4.35	0.02	2.04	36.48	10.70	>100	0.26			3.7	30.2
196						0.89	0.41	0.05	33.68	>100	2.37	>100	11.25	0.02	1.6	3	42.1
197						2.10	0.61		>100	>100	11.98	>100		<0.003	202	4.9	>50
198						7.44	5.84		0.04	>100	1.5	>100	2.34	0.29	1.7	4.2	>50
199						0.13	0.17		0.89	>100	2.83	>100		<0.03	0.1	3.5	>50
200						0.97	1.29		11.80	>100	5.62	>100		0.22	9.2	8.6	>50
201						0.16	0.57		1.73	>100	6.61	>100		<0.03	0.16	3.9	>50
202						0.63	2.54		9.58	>100		>100	>50	<0.03	2.1	5.2	>50
203						0.14		2.27	0.15	>12.5		>12.5					
204						0.36	0.07	0.59	0.68	7.09		>100	4.44				
205						0.52	0.44	2.02	5.43	16.21		>100	1.34				
206						6.08	0.06	1.18	0.04	>100		>100	0.74				
207						2.72	1.50	7.70	0.87	>100		>100	7.52				
208						<0.78	0.38		23.45	>100	0.13	>100	1.84				
209						50.64		30.24	>100	>100		>100					
214						0.02	0.08	0.07	0.50	>100		>100	>100				
215			>100			2.91			0.22	>100	6.90	>100	5.59	0.07	1.1	20	25
216			>50			1.47			0.21	>100	23.86	>100	4.75	<0.07	1.4	N/A *	N/A *
217			>50			0.20	0.48		0.21	>100	2.89	>100	3.68	0.62	4.8	8.7	>50
218			43.5			0.19		0.10	2.11	>100	32.27	>100		<0.07	0.35	24.8	>50
219			>50			0.35	0.11		0.47	98.06	4.15	>100	0.08	0.016	0.26	7.9	>50
220			>50			<0.78	0.18		0.62	>100	1.89	>100	6.48	0.04	0.49	8.5	>50
222			>50			28.25	0.57	1.22	3.75	>100		>100	5.77			>50	35
223						1.0	0.67	4.24	0.781	13.29		>100	1.06				
224									2.38	>100			1.16				

WO 00/08202

256

PCT/US99/17845 ,

**A. Assay Procedures.**

The following in vitro assays may be used to determine the level of activity and effect of the different compounds of the present invention on one or more of the PKs. Similar assays can be designed along the same lines for any PK using techniques well known in the art.

The cellular/catalytic assays described herein are performed in an ELISA format. The general procedure is as follows: a compound is introduced to cells expressing the test kinase, either naturally or recombinantly, for a selected period of time after which, if the test kinase is a receptor, a ligand known to activate the receptor is added.

The cells are lysed and the lysate is transferred to the wells of an ELISA plate previously coated with a specific antibody recognizing the substrate of the enzymatic phosphorylation reaction. Non-substrate components of the cell lysate are washed away and the amount of phosphorylation on the substrate is detected with an

antibody specifically recognizing phosphotyrosine compared with control cells that were not contacted with a test compound. The cellular/biologic assays described herein measure the amount of DNA made in response to activation of a test kinase, which is a general measure of a proliferative response. The general procedure for this assay is as follows: a compound is introduced to cells expressing the

test kinase, either naturally or recombinantly, for a selected period of time after which, if the test kinase is a receptor, a ligand known to activate the receptor is added.

After incubation at least overnight, a DNA labeling reagent such as Bromodeoxyuridine (BrdU) or 3H-thymidine is added. The amount of labeled DNA is detected with either an anti-BrdU antibody or by measuring radioactivity and is compared to control cells not contacted with a test compound.

WO 00/08202

257

PCT/US99/17845 ,

**Cellular/Catalytic Assays**

Enzyme linked immunosorbent assays (ELISA) may be used to detect and measure the presence of PK activity. The ELISA may be conducted according to known protocols which are described  
5 in, for example, Voller, et al., 1980, "Enzyme-Linked Immunosorbent Assay," In: Manual of Clinical Immunology, 2d ed., edited by Rose and Friedman, pp 359-371 Am. Soc. Of Microbiology, Washington, D.C.

The disclosed protocol may be adapted for determining  
10 activity with respect to a specific PK. That is, the preferred protocols for conducting the ELISA experiments for specific PKs is provided below. However, adaptation of these protocols for determining a compound's activity for other members of the RTK family, as well as for CTKs and STKs, is  
15 well within the scope of knowledge of those skilled in the art.

**FLK-1 Assay**

An ELISA assay is conducted to measure the kinase activity of the FLK-1 receptor and more specifically, the  
20 inhibition or activation of TK activity on the FLK-1 receptor.

Specifically, the following assay can be conducted to measure kinase activity of the FLK-1 receptor in cells genetically engineered to express Flk-1.

**Materials and Reagents.**

- 25 a. Corning 96-well ELISA plates (Corning Catalog No. 25805-96),
- b. Cappel goat anti-rabbit IgG (catalog no. 55641),
- c. PBS (Gibco Catalog No. 450-1300EB),
- d. TBSW Buffer (50 mM Tris (pH 7.2), 150 mM NaCl and  
30 0.1% Tween-20),
- e. Ethanolamine stock (10% ethanolamine (pH 7.0), stored at 4°C),
- f. HNTG buffer (20mM HEPES buffer (pH 7.5), 150mM NaCl, 0.2% Triton X-100, and 10% glycerol),



WO 00/08202

258

PCT/US99/17845 ,

- g. EDTA (0.5 M (pH 7.0) as a 100X stock),
- h. Sodium orthovanadate (0.5 M as a 100X stock),
- i. Sodium pyrophosphate (0.2 M as a 100X stock),
- j. NUNC 96 well V bottom polypropylene plates (Applied  
5 Scientific Catalog No. AS-72092),
- k. NIH3T3 C7#3 Cells (FLK-1 expressing cells),
- l. DMEM with 1X high glucose L-Glutamine (catalog No.  
11965-050),
- m. FBS, Gibco (catalog no. 16000-028),
- 10 n. L-glutamine, Gibco (catalog no. 25030-016),
- o. VEGF, PeproTech, Inc. (catalog no. 100-20) (kept as 1  
µg/100 µl stock in Milli-Q dH<sub>2</sub>O and stored at -20° C,
- p. Affinity purified anti-FLK-1 antiserum,
- q. UB40 monoclonal antibody specific for  
15 phosphotyrosine (see, Fendley, et al., 1990, Cancer Research  
50:1550-1558),
- r. EIA grade Goat anti-mouse IgG-POD (BioRad catalog  
no. 172-1011),
- s. 2,2-azino-bis(3-ethylbenz-thiazoline-6-sulfonic acid  
20 (ABTS) solution (100mM citric acid (anhydrous), 250 mM Na<sub>2</sub>HPO<sub>4</sub>  
(pH 4.0), 0.5 mg/ml ABTS (Sigma catalog no. A-1888)), solution  
should be stored in dark at 4° C until ready for use,
- t. H<sub>2</sub>O<sub>2</sub> (30% solution) (Fisher catalog no. H325),
- u. ABTS/ H<sub>2</sub>O<sub>2</sub> (15ml ABTS solution, 2 µl H<sub>2</sub>O<sub>2</sub>) prepared 5  
25 minutes before use and left at room temperature,
- v. 0.2 M HCl stock in H<sub>2</sub>O,
- w. dimethylsulfoxide (100%) (Sigma Catalog No. D-8418),  
and
- y. Trypsin-EDTA (Gibco BRL Catalog No. 25200-049).
- 30 **Protocol.**
- 1. Coat Corning 96-well ELISA plates with 1.0 µg per  
well Cappel Anti-rabbit IgG antibody in 0.1M Na<sub>2</sub>CO<sub>3</sub> pH 9.6.  
Bring final volume to 150 µl per well. Coat plates overnight

WO 00/08202

259

PCT/US99/17845 ,

at 4°C. Plates can be kept up to two weeks when stored at 4°C.

2. Grow cells in Growth media (DMEM, supplemented with 2.0 mM L-Glutamine, 10% FBS) in suitable culture dishes until  
5 confluent at 37°C, 5% CO<sub>2</sub>.

3. Harvest cells by trypsinization and seed in Corning 25850 polystyrene 96-well round bottom cell plates, 25,000 cells/well in 200 µl of growth media.

4. Grow cells at least one day at 37°C, 5% CO<sub>2</sub>.

10 5. Wash cells with D-PBS 1X.

6. Add 200 µl/well of starvation media (DMEM, 2.0mM l-Glutamine, 0.1% FBS). Incubate overnight at 37°C, 5% CO<sub>2</sub>.

7. Dilute Compounds 1:20 in polypropylene 96 well plates using starvation media. Dilute dimethylsulfoxide 1:20  
15 for use in control wells.

8. Remove starvation media from 96 well cell culture plates and add 162 µl of fresh starvation media to each well.

9. Add 18 µl of 1:20 diluted compound dilution (from step 7) to each well plus the 1:20 dimethylsulfoxide dilution  
20 to the control wells (+/- VEGF), for a final dilution of 1:200 after cell stimulation. Final dimethylsulfoxide is 0.5%. Incubate the plate at 37°C, 5% CO<sub>2</sub> for two hours.

10. Remove unbound antibody from ELISA plates by inverting plate to remove liquid. Wash 3 times with TBSW +  
25 0.5% ethanolamine, pH 7.0. Pat the plate on a paper towel to remove excess liquid and bubbles.

11. Block plates with TBSW + 0.5% ethanolamine, pH 7.0, 150 µl per well. Incubate plate thirty minutes while shaking on a microtiter plate shaker.

30 12. Wash plate 3 times as described in step 10.

13. Add 0.5 µg/well affinity purified anti-FLU-1 polyclonal rabbit antiserum. Bring final volume to 150

WO 00/08202

260

PCT/US99/17845 ,

µl/well with TBSW + 0.5% ethanolamine pH 7.0. Incubate plate for thirty minutes while shaking.

14. Add 180 µl starvation medium to the cells and stimulate cells with 20 µl/well 10.0 mM sodium orthovanadate and 500 ng/ml VEGF (resulting in a final concentration of 1.0 mM sodium orthovanadate and 50 ng/ml VEGF per well) for eight minutes at 37°C, 5% CO<sub>2</sub>. Negative control wells receive only starvation medium.

15. After eight minutes, media should be removed from the cells and washed one time with 200 µl/well PBS.

16. Lyse cells in 150 µl/well HNTG while shaking at room temperature for five minutes. HNTG formulation includes sodium ortho vanadate, sodium pyrophosphate and EDTA.

17. Wash ELISA plate three times as described in step 10.

18. Transfer cell lysates from the cell plate to ELISA plate and incubate while shaking for two hours. To transfer cell lysate pipette up and down while scrapping the wells.

19. Wash plate three times as described in step 10.

20. Incubate ELISA plate with 0.02 µg/well UB40 in TBSW + 0.5% ethanolamine. Bring final volume to 150 µl/well. Incubate while shaking for 30 minutes.

21. Wash plate three times as described in step 10.

22. Incubate ELISA plate with 1:10,000 diluted EIA grade goat anti-mouse IgG conjugated horseradish peroxidase in TBSW plus 0.5% ethanolamine, pH 7.0. Bring final volume to 150 µl/well. Incubate while shaking for thirty minutes.

23. Wash plate as described in step 10.

24. Add 100 µl of ABTS/H<sub>2</sub>O<sub>2</sub> solution to well. Incubate ten minutes while shaking.

25. Add 100 µl of 0.2 M HCl for 0.1 M HCl final concentration to stop the color development reaction. Shake 1

WO 00/08202

261

PCT/US99/17845 ,

minute at room temperature. Remove bubbles with slow stream of air and read the ELISA plate in an ELISA plate reader at 410 nm.

**EGF Receptor-HER2 Chimeric Receptor Assay In Whole Cells.**

5 HER2 kinase activity in whole EGFR-NIH3T3 cells are measured as described below:

**Materials and Reagents.**

a. EGF: stock concentration: 16.5 ILM, EGF 201, TOYOBO, Co., Ltd. Japan.

10 b. 05-101 (UBI) (a monoclonal antibody recognizing an EGFR extracellular domain).

c. Anti-phosphotyrosine antibody (anti-Ptyr) (polyclonal) (see, Fendley, et al., supra).

d. Detection antibody: Goat anti-rabbit IgG  
15 horseradish peroxidase conjugate, TAGO, Inc., Burlingame, CA.

e. TBST buffer:

Tris-HCl, pH 7.2	50 mM
NaCl	150 mM
Triton X-100	0.1

20

f. HNTG 5X stock:

HEPES	0.1 M
NaCl	0.75 M
Glycerol	50%
25 Triton X-100	1.0%

g. ABTS stock:

Citric Acid	100 mM
Na <sub>2</sub> HPO <sub>4</sub>	250 mM
HCl, conc.	0.5 mM
30 ABTS*	0.5mg/ml

\* (2,2'-azinobis(3-ethylbenzthiazolinesulfonic acid)). Keep solution in dark at 4°C until use.

h. Stock reagents of:

WO 00/08202

262

PCT/US99/17845

EDTA 100 mM pH 7.0

 $\text{Na}_3\text{VO}_4$  0.5 M $\text{Na}_4(\text{P}_2\text{O}_7)$  0.2 M**Procedure.****5 Pre-coat ELISA Plate**

1. Coat ELISA plates (Corning, 96 well, Cat. #25805-96) with 05-101 antibody at 0.5  $\mu\text{g}$  per well in PBS, 100  $\mu\text{l}$  final volume/well, and store overnight at 4°C. Coated plates are good for up to 10 days when stored at 4°C.

10 2. On day of use, remove coating buffer and replace with 100  $\mu\text{l}$  blocking buffer (5% Carnation Instant Non-Fat Dry Milk in PBS). Incubate the plate, shaking, at room temperature (about 23°C to 25°C) for 30 minutes. Just prior to use, remove blocking buffer and wash plate 4 times with TBST  
15 buffer.

**Seeding Cells**

1. An NIH3T3 cell line overexpressing a chimeric receptor containing the EGFR extracellular domain and intracellular HER2 kinase domain can be used for this assay.

20 2. Choose dishes having 80-90% confluence for the experiment. Trypsinize cells and stop reaction by adding 10% fetal bovine serum. Suspend cells in DMEM medium (10% CS DMEM medium) and centrifuge once at 1500 rpm, at room temperature for 5 minutes.

25 3. Resuspend cells in seeding medium (DMEM, 0.5% bovine serum), and count the cells using trypan blue. Viability above 90% is acceptable. Seed cells in DMEM medium (0.5% bovine serum) at a density of 10,000 cells per well, 100  $\mu\text{l}$  per well, in a 96 well microtiter plate. Incubate seeded  
30 cells in 5%  $\text{CO}_2$  at 37°C for about 40 hours.

**Assay Procedures**

1. Check seeded cells for contamination using an inverted microscope. Dilute drug stock (10 mg/ml in DMSO) 1:10

WO 00/08202

263

PCT/US99/17845 ,

in DMEM medium, then transfer 5  $\mu$ l to a TBST well for a final drug dilution of 1:200 and a final DMSO concentration of 1%. Control wells receive DMSO alone. Incubate in 5% CO<sub>2</sub> at 37°C for two hours.

- 5                   2.     Prepare EGF ligand: dilute stock EGF in DMEM so that upon transfer of 10  $\mu$ l dilute EGF (1:12 dilution), 100 nM final concentration is attained.

                  3.     Prepare fresh HNTG\* sufficient for 100  $\mu$ l per well, and place on ice.

10                  HNTG\* (10 ml):

                  HNTG stock                               2.0 ml

                  milli-Q H<sub>2</sub>O                               7.3 ml

                  EDTA, 100 mM, pH 7.0                   0.5 ml

                  Na<sub>3</sub>VO<sub>4</sub> (0.5 M)                           0.1 ml

15                  Na<sub>4</sub>(P<sub>2</sub>O<sub>7</sub>) (0.2 M)                       0.1 ml

                  4.     After 120 minutes incubation with drug, add prepared SGF ligand to cells, 10  $\mu$ l per well, to a final concentration of 100 nM. Control wells receive DMEM alone. Incubate with shaking, at room temperature, for 5 minutes.

- 20                  5.     Remove drug, EGF, and DMEM. Wash cells twice with PBS. Transfer HNTG\* to cells, 100  $\mu$ l per well. Place on ice for 5 minutes. Meanwhile, remove blocking buffer from other ELISA plate and wash with TBST as described above.

                  6.     With a pipette tip securely fitted to a  
25 micropipettor, scrape cells from plate and homogenize cell material by repeatedly aspirating and dispensing the HNTG\* lysis buffer. Transfer lysate to a coated, blocked, and washed ELISA plate. Incubate shaking at room temperature for one hour.

- 30                  7.     Remove lysate and wash 4 times with TBST. Transfer freshly diluted anti-Ptyr antibody to ELISA plate at 100  $\mu$ l per well. Incubate shaking at room temperature for 30 minutes in the presence of the anti-Ptyr antiserum (1:3000

WO 00/08202

264

PCT/US99/17845 ,

dilution in TBST).

8. Remove the anti-Ptyr antibody and wash 4 times with TBST. Transfer the freshly diluted TAGO anti-rabbit IgG antibody to the ELISA plate at 100  $\mu$ l per well. Incubate shaking at room temperature for 30 minutes (anti-rabbit IgG antibody: 1:3000 dilution in TBST).

9. Remove TAGO detection antibody and wash 4 times with TBST. Transfer freshly prepared ABTS/H<sub>2</sub>O<sub>2</sub> solution to ELISA plate, 100  $\mu$ l per well. Incubate shaking at room temperature for 20 minutes. (ABTS/H<sub>2</sub>O<sub>2</sub> solution: 1.0  $\mu$ l 30% H<sub>2</sub>O<sub>2</sub> in 10 ml ABTS stock).

10. Stop reaction by adding 50  $\mu$ l 5N H<sub>2</sub>SO<sub>4</sub> (optional), and determine O.D. at 410 nm.

11. The maximal phosphotyrosine signal is determined by subtracting the value of the negative controls from the positive controls. The percent inhibition of phosphotyrosine content for extract-containing wells is then calculated, after subtraction of the negative controls.

#### PDGF-R Assay

All cell culture media, glutamine, and fetal bovine serum can be purchased from Gibco Life Technologies (Grand Island, NY) unless otherwise specified. All cells are grown in a humid atmosphere of 90-95% air and 5-10% CO<sub>2</sub> at 37°C. All cell lines are routinely subcultured twice a week and are negative for mycoplasma as determined by the Mycotect method (Gibco).

For ELISA assays, cells (U1242, obtained from Joseph Schlessinger, NYU) are grown to 80-90% confluency in growth medium (MEM with 10% FBS, NEAA, 1 mM NaPyr and 2 mM GLN) and seeded in 96-well tissue culture plates in 0.5% serum at 25,000 to 30,000 cells per well. After overnight incubation in 0.5% serum-containing medium, cells are changed to serum-free medium and treated with test compound for 2 hr in a 5%

WO 00/08202

265

PCT/US99/17845 ,

CO<sub>2</sub>, 37°C incubator. Cells are then stimulated with ligand for 5-10 minute followed by lysis with HNTG (20 mM Hepes, 150 mM NaCl, 10% glycerol, 5 mM EDTA, 5 mM Na<sub>3</sub>VO<sub>4</sub>, 0.2% Triton X-100, and 2 mM NaPyr). Cell lysates (0.5 mg/well in 5 PBS) are transferred to ELISA plates previously coated with receptor-specific antibody and which had been blocked with 5% milk in TBST (50 mM Tris-HCl pH 7.2, 150 mM NaCl and 0.1% Triton X-100) at room temperature for 30 min. Lysates are incubated with shaking for 1 hour at room temperature. The 10 plates are washed with TBST four times and then incubated with polyclonal anti-phosphotyrosine antibody at room temperature for 30 minutes. Excess anti-phosphotyrosine antibody is removed by rinsing the plate with TBST four times. Goat anti-rabbit IgG antibody is added to the ELISA 15 plate for 30 min at room temperature followed by rinsing with TBST four more times. ABTS (100 mM citric acid, 250 mM Na<sub>2</sub>HPO<sub>4</sub> and 0.5 mg/mL 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid)) plus H<sub>2</sub>O<sub>2</sub> (1.2 mL 30% H<sub>2</sub>O<sub>2</sub> to 10 ml ABTS) is added to the ELISA plates to start color development. 20 Absorbance at 410 nm with a reference wavelength of 630 nm is recorded about 15 to 30 min after ABTS addition.

#### IGF-1 RECEPTOR Assay

The following protocol may be used to measure phosphotyrosine level on IGF-1 receptor, which indicates IGF-1 25 receptor tyrosine kinase activity.

#### **Materials and Reagents.**

- a. The cell line used in this assay is 3T3/IGF-1R, a cell line genetically engineered to overexpresses IGF-1 receptor.
- 30 b. NIH3T3/IGF-1R is grown in an incubator with 5% CO<sub>2</sub> at 37°C. The growth media is DMEM + 10% FBS (heat inactivated)+ 2mM L-glutamine.
- c. Affinity purified anti-IGF-1R antibody 17-69.
- d. D-PBS:



WO 00/08202

266

PCT/US99/17845

- $\text{KH}_2\text{PO}_4$  0.20 g/l  
 $\text{KH}_2\text{PO}_4$  2.16 g/l  
KCl 0.20 g/l  
NaCl 8.00 g/l (pH 7.2)
- 5 e. Blocking Buffer: TBST plus 5% Milk (Carnation Instant Non-Fat Dry Milk).
- f. TBST buffer:
- Tris-HCl 50 mM  
NaCl 150mM (pH 7.2/HCl 10N)
- 10 Triton X-100 0.1%  
Stock solution of TBS (10X) is prepared, and Triton X-100 is added to the buffer during dilution.
- g. HNTG buffer:
- HEPES 20 mM
- 15 NaCl 150 mM (pH 7.2/HCl 1N)  
Glycerol 10%  
Triton X-100 0.2%  
Stock solution (5X) is prepared and kept at 4°C.
- h. EDTA/HCl: 0.5 M pH 7.0 (NaOH) as 100X stock.
- 20 i.  $\text{Na}_3\text{VO}_4$ : 0.5 M as 100X stock and aliquots are kept at 80°C.
- j.  $\text{Na}_4\text{P}_2\text{O}_7$ : 0.2 M as 100X stock.
- k. Insulin-like growth factor-1 from Promega (Cat# G5111).
- 25 l. Rabbit polyclonal anti-phosphotyrosine antiserum.
- m. Goat anti-rabbit IgG, POD conjugate (detection antibody), Tago (Cat. No. 4520, Lot No. 1802): Tago, Inc., Burlingame, CA.
- n. ABTS (2,2'-azinobis(3-ethylbenzthiazolinesulfonic acid)) solution:
- 30 Citric acid 100 mM  
 $\text{Na}_2\text{HPO}_4$  250 mM (pH 4.0/1 N HCl)  
ABTS 0.5 mg/ml  
ABTS solution should be kept in dark and 4°C. The

WO 00/08202

267

PCT/US99/17845

solution should be discarded when it turns green.

o. Hydrogen Peroxide: 30% solution is kept in the dark and at 4°C.

**Procedure.**

5 All the following steps are conducted at room temperature unless specifically indicated otherwise. All ELISA plate washings are performed by rinsing the plate with tap water three times, followed by one TBST rinse. Pat plate dry with paper towels.

10 **Cell Seeding:**

1. The cells, grown in tissue culture dish (Corning 25020-100) to 80-90% confluence, are harvested with Trypsin-EDTA (0.25%, 0.5 ml/D-100, GIBCO).

2. Resuspend the cells in fresh DMEM + 10% FBS +  
15 2mM L-Glutamine, and transfer to 96-well tissue culture plate (Corning, 25806-96) at 20,000 cells/well (100 µl/well). Incubate for 1 day then replace medium to serum-free medium (90/µl) and incubate in 5% CO<sub>2</sub> and 37°C overnight.

**ELISA Plate Coating and Blocking:**

20 1. Coat the ELISA plate (Corning 25805-96) with Anti-IGF-1R Antibody at 0.5 µg/well in 100 µl PBS at least 2 hours.

2. Remove the coating solution, and replace with 100 µl Blocking Buffer, and shake for 30 minutes. Remove the  
25 blocking buffer and wash the plate just before adding lysate.

**Assay Procedures:**

1. The drugs are tested under serum-free condition.  
2. Dilute drug stock (in 100% DMSO) 1:10 with DMEM in 96-well poly-propylene plate, and transfer 10 µl/well of this  
30 solution to the cells to achieve final drug dilution 1:100, and final DMSO concentration of 1.0%. Incubate the cells in 5% CO<sub>2</sub> at 37°C for 2 hours.

3. Prepare fresh cell lysis buffer (HNTG\*)

WO 00/08202

268

PCT/US99/17845 ,

	HNTG	2 ml
	EDTA	0.1 ml
	Na <sub>3</sub> VO <sub>4</sub>	0.1 ml
	Na <sub>4</sub> (P <sub>2</sub> O <sub>7</sub> )	0.1 ml
5	H <sub>2</sub> O	7.3 ml

4. After drug incubation for two hours, transfer 10  $\mu$ l/well of 200nM IGF-1 Ligand in PBS to the cells (Final Conc. is 20 nM), and incubate at 5% CO<sub>2</sub> at 37°C for 10 minutes.

5. Remove media and add 100  $\mu$ l/well HNTG\* and shake for 10 minutes. Look at cells under microscope to see if they are adequately lysed.

6. Use a 12-channel pipette to scrape the cells from the plate, and homogenize the lysate by repeated aspiration and dispensing. Transfer all the lysate to the antibody coated ELISA plate, and shake for 1 hour.

7. Remove the lysate, wash the plate, transfer anti-pTyr (1:3,000 with TBST) 100  $\mu$ l/well, and shake for 30 minutes.

8. Remove anti-pTyr, wash the plate, transfer TAGO (1:3,000 with TBST) 100  $\mu$ l/well, and shake for 30 minutes.

9. Remove detection antibody, wash the plate, and transfer fresh ABTS/H<sub>2</sub>O<sub>2</sub> (1.2  $\mu$ l H<sub>2</sub>O<sub>2</sub> to 10 ml ABTS) 100  $\mu$ l/well to the plate to start color development.

Measure OD at 410 nm with a reference wavelength of 630 nm in Dynatec MR5000.

#### EGFR Assay

EGF Receptor kinase activity in cells genetically engineered to express human EGF-R can be measured as described below:

#### 30 Materials and Reagents.

a. EGF Ligand: stock concentration = 16.5  $\mu$ M, EGF 201, TOYOBO, Co., Ltd. Japan.

b. 05-101 (UBI) (a monoclonal antibody recognizing an

WO 00/08202

269

PCT/US99/17845 ,

EGFR extracellular domain).

c. Anti-phosphotyrosine antibody (anti-Ptyr)  
(polyclonal).

d. Detection antibody: Goat anti-rabbit IgG horse  
5 radish peroxidase conjugate, TAGO, Inc., Burlingame, CA.

e. TBST buffer:

Tris-HCl, pH 7	50 mM
NaCl	150 mM
Triton X-100	0.1

10 f. HNTG 5X stock:

HEPES	0.1 M
NaCl	0.75 M
Glycerol	50
Triton X-100	1.0%

15 g. ABTS stock:

Citric Acid	100 mM
Na <sub>3</sub> VO <sub>4</sub>	250 mM
HCl, conc.	4.0 pH
ABTS*	0.5 mg/ml

20 Keep solution in dark at 4°C until used.

h. Stock reagents of:

EDTA 100 mM pH 7.0
Na <sub>3</sub> VO <sub>4</sub> 0.5 M
Na <sub>4</sub> (P <sub>2</sub> O <sub>7</sub> ) 0.2 M

25 **Procedure.**

**Pre-coat ELISA Plate**

1. Coat ELISA plates (Corning, 96 well, Cat. #25805-96)  
with 05-101 antibody at 0.5 µg per well in PBS, 150 µl final  
volume/well, and store overnight at 4°C. Coated plates are  
30 good for up to 10 days when stored at 4°C.

2. On day of use, remove coating buffer and replace  
with blocking buffer (5% Carnation Instant NonFat Dry Milk in  
PBS). Incubate the plate, shaking, at room temperature (about

WO 00/08202

270

PCT/US99/17845 ,

23°C to 25°C) for 30 minutes. Just prior to use, remove blocking buffer and wash plate 4 times with TBST buffer.

#### Seeding Cells

1. NIH 3T3/C7 cell line (Honegger, et al., Cell 51:199-  
5 209, 1987) can be use for this assay.

2. Choose dishes having 80-90% confluence for the experiment. Trypsinize cells and stop reaction by adding 10% CS DMEM medium. Suspend cells in DMEM medium (10% CS DMEM medium) and centrifuge once at 1000 rpm at room temperature  
10 for 5 minutes.

3. Resuspend cells in seeding medium (DMEM, 0.5% bovine serum), and count the cells using trypan blue. Viability above 90% is acceptable. Seed cells in DMEM medium (0.5% bovine serum) at a density of 10,000 cells per well, 100 µl per well,  
15 in a 96 well microtiter plate. Incubate seeded cells in 5% CO<sub>2</sub> at 37°C for about 40 hours.

#### Assay Procedures.

1. Check seeded cells for contamination using an inverted microscope. Dilute test compounds stock (10 mg/ml in  
20 DMSO) 1:10 in DMEM medium, then transfer 5 µl to a test well for a test compounds drug dilution of 1:200 and a final DMSO concentration of 1%. Control wells receive DMSO alone. Incubate in 5% CO<sub>2</sub> at 37°C for one hour.

2. Prepare EGF ligand: dilute stock EGF in DMEM so that  
25 upon transfer of 10 µl dilute EGF (1:12 dilution), 25 nM final concentration is attained.

3. Prepare fresh 10 ml HNTG\* sufficient for 100 µl per well wherein HNTG\* comprises: HNTG stock (2.0 ml), milli-Q H<sub>2</sub>O (7.3 ml), EDTA, 100 mM, pH 7.0 (0.5 ml), Na<sub>3</sub>VO<sub>4</sub> 0.5 M (0.1 ml)  
30 and Na<sub>4</sub>(P<sub>2</sub>O<sub>7</sub>), 0.2 M (0.1 ml).

4. Place on ice.

5. After two hours incubation with drug, add prepared EGF ligand to cells, 10 µl per well, to yield a final

WO 00/08202

271

PCT/US99/17845 ,

concentration of 25 nM. Control wells receive DMEM alone.

Incubate, shaking, at room temperature, for 5 minutes.

6. Remove test compound, EGF, and DMEM. Wash cells twice with PBS. Transfer HNTG\* to cells, 100 µl per well.

5 Place on ice for 5 minutes. Meanwhile, remove blocking buffer from other ELISA plate and wash with TBST as described above.

7. With a pipette tip securely fitted to a micropipettor, scrape cells from plate and homogenize cell material by repeatedly aspirating and dispensing the HNTG\*  
10 lysis buffer. Transfer lysate to a coated, blocked, and washed ELISA plate. Incubate shaking at room temperature for one hour.

8. Remove lysate and wash 4 times with TBST. Transfer freshly diluted anti-Ptyr antibody to ELISA plate at 100 µl  
15 per well. Incubate shaking at room temperature for 30 minutes in the presence of the anti-Ptyr antiserum (1:3000 dilution in TBST).

9. Remove the anti-Ptyr antibody and wash 4 times with TBST. Transfer the freshly diluted TAGO 30 anti-rabbit IgG  
20 antibody to the ELISA plate at 100 µl per well. Incubate shaking at room temperature for 30 minutes (anti-rabbit IgG antibody: 1:3000 dilution in TBST).

10. Remove detection antibody and wash 4 times with TBST. Transfer freshly prepared ABTS/H<sub>2</sub>O<sub>2</sub> solution to ELISA  
25 plate, 100 µl per well. Incubate at room temperature for 20 minutes. ABTS/H<sub>2</sub>O<sub>2</sub> solution: 1.2 µl 30% H<sub>2</sub>O<sub>2</sub> in 10 ml ABTS stock.

11. Stop reaction by adding 50 µl 5N H<sub>2</sub>SO<sub>4</sub> (optional), and determine O.D. at 410 nm.

30 12. The maximal phosphotyrosine signal is determined by subtracting the value of the negative controls from the positive controls. The percent inhibition of phosphotyrosine content for extract-containing wells is then calculated, after

WO 00/08202

272

PCT/US99/17845

subtraction of the negative controls.

**Met Autophosphorylation Assay**

This assay determines Met tyrosine kinase activity by analyzing Met protein tyrosine kinase levels on the Met  
5 receptor.

**Reagents**

a. HNTG (5X stock solution): Dissolve 23.83 g HEPES and 43.83 g NaCl in about 350 ml dH<sub>2</sub>O. Adjust pH to 7.2 with HCl or NaOH, add 500 ml glycerol and 10 ml Triton X-100, mix, add  
10 dH<sub>2</sub>O to 1 L total volume. To make 1 L of 1X working solution add 200 ml 5X stock solution to 800 ml dH<sub>2</sub>O, check and adjust pH as necessary, store at 4°C.

b. PBS (Dulbecco's Phosphate-Buffered Saline), Gibco Cat. # 450-1300EB (1X solution).

15 c. Blocking Buffer: in 500 ml dH<sub>2</sub>O place 100 g BSA, 12.1 g Tris-pH7.5, 58.44 g NaCl and 10 ml Tween-20, dilute to 1 L total volume.

d. Kinase Buffer: To 500 ml dH<sub>2</sub>O add 12.1 g TRIS (pH 7.2), 58.4 g NaCl, 40.7 g MgCl<sub>2</sub>, and 1.9 g EGTA, bring to 1 L  
20 total volume with dH<sub>2</sub>O.

e. PMSF (Phenylmethylsulfonyl fluoride), Sigma Cat. # P-7626, to 435.5 mg, add 100% ethanol to 25 ml total volume, vortex.

25 f. ATP (Bacterial Source), Sigma Cat. # A-7699, store powder at -20°C, to make up solution for use, dissolve 3.31 mg in 1 ml dH<sub>2</sub>O.

g. RC-20H HRPO Conjugated Anti-Phosphotyrosine, Transduction Laboratories Cat. # E120H.

30 h. Pierce 1-Step™ Turbo TMB-ELISA (3,3',5,5'-tetramethylbenzidine, Pierce Cat. # 34022.

i. H<sub>2</sub>SO<sub>4</sub>, add 1 ml conc. (18 N) to 35 ml dH<sub>2</sub>O.

j. TRIS HCL, Fischer Cat. # BP152-5, to 121.14 g of material, add 600 ml MilliQ H<sub>2</sub>O, adjust pH to 7.5 (or 7.2) with HCl, bring volume to 1 L with MilliQ H<sub>2</sub>O.

WO 00/08202

273

PCT/US99/17845

- k. NaCl, Fischer Cat. # S271-10, make up 5M solution.
- l. Tween-20, Fischer Cat. # S337-500.
- m.  $\text{Na}_3\text{VO}_4$ , Fischer Cat. # S454-50, to 1.8 g material add  
80 ml MilliQ  $\text{H}_2\text{O}$ , adjust pH to 10.0 with HCl or NaOH, boil in  
5 microwave, cool, check pH, repeat procedure until pH stable at  
10.0, add MilliQ  $\text{H}_2\text{O}$  to 100 ml total volume, make 1 ml aliquots  
and store at  $-80^\circ\text{C}$ .
- n.  $\text{MgCl}_2$ , Fischer Cat. # M33-500, make up 1M solution.
- o. HEPES, Fischer Cat. # BP310-500, to 200 ml MilliQ  
10  $\text{H}_2\text{O}$ , add 59.6 g material, adjust pH to 7.5, bring volume to 250  
ml total, sterile filter.
- p. Albumin, Bovine (BSA), Sigma Cat. # A-4503, to 30  
grams material add sterile distilled water to make total  
volume of 300 ml, store at  $4^\circ\text{C}$ .
- 15 q. TBST Buffer: to approx. 900 ml  $\text{dH}_2\text{O}$  in a 1 L  
graduated cylinder add 6.057 g TRIS and 8.766 g NaCl, when  
dissolved, adjust pH to 7.2 with HCl, add 1.0 ml Triton X-100  
and bring to 1 L total volume with  $\text{dH}_2\text{O}$ .
- r. Goat Affinity purified antibody Rabbit IgG (whole  
20 molecule), Cappel Cat. # 55641.
- s. Anti h-Met (C-28) rabbit polyclonal IgG antibody,  
Santa Cruz Chemical Cat. # SC-161.
- t. Transiently Transfected EGFR/Met chimeric cells  
(EMR) (Komada, et al., Oncogene, 8:2381-2390 (1993)).
- 25 u. Sodium Carbonate Buffer, ( $\text{Na}_2\text{CO}_3$ , Fischer Cat. #  
S495): to 10.6 g material add 800 ml MilliQ  $\text{H}_2\text{O}$ , when dissolved  
adjust pH to 9.6 with NaOH, bring up to 1 L total volume with  
MilliQ  $\text{H}_2\text{O}$ , filter, store at  $4^\circ\text{C}$ .

#### Procedure

- 30 All of the following steps are conducted at room  
temperature unless it is specifically indicated otherwise. All  
ELISA plate washing is by rinsing 4X with TBST.

#### EMR Lysis

This procedure can be performed the night before or



WO 00/08202

274

PCT/US99/17845

immediately prior to the start of receptor capture.

1. Quick thaw lysates in a 37° C waterbath with a swirling motion until the last crystals disappear.

2. Lyse cell pellet with 1X HNTG containing 1 mM PMSF.

5 Use 3 ml of HNTG per 15 cm dish of cells. Add 1/2 the calculated HNTG volume, vortex the tube for 1 min., add the remaining amount of HNTG, vortex for another min.

3. Balance tubes, centrifuge at 10,000 x g for 10 min at 4°C.

10 4. Pool supernatants, remove an aliquot for protein determination.

5. Quick freeze pooled sample in dry ice/ethanol bath.

This step is performed regardless of whether lysate will be stored overnight or used immediately following protein  
15 determination.

6. Perform protein determination using standard bicinchoninic acid (BCA) method (BCA Assay Reagent Kit from Pierce Chemical Cat. # 23225).

#### ELISA Procedure

20 1. Coat Corning 96 well ELISA plates with 5 µg per well Goat anti-Rabbit antibody in Carbonate Buffer for a total well volume of 50 µl. Store overnight at 4°C.

2. Remove unbound Goat anti-rabbit antibody by inverting plate to remove liquid.

25 3. Add 150 µl of Blocking Buffer to each well. Incubate for 30 min. with shaking.

4. Wash 4X with TBST. Pat plate on a paper towel to remove excess liquid and bubbles.

5. Add 1µg per well of Rabbit anti-Met antibody diluted  
30 in TBST for a total well volume of 100 µl.

6. Dilute lysate in HNTG (90 µg lysate/100µl)

7. Add 100 µl of diluted lysate to each well. Shake for 60 min.

WO 00/08202

275

PCT/US99/17845

8. Wash 4X with TBST. Pat on paper towel to remove excess liquid and bubbles.

9. Add 50 µl of 1X lysate buffer per well.

10. Dilute compounds/extracts 1:10 in 1X Kinase Buffer in a polypropylene 96 well plate.

11. Transfer 5.5 µl of diluted compound to ELISA plate wells. Incubate at room temperature with shaking for 20 min.

12. Add 5.5 µl of 60 µM ATP solution per well. Negative controls do not receive any ATP. Incubate for 90 min., with shaking.

13. Wash 4X with TBST. Pat plate on paper towel to remove excess liquid and bubbles.

14. Add 100 µl per well of RC20 (1:3000 dilution in Blocking Buffer). Incubate 30 min. with shaking.

15. 15. Wash 4X with TBST. Pat plate on paper towel to remove excess liquid and bubbles.

16. Add 100 µl per well of Turbo-TMB. Incubate with shaking for 30-60 min.

17. Add 100 µl per well of 1M H<sub>2</sub>SO<sub>4</sub> to stop reaction.

20. 18. Read assay on Dynatech MR7000 ELISA reader. Test Filter = 450 nm, reference filter = 410 nm.

#### Biochemical src assay

25 This assay is used to determine src protein kinase activity measuring phosphorylation of a biotinylated peptide as the readout.

#### **Materials and Reagents:**

a. Yeast transformed with src (Sugen, Inc., Redwood City, California).

30 b. Cell lysates: Yeast cells expressing src are pelleted, washed once with water, re-pelleted and stored at -80°C until use.

WO 00/08202

276

PCT/US99/17845 ,

c. N-terminus biotinylated EEEYEEYEEYEEYEEYEEY is prepared by standard procedures well known to those skilled in the art.

d. DMSO: Sigma, St. Louis, MO.

5 e. 96 Well ELISA Plate: Corning 96 Well Easy Wash, Modified flat Bottom Plate, Corning Cat. #25805-96.

f. NUNC 96-well V-bottom polypropylene plates for dilution of compounds: Applied Scientific Cat. # A-72092.

g. Vecastain ELITE ABC reagent: Vector, Burlingame, CA.

10 h. Anti-src (327) mab: Schizosaccharomyces Pombe is used to express recombinant Src (Superti-Furga, et al., EMBO J., 12:2625-2634, Superti-Furga, et al., Nature Biochem., 14:600-605). S. Pombe strain SP200 (h-s leul.32 ura4 ade210) is grown as described and transformations are pRSP expression  
15 plasmids are done by the lithium acetate method (Superti-Furga, supra). Cells are grown in the presence of 1  $\mu$ M thiamine to repress expression from the nmt1 promoter or in the absence of thiamine to induce expression.

i. Monoclonal anti-phosphotyrosine, UBI 05-321 (UB40  
20 may be used instead).

j. Turbo TMB-ELISA peroxidase substrate: Pierce Chemical.

#### Buffer Solutions

a. PBS (Dulbecco's Phosphate-Buffered Saline): GIBCO  
25 PBS, GIBCO Cat. # 450-1300EB.

b. Blocking Buffer: 5% Non-fat milk (Carnation) in PBS.

c. Carbonate Buffer:  $\text{Na}_2\text{CO}_3$  from Fischer, Cat. # S495, make up 100 mM stock solution.

d. Kinase Buffer: 1.0 ml (from 1M stock solution)  $\text{MgCl}_2$ ,  
30 0.2 ml (from a 1M stock solution)  $\text{MnCl}_2$ , 0.2 ml (from a 1M stock solution) DTT, 5.0 ml (from a 1M stock solution) HEPES, 0.1 ml TX-100, bring to 10 ml total volume with MilliQ  $\text{H}_2\text{O}$ .

WO 00/08202

277

PCT/US99/17845 ,

e. Lysis Buffer: 5.0 HEPES (from 1M stock solution.),  
2.74 ml NaCl (from 5M stock solution), 10 ml glycerol, 1.0 ml  
TX-100, 0.4 ml EDTA (from a 100 mM stock solution), 1.0 ml  
PMSF (from a 100 mM stock solution), 0.1 ml  $\text{Na}_3\text{VO}_4$  (from a 0.1  
5 M stock solution), bring to 100 ml total volume with MilliQ  
 $\text{H}_2\text{O}$ .

f. ATP: Sigma Cat. # A-7699, make up 10 mM stock  
solution (5.51 mg/ml).

g TRIS-HCl: Fischer Cat. # BP 152-5, to 600 ml MilliQ  
10  $\text{H}_2\text{O}$  add 121.14 g material, adjust pH to 7.5 with HCl, bring to  
1 L total volume with MilliQ  $\text{H}_2\text{O}$ .

h. NaCl: Fischer Cat. # S271-10, Make up 5M stock  
solution with MilliQ  $\text{H}_2\text{O}$ .

i.  $\text{Na}_3\text{VO}_4$ : Fischer Cat. # S454-50, to 80 ml MilliQ  $\text{H}_2\text{O}$ ,  
15 add 1.8 g material, adjust pH to 10.0 with HCl or NaOH, boil  
in a microwave, cool, check pH, repeat pH adjustment until pH  
remains stable after heating/cooling cycle, bring to 100 ml  
total volume with MilliQ  $\text{H}_2\text{O}$ , make 1 ml aliquots and store at -  
80°C.

20 j.  $\text{MgCl}_2$ : Fischer Cat. # M33-500, make up 1M stock  
solution with MilliQ  $\text{H}_2\text{O}$ .

k. HEPES: Fischer Cat. # BP 310-500, to 200 ml MilliQ  
 $\text{H}_2\text{O}$ , add 59.6 g material, adjust pH to 7.5, bring to 250 ml  
total volume with MilliQ  $\text{H}_2\text{O}$ , sterile filter (1M stock  
25 solution).

l. TBST Buffer: TBST Buffer: To 900 ml  $\text{dH}_2\text{O}$  add 6.057  
g TRIS and 8.766 g NaCl, adjust pH to 7.2 with HCl, add 1.0 ml  
Triton-X100, bring to 1 L total volume with  $\text{dH}_2\text{O}$ .

m.  $\text{MnCl}_2$ : Fischer Cat. # M87-100, make up 1M stock  
30 solution with MilliQ  $\text{H}_2\text{O}$ .

n. DTT: Fischer Cat. # BP172-5.

o. TBS (TRIS Buffered Saline): to 900 ml MilliQ  $\text{H}_2\text{O}$  add  
6.057 g TRIS and 8.777 g NaCl, bring to 1 L total volume with  
MilliQ  $\text{H}_2\text{O}$ .

WO 00/08202

278

PCT/US99/17845

p. Kinase Reaction Mixture: Amount per assay plate (100 wells): 1.0 ml Kinase Buffer, 200 µg GST-ζ, bring to final volume of 8.0 ml with MilliQ H<sub>2</sub>O.

q. Biotin labeled EEEYEEYEEYEEYEEY: Make peptide  
5 stock solution (1mM, 2.98 mg/ml) in water fresh just before use.

r. Vectastain ELITE ABC reagent: To prepare 14 ml of working reagent, add 1 drop of reagent A to 15 ml TBST and invert tube several times to mix. Then add 1 drop of reagent  
10 B. Put tube on orbital shaker at room temperature and mix for 30 minutes.

**Procedures:**

**Preparation of src coated ELISA plate.**

1. Coat ELISA plate with 0.5 µg/well anti-src mab in  
15 100 µl of pH 9.6 sodium carbonate buffer, hold at 4°C overnight.

2. Wash wells once with PBS.

3. Block plate with 0.15 ml 5% milk in PBS for 30 min.  
at room temperature.

20 4. Wash plate 5X with PBS.

5. Add 10 µg/well of src transformed yeast lysates diluted in Lysis Buffer (0.1 ml total volume per well).  
(Amount of lysate may vary between batches.) Shake plate for  
20 minutes at room temperature.

25 **Preparation of phosphotyrosine antibody-coated ELISA plate.**

1. 4G10 plate: coat 0.5 µg/well 4G10 in 100 µl PBS overnight at 4°C and block with 150 µl of 5% milk in PBS for 30 minutes at room temperature.

30 **Kinase assay procedure.**

1. Remove unbound proteins from plates and wash plates 5X with PBS.

WO 00/08202

279

PCT/US99/17845 ;

2. Add 0.08 ml Kinase Reaction Mixture per well (containing 10  $\mu$ l of 10X Kinase Buffer and 10  $\mu$ M (final concentration) biotin-EEEEEEEEEEEEEEEEEEEE per well diluted in water.

5        3. Add 10  $\mu$ l of compound diluted in water containing 10% DMSO and pre-incubate for 15 minutes at room temperature.

4. Start kinase reaction by adding 10  $\mu$ l/well of 0.05 mM ATP in water (5  $\mu$ M ATP final).

5. Shake ELISA plate for 15 min. at room temperature.

10       6. Stop kinase reaction by adding 10  $\mu$ l of 0.5 M EDTA per well.

7. Transfer 90  $\mu$ l supernatant to a blocked 4G10 coated ELISA plate.

15       8. Incubate for 30 min. while shaking at room temperature.

9. Wash plate 5X with TBST.

10. Incubate with Vectastain ELITE ABC reagent (100  $\mu$ l/well) for 30 min. at room temperature.

11. Wash the wells 5X with TBST.

20       12. Develop with Turbo TMB.

#### Biochemical lck Assay

This assay is used to determine lck protein kinase activities measuring phosphorylation of GST- $\zeta$  as the readout.

#### **Materials and Reagents:**

25       a. Yeast transformed with lck. Schizosaccharomyces Pombe is used to express recombinant Lck (Superti-Furga, et al., EMBO J, 12:2625-2634, Superti-Furga, et al., Nature Biotech., 14:600-605). S. Pombe strain SP200 (h-s leul.32 ura4 ade210) is grown as described and transformations with  
30       pRSP expression plasmids are done by the lithium acetate method (Superti-Furga, supra). Cells are grown in the presence of 1  $\mu$ M thiamine to induce expression.

WO 00/08202

280

PCT/US99/17845 ,

b. Cell lysates: Yeast cells expressing lck are pelleted, washed once in water, re-pelleted and stored frozen at -80°C until use.

c. GST- $\zeta$ : DNA encoding for GST- $\zeta$  fusion protein for  
5 expression in bacteria obtained from Arthur Weiss of the  
Howard Hughes Medical Institute at the University of  
California, San Francisco. Transformed bacteria are grown  
overnight while shaking at 25°C. GST- $\zeta$  is purified by  
glutathione affinity chromatography, Pharmacia, Alameda, CA.

10 d. DMSO: Sigma, St. Louis, MO.

e. 96-Well ELISA plate: Corning 96 Well Easy Wash,  
Modified Flat Bottom Plate, Corning Cat. #25805-96.

f. NUNC 96-well V-bottom polypropylene plates for  
dilution of compounds: Applied Scientific Cat. # AS-72092.

15 g. Purified Rabbit anti-GST antiserum: Amrad  
Corporation (Australia) Cat. #90001605.

h. Goat anti-Rabbit-IgG-HRP: Amersham Cat. # V010301.

i. Sheep anti-mouse IgG (H+L): Jackson Labs Cat. #  
5215-005-003.

20 j. Anti-Lck (3A5) mab: Santa Cruz Biotechnology Cat #  
sc-433.

k. Monoclonal anti-phosphotyrosine UBI 05-321 (UB40 may  
be used instead).

**Buffer solutions:**

25 a. PBS (Dulbecco's Phosphate-Buffered Saline) 1X  
solution: GIBCO PBS, GIBCO Cat. # 450-1300EB.

b. Blocking Buffer: 100 g. BSA, 12.1 g. TRIS (pH7.5),  
58.44 g NaCl, 10 ml Tween-20, bring up to 1 L total volume  
with MilliQ H<sub>2</sub>O.

30 c. Carbonate Buffer: Na<sub>2</sub>CO<sub>3</sub> from Fischer, Cat. # S495,  
make up 100 mM solution with MilliQ H<sub>2</sub>O.

d. Kinase Buffer: 1.0 ml (from 1M stock solution) MgCl<sub>2</sub>,  
0.2 ml (from a 1M stock solution) MnCl<sub>2</sub>, 0.2 ml (from a 1M

WO 00/08202

281

PCT/US99/17845 ,

stock solution) DTT, 5.0 ml (from a 1M stock solution) HEPES, 0.1 ml TX-100, bring to 10 ml total volume with MilliQ H<sub>2</sub>O.

e. Lysis Buffer: 5.0 HEPES (from 1M stock solution.), 2.74 ml NaCl (from 5M stock solution), 10 ml glycerol, 1.0 ml TX-100, 0.4 ml EDTA (from a 100 mM stock solution), 1.0 ml PMSF (from a 100 mM stock solution), 0.1 ml Na<sub>3</sub>VO<sub>4</sub> (from a 0.1 M stock solution), bring to 100 ml total volume with MilliQ H<sub>2</sub>O.

f. ATP: Sigma Cat. # A-7699, make up 10 mM stock solution (5.51 mg/ml).

g TRIS-HCl: Fischer Cat. # BP 152-5, to 600 ml MilliQ H<sub>2</sub>O add 121.14 g material, adjust pH to 7.5 with HCl, bring to 1 L total volume with MilliQ H<sub>2</sub>O.

h. NaCl: Fischer Cat. # S271-10, Make up 5M stock solution with MilliQ H<sub>2</sub>O.

i Na<sub>3</sub>VO<sub>4</sub>: Fischer Cat. # S454-50, to 80 ml MilliQ H<sub>2</sub>O, add 1.8 g material, adjust pH to 10.0 with HCl or NaOH, boil in a microwave, cool, check pH, repeat pH adjustment until pH remains stable after heating/cooling cycle, bring to 100 ml total volume with MilliQ H<sub>2</sub>O, make 1 ml aliquots and store at -80°C.

j. MgCl<sub>2</sub>: Fischer Cat. # M33-500, make up 1M stock solution with MilliQ H<sub>2</sub>O.

k. HEPES: Fischer Cat. # BP 310-500, to 200 ml MilliQ H<sub>2</sub>O, add 59.6 g material, adjust pH to 7.5, bring to 250 ml total volume with MilliQ H<sub>2</sub>O, sterile filter (1M stock solution).

l. Albumin, Bovine (BSA), Sigma Cat. # A4503, to 150 ml MilliQ H<sub>2</sub>O add 30 g material, bring 300 ml total volume with MilliQ H<sub>2</sub>O, filter through 0.22 µm filter, store at 4°C.

m. TBST Buffer: To 900 ml dH<sub>2</sub>O add 6.057 g TRIS and 8.766 g NaCl, adjust pH to 7.2 with HCl, add 1.0 ml Triton-X100, bring to 1 L total volume with dH<sub>2</sub>O.



WO 00/08202

282

PCT/US99/17845 ,

n.  $\text{MnCl}_2$ : Fischer Cat. # M87-100, make up 1M stock solution with MilliQ  $\text{H}_2\text{O}$ .

o. DTT: Fischer Cat. # BP172-5.

p. TBS (TRIS Buffered Saline): to 900 ml MilliQ  $\text{H}_2\text{O}$  add  
5 6.057 g TRIS and 8.777 g NaCl, bring to 1 L total volume with MilliQ  $\text{H}_2\text{O}$ .

q Kinase Reaction Mixture: Amount per assay plate (100 wells): 1.0 ml Kinase Buffer, 200  $\mu\text{g}$  GST- $\zeta$ , bring to final volume of 8.0 ml with MilliQ  $\text{H}_2\text{O}$ .

10 **Procedures:**

**Preparation of Lck coated ELISA plate.**

1. Coat 2.0  $\mu\text{g}$ /well Sheep anti-mouse IgG in 100  $\mu\text{l}$  of pH 9.6 sodium carbonate buffer at 4°C overnight.

2. Wash well once with PBS.

15 3. Block plate with 0.15 ml of blocking Buffer for 30 min. at room temp.

4. Wash plate 5X with PBS.

5. Add 0.5  $\mu\text{g}$ /well of anti-lck (mab 3A5) in 0.1 ml PBS at room temperature for 1-2 hours.

20 6. Wash plate 5X with PBS.

7. Add 20  $\mu\text{g}$ /well of lck transformed yeast lysates diluted in Lysis Buffer (0.1 ml total volume per well). Shake plate at 4°C overnight to prevent loss of activity.

25 **Preparation of phosphotyrosine antibody-coated ELISA plate.**

1. UB40 plate: 1.0  $\mu\text{g}$ /well UB40 in 100  $\mu\text{l}$  of PBS overnight at 4°C and block with 150  $\mu\text{l}$  of Blocking Buffer for at least 1 hour.

**Kinase assay procedure.**

30 1. Remove unbound proteins from plates and wash plates 5X with PBS.

WO 00/08202

283

PCT/US99/17845

2. Add 0.08 ml Kinase Reaction Mixture per well  
(containing 10  $\mu$ l of 10X Kinase Buffer and 2  $\mu$ g GST- $\zeta$  per well  
diluted with water).

3. Add 10  $\mu$ l of compound diluted in water containing  
5 10% DMSO and pre-incubate for 15 minutes at room temperature.

4. Start kinase reaction by adding 10 $\mu$ l/well of 0.1 mM  
ATP in water (10  $\mu$ M ATP final).

5. Shake ELISA plate for 60 min. at room temperature.

6. Stop kinase reaction by adding 10  $\mu$ l of 0.5 M EDTA  
10 per well.

7. Transfer 90  $\mu$ l supernatant to a blocked 4G10 coated  
ELISA plate from section B, above.

8. Incubate while shaking for 30 min. at room  
temperature.

15 9. Wash plate 5X with TBST.

10. Incubate with Rabbit anti-GST antibody at 1:5000  
dilution in 100  $\mu$ l TBST for 30 min. at room temperature.

11. Wash the wells 5X with TBST.

12. Incubate with Goat anti-Rabbit-IgG-HRP at 1:20,000  
20 dilution in 100  $\mu$ l of TBST for 30 min. at room temperature.

13. Wash the wells 5X with TBST.

14. Develop with Turbo TMB.

**Assay measuring phosphorylating function of RAF.**

The following assay reports the amount of RAF-catalyzed  
25 phosphorylation of its target protein MEK as well as MEK's  
target MAPK. The RAF gene sequence is described in Bonner et  
al., 1985, Molec. Cell. Biol., 5:1400-1407, and is readily  
accessible in multiple gene sequence data banks. Construction  
of the nucleic acid vector and cell lines utilized for this  
30 portion of the invention are fully described in Morrison et  
al., 1988, Proc. Natl. Acad. Sci. USA, 85:8855-8859.

WO 00/08202

284

PCT/US99/17845 ,

**Materials and Reagents**

1. Sf9 (*Spodoptera frugiperda*) cells, GIBCO-BRL, Gaithersburg, MD.
2. RIPA buffer: 20 mM Tris/HCl pH 7.4, 137 mM NaCl, 10% glycerol, 1 mM PMSF, 5 mg/L Aprotinin, 0.5 % Triton X-100,
3. Thioredoxin-MEK fusion protein (T-MEK): T-MEK expression and purification by affinity chromatography are performed according to the manufacturer's procedures. Catalog# K 350-01 and R 350-40, Invitrogen Corp., San Diego, CA
4. His-MAPK (ERK 2), His-tagged MAPK is expressed in XL1 Blue cells transformed with pUC18 vector encoding His-MAPK. His-MAPK is purified by Ni-affinity chromatography. Cat# 27-4949-01, Pharmacia, Alameda, CA, as described herein.
5. Sheep anti mouse IgG: Jackson laboratories, West Grove, PA. Catalog, # 515-006-008, Lot# 28563
6. RAF-1 protein kinase specific antibody: URP2653 from UBI.
7. Coating buffer: PBS, phosphate buffered saline, GIBCO-BRL, Gaithersburg, MD.
8. Wash buffer: TBST (50 mM Tris/HCL pH 7.2, 150 mM NaCl, 0.1 % Triton X-100).
9. Block buffer: TBST, 0.1 % ethanolamine pH 7.4
10. DMSO, Sigma, St. Louis, MO
11. Kinase buffer (KB): 20 mM HEPES/HCl pH 7.2, 150 mM NaCl, 0.1 % Triton X-100, 1 mM PMSF, 5 mg/L Aprotinin, 75 mM sodium orthovanadate, 0.5 mM DTT and 10 mM MgCl<sub>2</sub>.
12. ATP mix: 100 mM MgCl<sub>2</sub>, 300 mM ATP, 10 mCi  $\gamma^{33}\text{P}$  ATP (Dupont-NEN)/mL.
- 13 Stop solution: 1% phosphoric acid, Fisher, Pittsburgh, PA.
14. Wallac Cellulose Phosphate Filter mats, Wallac, Turku, Finland.

WO 00/08202

285

PCT/US99/17845 ,

15. Filter wash solution: 1% phosphoric acid, Fisher, Pittsburgh, PA.

16. Tomtec plate harvester, Wallac, Turku, Finland.

17. Wallac beta plate reader # 1205, Wallac, Turku, Finland.

18. NUNC 96-well V bottom polypropylene plates for compounds Applied Scientific Catalog # AS-72092.

#### Procedure

All of the following steps are conducted at room temperature unless specifically indicated otherwise.

1. ELISA plate coating: ELISA wells are coated with 100 ml of Sheep anti mouse affinity purified antiserum (1 mg/100 mL coating buffer) over night at 4° C. ELISA plates can be used for two weeks when stored at 4° C.

2. Invert the plate and remove liquid. Add 100 mL of blocking solution and incubate for 30 min.

3. Remove blocking solution and wash four times with wash buffer. Pat the plate on a paper towel to remove excess liquid.

4. Add 1 mg of antibody specific for RAF-1 to each well and incubate for 1 hour. Wash as described in step 3.

5. Thaw lysates from RAS/RAF infected Sf9 cells and dilute with TBST to 10 mg/100 mL. Add 10 mg of diluted lysate to the wells and incubate for 1 hour. Shake the plate during incubation. Negative controls receive no lysate. Lysates from RAS/RAF infected Sf9 insect cells are prepared after cells are infected with recombinant baculoviruses at a MOI of 5 for each virus, and harvested 48 hours later. The cells are washed once with PBS and lysed in RIPA buffer. Insoluble material is removed by centrifugation (5 min at 10,000 x g). Aliquots of lysates are frozen in dry ice/ethanol and stored at -80 °C until use.

WO 00/08202

286

PCT/US99/17845

6. Remove non-bound material and wash as outlined above (step 3).

7. Add 2 mg of T-MEK and 2 mg of His-MAEPK per well and adjust the volume to 40 ml with kinase buffer. Methods for  
5 purifying T-MEK and MAPK from cell extracts are provided herein by example.

8. Pre-dilute compounds (stock solution 10 mg/ml DMSO) or extracts 20 fold in TBST plus 1% DMSO. Add 5 ml of the pre-diluted compounds/extracts to the wells described in step  
10 6. Incubate for 20 min. Controls receive no drug.

9. Start the kinase reaction by addition of 5 ml ATP mix, Shake the plates on an ELISA plate shaker during incubation.

10. Stop the kinase reaction after 60 min by addition of  
15 30 mL stop solution to each well.

11. Place the phosphocellulose mat and the ELISA plate in the Tomtec plate harvester. Harvest and wash the filter with the filter wash solution according to the manufacturer's recommendation. Dry the filter mats. Seal the filter mats  
20 and place them in the holder. Insert the holder into radioactive detection apparatus and quantify the radioactive phosphorous on the filter mats.

Alternatively, 40 mL aliquots from individual wells of the assay plate can be transferred to the corresponding  
25 positions on the phosphocellulose filter mat. After air drying the filters, put the filters in a tray. Gently rock the tray, changing the wash solution at 15 min intervals for 1 hour. Air-dry the filter mats. Seal the filter mats and place them in a holder suitable for measuring the radioactive  
30 phosphorous in the samples. Insert the holder into a detection device and quantify the radioactive phosphorous on the filter mats.

WO 00/08202

287

PCT/US99/17845 ,

**CDK2/Cyclin A - Inhibition Assay**

This assay analyzes the protein kinase activity of CDK2 in exogenous substrate.

**Reagents:**

5        A.    Buffer A: (80 mM Tris ( pH 7.2), 40 mM MgCl<sub>2</sub>), 4.84 g. Tris (F.W. =121.1 g/mol), 4.07 g. MgCl<sub>2</sub> (F.W.=203.31 g/mol) dissolved in 500 ml H<sub>2</sub>O. Adjust pH to 7.2 with HCl.

      B.    Histone H1 solution (0.45 mg/ml Histone H1 and 20 mM HEPES pH 7.2: 5 mg Histone H1 (Boehinger Mannheim) in 11.111  
10 ml 20 mM HEPES pH 7.2 (477 mg HEPES (F.W.= 238.3 g/mol) dissolved in 100 ml ddH<sub>2</sub>O, stored in 1 ml aliquots at -80° C.

      C.    ATP solution (60 µM ATP, 300 µg/ml BSA, 3 mM DTT): 120 µl 10 mM ATP, 600 µl 10 mg/ml BSA to 20 ml, stored in 1 ml aliquots at -80° C.

15        D.    CDK2 solution: cdk2/cyclin A in 10 mM HEPES pH 7.2, 25 mM NaCl, 0.5 mM DTT, 10% glycerol, stored in 9 µl aliquots at  
-80° C.

**Protocol**

20        1.    Prepare solutions of inhibitors at three times the desired final assay concentration in ddH<sub>2</sub>O/15% DMSO by volume.

      2.    Dispense 20 µl of inhibitors to wells of polypropylene 96-well plates (or 20 µl 15% DMSO for positive and negative controls).

25        3.    Thaw Histone H1 solution (1 ml/plate), ATP solution (1 ml/plate plus 1 aliquot for negative control), and CDK2 solution (9 µl/plate). Keep CDK2 on ice until use. Aliquot CDK2 solution appropriately to avoid repeated freeze-thaw cycles.

30        4.    Dilute 9 µl CDK2 solution into 2.1 ml Buffer A (per plate). Mix. Dispense 20 µl into each well.

WO 00/08202

288

PCT/US99/17845 ,

5. Mix 1 ml Histone H1 solution with 1 ml ATP solution (per plate) into a 10 ml screw cap tube. Add  $\gamma^{33}\text{P}$  ATP to a concentration of 0.15  $\mu\text{Ci}/20\mu\text{l}$  (0.15  $\mu\text{Ci}/\text{well}$  in assay). Mix carefully to avoid BSA frothing. Add 20  $\mu\text{l}$  to appropriate wells. Mix plates on plate shaker. For negative control, mix ATP solution with an equal amount of 20 mM HEPES pH 7.2 and add  $\gamma^{33}\text{P}$  ATP to a concentration of 0.15  $\mu\text{Ci}/20\mu\text{l}$  solution. Add 20  $\mu\text{l}$  to appropriate wells.
6. Let reactions proceed for 60 minutes.
7. Add 35  $\mu\text{l}$  10% TCA to each well. Mix plates on plate shaker.
8. Spot 40  $\mu\text{l}$  of each sample onto P30 filter mat squares. Allow mats to dry (approx. 10-20 minutes).
- 9 Wash filter mats 4 X 10 minutes with 250 ml 1% phosphoric acid (10 ml phosphoric acid per liter  $\text{ddH}_2\text{O}$ ).
10. Count filter mats with beta plate reader.

#### Cellular/Biologic Assays

##### PDGF-Induced BrdU Incorporation Assay

##### Materials and Reagents:

- (1) PDGF: human PDGF B/B, 1276-956, Boehringer Mannheim, Germany.
- (2) BrdU Labeling Reagent: 10 mM, in PBS (pH7.4), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (3) FixDenat: fixation solution (ready to use), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (4) Anti-BrdU-POD: mouse monoclonal antibody conjugated with peroxidase, Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (5) TMB Substrate Solution: tetramethylbenzidine (TMB), ready to use, Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (6) PBS Washing Solution : 1X PBS, pH 7.4 (Sugen, Inc., Redwood City, California).

WO 00/08202

289

PCT/US99/17845 ,

(7) Albumin, Bovine (BSA): fraction V powder, A-8551, Sigma Chemical Co., USA.

(8) 3T3 cell line genetically engineered to express human PDGF-R.

5           **Protocol**

(1) Cells are seeded at 8000 cells/well in DMEM, 10% CS, 2mM Gln in a 96 well plate. Cells are incubated overnight at 37°C in 5% CO<sub>2</sub>.

10           (2) After 24 hours, the cells are washed with PBS, and then are serum starved in serum free medium (0%CS DMEM with 0.1% BSA) for 24 hours.

            (3) On day 3, ligand (PDGF, 3.8 nM, prepared in DMEM with 0.1% BSA) and test compounds are added to the cells simultaneously. The negative control wells receive serum free  
15           DMEM with 0.1% BSA only, the positive control cells receive the ligand (PDGF) but no test compound. Test compounds are prepared in serum free DMEM with ligand in a 96 well plate, and serially diluted for 7 test concentrations.

            (4) After 20 hours of ligand activation, diluted BrdU  
20           labeling reagent (1:100 in DMEM, 0.1% BSA) is added and the cells are incubated with BrdU (final concentration=10 µM) for 1.5 hours.

            (5) After incubation with labeling reagent, the medium is removed by decanting and tapping the inverted plate on a  
25           paper towel. FixDenat solution is added (50 µl/well) and the plates are incubated at room temperature for 45 minutes on a plate shaker.

            (6) The FixDenat solution is thoroughly removed by decanting and tapping the inverted plate on a paper towel.  
30           Milk is added (5% dehydrated milk in PBS, 200 µl/well) as a blocking solution and the plate is incubated for 30 minutes at room temperature on a plate shaker.

            (7) The blocking solution is removed by decanting and



WO 00/08202

290

PCT/US99/17845 ;

the wells are washed once with PBS. Anti-BrdU-POD solution (1:100 dilution in PBS, 1% BSA) is added (100 µl/well) and the plate is incubated for 90 minutes at room temperature on a plate shaker.

- 5           (8) The antibody conjugate is thoroughly removed by decanting and rinsing the wells 5 times with PBS, and the plate is dried by inverting and tapping on a paper towel.

- (9) TMB substrate solution is added (100 µl/well) and incubated for 20 minutes at room temperature on a plate shaker  
10 until color development is sufficient for photometric detection.

- (10) The absorbance of the samples are measured at 410 nm (in "dual wavelength" mode with a filter reading at 490 nm, as a reference wavelength) on a Dynatech ELISA plate reader.

15           **EGF-Induced BrdU Incorporation Assay**

**Materials and Reagents**

- (1) EGF: mouse EGF, 201, Toyobo, Co., Ltd. Japan.  
          (2) BrdU Labeling Reagent: 10 mM, in PBS (pH7.4), Cat. No. 1 647 229, Boehringer Mannheim, Germany.  
20           (3) FixDenat: fixation solution (ready to use), Cat. No. 1 647 229, Boehringer Mannheim, Germany.  
          (4) Anti-BrdU-POD: mouse monoclonal antibody conjugated with peroxidase, Cat. No. 1 647 229, Boehringer Mannheim, Germany.  
25           (5) TMB Substrate Solution: tetramethylbenzidine (TMB), ready to use, Cat. No. 1 647 229, Boehringer Mannheim, Germany.  
          (6) PBS Washing Solution : 1X PBS, pH 7.4 (Sugen, Inc., Redwood City, California).  
30           (7) Albumin, Bovine (BSA): fraction V powder, A-8551, Sigma Chemical Co., USA.  
          (8) 3T3 cell line genetically engineered to express human EGF-R.

WO 00/08202

291

PCT/US99/17845 ,

**Protocol**

(1) Cells are seeded at 8000 cells/well in 10% CS, 2mM Gln in DMEM, in a 96 well plate. Cells are incubated overnight at 37°C in 5% CO<sub>2</sub>.

5 (2) After 24 hours, the cells are washed with PBS, and then are serum starved in serum free medium (0% CS DMEM with 0.1% BSA) for 24 hours.

(3) On day 3, ligand (EGF, 2 nM, prepared in DMEM with 0.1% BSA) and test compounds are added to the cells  
10 simultaneously. The negative control wells receive serum free DMEM with 0.1% BSA only, the positive control cells receive the ligand (EGF) but no test compound. Test compounds are prepared in serum free DMEM with ligand in a 96 well plate, and serially diluted for 7 test concentrations.

15 (4) After 20 hours of ligand activation, diluted BrdU labeling reagent (1:100 in DMEM, 0.1% BSA) is added and the cells are incubated with BrdU (final concentration = 10 µM) for 1.5 hours.

(5) After incubation with labeling reagent, the medium  
20 is removed by decanting and tapping the inverted plate on a paper towel. FixDenat solution is added (50 µl/well) and the plates are incubated at room temperature for 45 minutes on a plate shaker.

(6) The FixDenat solution is thoroughly removed by  
25 decanting and tapping the inverted plate on a paper towel. Milk is added (5% dehydrated milk in PBS, 200 µl/well) as a blocking solution and the plate is incubated for 30 minutes at room temperature on a plate shaker.

(7) The blocking solution is removed by decanting and  
30 the wells are washed once with PBS. Anti-BrdU-POD solution (1:100 dilution in PBS, 1% BSA) is added (100 µl/well) and the plate is incubated for 90 minutes at room temperature on a plate shaker.

WO 00/08202

292

PCT/US99/17845 ,

(8) The antibody conjugate is thoroughly removed by decanting and rinsing the wells 5 times with PBS, and the plate is dried by inverting and tapping on a paper towel.

(9) TMB substrate solution is added (100 µl/well) and  
5 incubated for 20 minutes at room temperature on a plate shaker until color development is sufficient for photometric detection.

(10) The absorbance of the samples are measured at 410 nm (in "dual wavelength" mode with a filter reading at 490 nm, as  
10 a reference wavelength) on a Dynatech ELISA plate reader.

#### EGF-Induced Her2-Driven BrdU Incorporation

##### **Materials and Reagents**

- (1) EGF: mouse EGF, 201, Toyobo, Co., Ltd. Japan
- (2) BrdU Labeling Reagent: 10 mM, in PBS (pH7.4), Cat.  
15 No. 1 647 229, Boehringer Mannheim, Germany.
- (3) FixDenat: fixation solution (ready to use), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (4) Anti-BrdU-POD: mouse monoclonal antibody conjugated with peroxidase, Cat. No. 1 647 229, Boehringer Mannheim,  
20 Germany.
- (5) TMB Substrate Solution: tetramethylbenzidine (TMB), ready to use, Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (6) PBS Washing Solution: 1X PBS, pH 7.4, made in house.
- 25 (7) Albumin, Bovine (BSA): fraction V powder, A-8551, Sigma Chemical Co., USA.
- (8) 3T3 cell line engineered to express a chimeric receptor having the extra-cellular domain of EGF-R and the intra-cellular domain of Her2.

##### **Protocol**

- (1) Cells are seeded at 8000 cells/well in DMEM, 10% CS, 2mM Gln in a 96- well plate. Cells are incubated overnight at 37° C in 5% CO<sub>2</sub>.

WO 00/08202

293

PCT/US99/17845

(2) After 24 hours, the cells are washed with PBS, and then are serum starved in serum free medium (0% CS DMEM with 0.1% BSA) for 24 hours.

(3) On day 3, ligand (EGF = 2 nM, prepared in DMEM with 0.1% BSA) and test compounds are added to the cells simultaneously. The negative control wells receive serum free DMEM with 0.1% BSA only, the positive control cells receive the ligand (EGF) but no test compound. Test compounds are prepared in serum free DMEM with ligand in a 96 well plate, and serially diluted for 7 test concentrations.

(4) After 20 hours of ligand activation, diluted BrdU labeling reagent (1:100 in DMEM, 0.1% BSA) is added and the cells are incubated with BrdU (final concentration = 10  $\mu$ M) for 1.5 hours.

(5) After incubation with labeling reagent, the medium is removed by decanting and tapping the inverted plate on a paper towel. FixDenat solution is added (50  $\mu$ l/well) and the plates are incubated at room temperature for 45 minutes on a plate shaker.

(6) The FixDenat solution is thoroughly removed by decanting and tapping the inverted plate on a paper towel. Milk is added (5% dehydrated milk in PBS, 200  $\mu$ l/well) as a blocking solution and the plate is incubated for 30 minutes at room temperature on a plate shaker.

(7) The blocking solution is removed by decanting and the wells are washed once with PBS. Anti-BrdU-POD solution (1:100 dilution in PBS, 1% BSA) is added (100  $\mu$ l/well) and the plate is incubated for 90 minutes at room temperature on a plate shaker.

(8) The antibody conjugate is thoroughly removed by decanting and rinsing the wells 5 times with PBS, and the plate is dried by inverting and tapping on a paper towel.

WO 00/08202

294

PCT/US99/17845

(9) TMB substrate solution is added (100 µl/well) and incubated for 20 minutes at room temperature on a plate shaker until color development is sufficient for photometric detection.

- 5 (10) The absorbance of the samples are measured at 410 nm (in "dual wavelength" mode with a filter reading at 490 nm, as a reference wavelength) on a Dynatech ELISA plate reader.

**IGF1-Induced BrdU Incorporation Assay**

**Materials and Reagents**

- 10 (1) IGF1 Ligand: human, recombinant, G511, Promega Corp, USA.
- (2) BrdU Labeling Reagent: 10 mM, in PBS (pH7.4), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (3) FixDenat: fixation solution (ready to use), Cat. No. 15 1 647 229, Boehringer Mannheim, Germany.
- (4) Anti-BrdU-POD: mouse monoclonal antibody conjugated with peroxidase, Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (5) TMB Substrate Solution: tetramethylbenzidine (TMB), 20 ready to use, Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (6) PBS Washing Solution: 1X PBS, pH 7.4 (Sugen, Inc., Redwood City, California).
- (7) Albumin, Bovine (BSA): fraction V powder, A-8551, 25 Sigma Chemical Co., USA.
- (8) 3T3 cell line genetically engineered to express human IGF-1 receptor.

**Protocol**

- 30 (1) Cells are seeded at 8000 cells/well in DMEM, 10% CS, 2mM Gln in a 96- well plate. Cells are incubated overnight at 37°C in 5% CO<sub>2</sub>.

WO 00/08202

295

PCT/US99/17845 ,

(2) After 24 hours, the cells are washed with PBS, and then are serum starved in serum free medium (0%CS DMEM with 0.1% BSA) for 24 hours.

(3) On day 3, ligand (IGF1 = 3.3 nM, prepared in DMEM with 0.1% BSA) and test compounds are added to the cells simultaneously. The negative control wells receive serum free DMEM with 0.1% BSA only, the positive control cells receive the ligand (IGF1) but no test compound. Test compounds are prepared in serum free DMEM with ligand in a 96 well plate, and serially diluted for 7 test concentrations.

(4) After 16 hours of ligand activation, diluted BrdU labeling reagent (1:100 in DMEM, 0.1% BSA) is added and the cells are incubated with BrdU (final concentration=10  $\mu$ M) for 1.5 hours.

(5) After incubation with labeling reagent, the medium is removed by decanting and tapping the inverted plate on a paper towel. FixDenat solution is added (50  $\mu$ l/well) and the plates are incubated at room temperature for 45 minutes on a plate shaker.

(6) The FixDenat solution is thoroughly removed by decanting and tapping the inverted plate on a paper towel. Milk is added (5% dehydrated milk in PBS, 200  $\mu$ l/well) as a blocking solution and the plate is incubated for 30 minutes at room temperature on a plate shaker.

(7) The blocking solution is removed by decanting and the wells are washed once with PBS. Anti-BrdU-POD solution (1:100 dilution in PBS, 1% BSA) is added (100  $\mu$ l/well) and the plate is incubated for 90 minutes at room temperature on a plate shaker.

(8) The antibody conjugate is thoroughly removed by decanting and rinsing the wells 5 times with PBS, and the plate is dried by inverting and tapping on a paper towel.

WO 00/08202

296

PCT/US99/17845 ,

(9) TMB substrate solution is added (100 µl/well) and incubated for 20 minutes at room temperature on a plate shaker until color development is sufficient for photometric detection.

- 5 (10) The absorbance of the samples are measured at 410 nm (in "dual wavelength" mode with a filter reading at 490 nm, as a reference wavelength) on a Dynatech ELISA plate reader.

**FGF-Induced BrdU incorporation Assay**

This assay measures FGF-induced DNA synthesis in  
10 3Tc7/EGFr cells that express endogenous FGF receptors.

**Materials and Reagents:**

1. FGF: human FGF2/bFGF (Gibco BRL, No. 13256-029).
2. BrdU Labeling reagent, (10 mM PBS (pH 7.4),  
Boehringer Mannheim Cat No. 1 647 229).
- 15 3. Fixdenat fixation solution (Boehringer Mannheim Cat  
No. 1 647 229).
4. Anti-BrdU-POD (mouse monoclonal antibody conjugated  
with peroxidase, Boehringer Mannheim Cat. No. 1 647 229).
5. TMB (tetramethylbenzidine, Boehringer Mannheim Cat.  
20 No. 1 647 229).
6. PBS washing solution, pH 7.4 (Sugen, Inc.).
7. Albumin, bovine (BSA), fraction V powder (Sigma  
Chemical Co., Cat. No. A-8551)

**Procedure**

- 25 1. 3T3 engineered cell line: 3T3c7/EGFr.
2. Cells are seeded at 8,000 cells/well in DMEM, 10% CS  
and 2 mM Gln in a 96-well plate. Incubate 24 hours at 37° C in  
5% CO<sub>2</sub>.
3. After 24 hours, wash cells with PBS then serum  
30 starve in serum free medium (0% DMEM, 0.1% BSA) for 24 hours.
4. Add ligand (FGF2 (1.5 nM in DMEM with 0.1% BSA) and  
test compound simultaneously. Negative control wells receive  
serum free DMEM with 0.1% BSA only, positive control wells

WO 00/08202

297

PCT/US99/17845 ,

receive FGF2 ligand but no test compound. Test compounds are prepared in serum-free DMEM with ligand in a 96-well plate and serially diluted to make seven (7) test concentrations.

5. After 20 hours, add diluted BrdU labeling reagent  
5 (1:100 BrdU:DMEM, 0.1% BSA, final concentration is 10  $\mu$ M) to the cells and incubate for 1.5 hours.

6. Decant medium. Remove traces of material with paper towel. Add FixDenat (50  $\mu$ l/well) and incubate at room temperature for 45 minutes on a plate shaker.

10 7. Remove Fixdenat solution. Add blocking solution (5% dehydrated milk in PBS (200  $\mu$ l/well)) and incubate for 30 minutes at room temperature on a plate shaker.

8. Decant blocking solution, wash wells once with PBS. Add anti-BrdU-POD solution (1:100 dilution in PBS, 0.1% BSA),  
15 incubate for 90 minutes at room temperature on a plate shaker.

9. Decant antibody conjugate, rinse wells 5 times with PBS. Dry plate by inverting on paper towel and tapping.

10. Add TMB solution (100  $\mu$ l/well), incubate 20 minutes at room temperature on a plate shaker until color development  
20 is sufficient for photometric detection.

11. Measure absorbance at 410 nm on a Dynatech ELISA plate reader using "Dual wavelength" mode with a filter at 490 nm.

#### Biochemical EGFR Assay

This assay measures the in vitro kinase activity of  
25 EGFR using ELISA.

#### **Materials and Reagents**

1. Corning 96-well Elisa plates (Corning Catalog No. 25805-96).

2. SUMO1 monoclonal anti-EGFR antibody (Biochemistry  
30 Lab, SUGEN, Inc.).

3. PBS (Dulbecco's Phosphate-Buffered Saline, Gibco Catalog No. 450-1300EB).



WO 00/08202

298

PCT/US99/17845

## 4. TBST Buffer

Reagent	M.W.	Working Concentration	Amount per L
---------	------	--------------------------	-----------------

5	Tris	121.14	50 mM	6.057 g
	NaCl	58.44	150 mM	8.766 g
	Triton X-100 NA	0.1%		1.0 ml

## 5. Blocking Buffer:

Reagent	M.W.	Working Concentration	Amount per 100 ml
---------	------	--------------------------	----------------------

10	Carnation Instant Non-Fat Milk	5%	5.0 g
	PBS	NA	100 ml

## 6. A431 cell lysate (Screening Lab, SUGEN, Inc.)

## 7. TBS Buffer:

15	Reagent	M.W.	Working Concentration	Amount per L
	Tris	121.14	50 mM	6.057 g
	NaCl	58.44	150 mM	8.766 g

## 8. TBS + 10% DMSO

20	Reagent	M.W.	Working Concentration	Amount per L
	Tris	121.14	50 mM	1.514 g
	NaCl	58.44	150 mM	2.192 g
	DMSO	NA	10%	25 ml

## 25 9. Adenosine-5'-triphosphate (ATP, from Equine muscle, Sigma Cat. No. A-5394).

Prepare a 1.0 mM solution in dH<sub>2</sub>O. This reagent should be made up immediately prior to use and kept on ice.

10. MnCl<sub>2</sub>.

30 Prepare a 1.0 M stock solution in dH<sub>2</sub>O.

11. ATP/MnCl<sub>2</sub> phosphorylation mix

Reagent	Stock solution	Amount per 10 ml	Working Concentration
---------	-------------------	---------------------	--------------------------

WO 00/08202

299

PCT/US99/17845

ATP	1.0 mM	300 $\mu$ l	30 $\mu$ M
MnCl <sub>2</sub>	1.0 M	500 $\mu$ l	50 mM
dH <sub>2</sub> O			9.2 ml

This reagent should be prepared immediately before  
 5 use and kept on ice.

12. NUNC 96-well V bottom polypropylene plates (Applied Scientific Cat. No. AS-72092).

13. Ethylenediaminetetraacetic acid (EDTA)

Prepare 200 mM working solution in dH<sub>2</sub>O. Adjust to pH  
 10 8.0 with 10 N NaOH.

14. Rabbit polyclonal anti-phosphotyrosine serum (Biochemistry Lab, SUGEN, Inc.)

15. Goat anti-rabbit IgG peroxidase conjugate (Biosource Cat. No. ALI0404)

15 16. ABTS (2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid), Sigma Cat. No. A-1888).

Reagent	M.W.	Working Concentration	Amount per L
Citric Acid	192.12	100 mM	19.21 g
Na <sub>2</sub> HP04	141.96	250 mM	35.49 g
ABTS	NA	0.5 mg/ml	500 mg

Mix first two ingredients in about 900 ml dH<sub>2</sub>O, adjust pH to 4.0 with phosphoric acid. Add ABTS, cover, let sit about 0.5 hr., filter. The solution should be kept in the dark at  
 25 4° C until ready to use.

17. Hydrogen peroxide 30% solution (Fisher Cat. No. H325)

18. ABTS/H<sub>2</sub>O<sub>2</sub>

Mix 15 ml ABTS solution and 2.0  $\mu$ l H<sub>2</sub>O<sub>2</sub>. Prepare 5  
 30 minutes before use.

19. 0.2 M HCl

WO 00/08202

300

PCT/US99/17845 ,

**Procedure**

1. Coat Corning 96 well ELISA plates with 0.5 µg SUMO1 in 100 µl PBS per well, store overnight at 4° C.
2. Remove unbound SUMO1 from wells by inverting plate to remove liquid. Wash 1x with dH<sub>2</sub>O. Pat the plate on a paper towel to remove excess liquid.
3. Add 150 µl of Blocking Buffer to each well. Incubate for 30 min. at room temperature with shaking.
4. Wash plate 3x with deionized water, then once with TBST. Pat plate on a paper towel to remove excess liquid and bubbles.
5. Dilute lysate in PBS (7 µg lysate/100 µl PBS).
6. Add 100 µl of diluted lysate to each well. Shake at room temperature for 60 min.
7. Wash plates as described in 4, above.
8. Add 120 µl TBS to ELISA plate containing captured EGFR.
9. Dilute test compound 1:10 in TBS in 96-well polypropylene plates (ie. 10 µl compound + 90 µl TBS).
10. Add 13.5 µl diluted test compound to ELISA plate. To control wells (wells which do not receive any test compound), add 13.5 µl TBS + 10% DMSO.
11. Incubate for 30 minutes while shaking at room temperature.
12. Add 15 µl phosphorylation mix directly to all wells except negative control well which does not receive ATP/MnCl<sub>2</sub> (final well volume should be approximately 150 µl with 3 µM ATP/5 mM MnCl<sub>2</sub> final concentration in each well.) Incubate 5 minutes while shaking.
13. After 5 minutes, stop reaction by adding 16.5 µl of 200 mM EDTA (pH 8.0) to each well, shaking continuously. After the EDTA has been added, shake for 1 min.

WO 00/08202

301

PCT/US99/17845 ,

14. Wash 4x with deionized water, twice with TBST.
15. Add 100 µl anti-phosphotyrosine (1:3000 dilution in TBST) per well. Incubate 30-45 min. at room temperature, with shaking.
- 5 16. Wash as described in 4, above.
17. Add 100 µl Biosource Goat anti-rabbit IgG peroxidase conjugate (1:2000 dilution in TBST) to each well. Incubate 30 min. at room temperature, with shaking.
18. Wash as described in 4, above.
- 10 19. Add 100 µl of ABTS/H<sub>2</sub>O<sub>2</sub> solution to each well.
20. Incubate 5 to 10 minutes with shaking. Remove any bubbles.
21. If necessary stop reaction with the addition of 100 µl 0.2 M HCl per well.
- 15 22. Read assay on Dynatech MR7000 ELISA reader. Test Filter: 410 nM Reference Filter: 630 Nm.

#### Biochemical PDGFR Assay

This assay measures the in vitro kinase activity of PDGFR using ELISA.

#### 20 **Materials and Reagents**

Unless otherwise noted, the preparation of working solution of the following reagents is the same as that for the Biochemical EGFR assay, above.

- 25 1. Corning 96-well Elisa plates (Corning Catalog No. 25805-96).
2. 28D4C10 monoclonal anti-PDGFR antibody (Biochemistry Lab, SUGEN, Inc.).
3. PBS (Dulbecco's Phosphate-Buffered Saline, Gibco Catalog No. 450-1300EB)
- 30 4. TBST Buffer.
5. Blocking Buffer.
6. PDGFR-β expressing NIH 3T3 cell lysate (Screening Lab, SUGEN, Inc.).

WO 00/08202

302

PCT/US99/17845

7. TBS Buffer.
8. TBS + 10% DMSO.
9. Adenosine-5'-triphosphate (ATP, from Equine muscle, Sigma Cat. No. A-5394).
- 5 10.  $\text{MnCl}_2$ .
11. Kinase buffer phosphorylation mix.
- | Reagent         | Stock<br>solution | Amount<br>per 10 ml | Working<br>Concentration |
|-----------------|-------------------|---------------------|--------------------------|
| Tris            | 1 M               | 250 $\mu\text{l}$   | 25 mM                    |
| 10 NaCl         | 5 M               | 200 $\mu\text{l}$   | 100 mM                   |
| $\text{MnCl}_2$ | 1 M               | 100 $\mu\text{l}$   | 10 mM                    |
| TX-100          | 100 mM            | 50 $\mu\text{l}$    | 0.5 mM                   |
12. NUNC 96-well V bottom polypropylene plates (Applied Scientific Cat. No. AS-72092).
- 15 13. Ethylenediaminetetraacetic acid (EDTA).
14. Rabbit polyclonal anti-phosphotyrosine serum (Biochemistry Lab, SUGEN, Inc.).
15. Goat anti-rabbit IgG peroxidase conjugate (Biosource Cat. No. ALI0404).
- 20 16. 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid) (ABTS, Sigma Cat. No. A-1888).
17. Hydrogen peroxide 30% solution (Fisher Cat. No. H325).
18. ABTS/ $\text{H}_2\text{O}_2$ .
- 25 19. 0.2 M HCl.

#### Procedure

1. Coat Corning 96 well ELISA plates with 0.5  $\mu\text{g}$  28D4C10 in 100  $\mu\text{l}$  PBS per well, store overnight at 4° C.
2. Remove unbound 28D4C10 from wells by inverting plate to remove liquid. Wash 1x with  $\text{dH}_2\text{O}$ . Pat the plate on a paper towel to remove excess liquid.
- 30 3. Add 150  $\mu\text{l}$  of Blocking Buffer to each well. Incubate for 30 min. at room temperature with shaking.

WO 00/08202

303

PCT/US99/17845

4. Wash plate 3x with deionized water, then once with TBST. Pat plate on a paper towel to remove excess liquid and bubbles.
5. Dilute lysate in HNTG (10 µg lysate/100 µl HNTG)
6. Add 100 µl of diluted lysate to each well. Shake at room temperature for 60 min.
7. Wash plates as described in 4, above.
8. Add 80 µl working kinase buffer mix to ELISA plate containing captured PDGFR.
9. Dilute test compound 1:10 in TBS in 96-well polypropylene plates (i.e., 10 µl compound + 90 µl TBS).
10. Add 10 µl diluted test compound to ELISA plate. To control wells (wells which do not receive any test compound), add 10 µl TBS + 10% DMSO.
11. Incubate for 30 minutes while shaking at room temperature.
12. Add 10 µl ATP directly to all wells except negative control well (final well volume should be approximately 100 µl with 20 µM ATP in each well.) Incubate 30 minutes while shaking.
13. After 30 minutes, stop reaction by adding 10 µl of 200 mM EDTA (pH 8.0) to each well.
14. Wash 4x with deionized water, twice with TBST.
15. Add 100 µl anti-phosphotyrosine (1:3000 dilution in TBST) per well. Incubate 30-45 min. at room temperature, with shaking.
16. Wash as described in 4, above.
17. Add 100 µl Biosource Goat anti-rabbit IgG peroxidase conjugate (1:2000 dilution in TBST) to each well. Incubate 30 min. at room temperature, with shaking.
18. Wash as described in 4, above.
19. Add 100 µl of ABTS/H<sub>2</sub>O<sub>2</sub> solution to each well.

WO 00/08202

304

PCT/US99/17845

20. Incubate 10 to 30 minutes with shaking. Remove any bubbles.

21. If necessary stop reaction with the addition of 100  $\mu$ l 0.2 M HCl per well.

5 22. Read assay on Dynatech MR7000 ELISA reader: test filter: 410 nm, reference filter: 630 nm.

#### Biochemical FGFR Assay

This assay measures in vitro kinase activity of the Myc-GyrB-FGFR fusion protein using ELISA.

#### 10 Materials and Reagents

##### 1. HNTG

	Reagent	M.W.	5x Stock Concentration	Amount per L	1x Working Concentration
	HEPES	238.3	100 mM	23.83 g	20 mM
15	NaCl	58.44	750 mM	43.83 g	150 mM
	Glycerol	NA	50%	500 ml	10%
	Triton X-100	NA	5%	10 ml	1.0%

To make a liter of 5x stock solution, dissolve HEPES and NaCl in about 350 ml dH<sub>2</sub>O, adjust pH to 7.2 with HCl or NaOH  
 20 (depending on the HEPES that is used), add glycerol, Triton X-100 and then dH<sub>2</sub>O to volume.

2. PBS (Dulbecco's Phosphate-Buffered Saline, Gibco Catalog # 450-1300EB).

3. Blocking Buffer.

25 4. Kinase Buffer.

	Reagent	M.W.	10x Stock Concentration	1x Working Concentration
	HEPES (pH 7.2)	238.3	500 mM	50 mM
	MnCl <sub>2</sub>		20 mM	2 mM
30	MgCl <sub>2</sub>	203.32	200 mM	10 mM
	Triton-X-100		1 %	0.1%
	DTT	380.35	5 mM	0.5 mM

WO 00/08202

305

PCT/US99/17845 ,

5. Phenylmethylsulfonyl fluoride (PMSF, Sigma, Cat. No. P-7626):

Working solution: 100 mM in ethanol.

6. ATP (Bacterial source, Sigma Cat. No. A-7699)

5 Use 3.31 mg per ml MilliQ H<sub>2</sub>O for a stock concentration of 6 mM.

7. Biotin conjugated anti-phosphotyrosine mab (clone 4G10, Upstate Biotechnology Inc. Cat. No. 16-103, Ser. No. 14495).

10 8. Vectastain Elite ABC reagent (Avidin peroxidase conjugate, Vector Laboratories Cat. No. PK-6 100).

9. ABTS Solution.

10. Hydrogen peroxide 30% solution (Fisher Catalog # H325).

15 11. ABTS/H<sub>2</sub>O<sub>2</sub>.

12. 0.2 M HCl.

13. TRIS HCl (Fischer Cat. No. BP 152-5).

Prepare 1.0 mM solution in MilliQ H<sub>2</sub>O, adjust pH to 7.2 with HCl.

20 14. NaCl (Fisher Cat. No. S271-10).

Prepare 5 M solution in MilliQ H<sub>2</sub>O.

15. MgCl<sub>2</sub> (Fisher Cat. No. M33-500).

Prepare 1 M solution in MilliQ H<sub>2</sub>O.

16. HEPES (Fisher Cat. No. BP310-500).

25 Prepare 1 M solution in MilliQ H<sub>2</sub>O, adjust pH to 7.5, sterile filter.

17. TBST Buffer.

18. Sodium Carbonate Buffer (Fisher Cat. No. S495).

30 Prepare 0.1 M solution in MilliQ H<sub>2</sub>O, adjust pH to 9.6 with NaOH, filter.

19. Dithiothreitol (DTT, Fisher Cat. No. BP172-25).



WO 00/08202

306

PCT/US99/17845 ,

Prepare 0.5 mM working solution in MilliQ H<sub>2</sub>O just prior to use. Store at -20° C until used, discard any leftover.

20. MnCl<sub>2</sub>.
- 5 21. Triton X-100.
22. Goat α-Rabbit IgG (Cappel).
23. Affinity purified Rabbit α GST GyrB (Biochemistry Lab. SUGEN, Inc.).

#### Procedure

- 10 All of the following steps are conducted at room temperature unless otherwise indicated.

1. Coat Corning 96-well ELISA plates with 2 µg Goat α-Rabbit antibody per well in Carbonate Buffer such that total well volume is 100 µl. Store overnight at 4° C.
- 15 2. Remove unbound Goat α-Rabbit antibody by inverting plate to remove liquid. Pat plate on a paper towel to remove excess liquid and bubbles
3. Add 150 µl Blocking Buffer (5% Low Fat Milk in PBS) to each well. Incubate while shaking on a micro-titer plate
- 20 shaker for 30 min.
4. Wash 4x with TBST. Pat plate on a paper towel to remove excess liquid and bubbles.
5. Add 0.5 µg Rabbit α-GyrB antibody per well. Dilute antibody in DPBS to a final volume of 100 µl per well.
- 25 Incubate with shaking on a micro-titer plate shaker at room temperature for 1 hour.
6. Wash 4x with TBST as described in step 4.
7. Add 2 µg COS/FGFR cell lysate (Myc-GyrB-FGFR source) in HNTG to each well to give a final volume of 100 µl per
- 30 well. Incubate with shaking on a micro-titer plate shaker for 1 hour.
8. Wash 4X with TBST as described in step 4.

WO 00/08202

307

PCT/US99/17845 ,

9. Add 80  $\mu$ l of 1x kinase buffer per well.
10. Dilute test compound 1:10 in 1x kinase buffer + 1% DMSO in a polypropylene 96 well plate.
11. Transfer 10  $\mu$ l of diluted test compound solution and  
5 control wells from polypropylene plate wells to the corresponding ELISA plate wells, incubate with shaking on a micro-titer plate shaker for 20 minutes.
12. Add 10  $\mu$ l of 70  $\mu$ M ATP diluted in kinase buffer to positive control and test wells (Final ATP concentration is 7  
10  $\mu$ M/well). Add 10  $\mu$ l 1x kinase buffer to negative control wells. Incubate with shaking on a micro-titer plate shaker for 15 min.
13. Stop kinase reaction by adding 5  $\mu$ l 0.5 M EDTA to all wells.
14. Wash 4x with TBST as described in step 4.
15. Add 100  $\mu$ l biotin conjugated  $\alpha$ -phosphotyrosine mab (b4G10) diluted in TBST to each well. Incubate with shaking on a micro-titer plate shaker for 30 minutes.
16. Prepare Vectastain ABC reagent. Add 1 drop reagent  
20 A to 15 ml TBST. Mix by inverting tube several times. Add 1 drop reagent B and mix again.
17. Wash 4x with TBST as described in step 4.
18. Add 100  $\mu$ l ABC HRP reagent to each well. Incubate with shaking on a micro-titer plate shaker for 30 minutes.
19. Wash 4x with TBST as described in step 4.
20. Add 100  $\mu$ l of ABTS/H<sub>2</sub>O<sub>2</sub> solution to each well.
22. Incubate 5 to 15 minutes with shaking. Remove any bubbles.
23. If necessary stop reaction by adding 100  $\mu$ l of 0.2M  
30 HCl/well.
24. Read assay on Dynatech MR7000 ELISA Plate Reader, test filter: 410 nM, reference filter: 630 nM.

WO 00/08202

308

PCT/US99/17845

**Biochemical FLK-1 Assay**

This assay evaluates flk-1 autophosphorylation activity in vitro using ELISA.

**Materials And Reagents**

- 5        1.    15 cm tissue culture dishes
2.    Flk-1/NIH cells: NIH fibroblast line over-expressing  
         human flk-1 clone 3 (SUGEN, Inc., obtained from MPI,  
         Martinsried, Germany).
3.    Growth medium: DMEM plus heat inactivated 10% FBS  
10      and 2 mM Glutamine (Gibco-BRL).
4.    Starvation medium: DMEM plus 0.5% heat-inactivated  
         FBS, 2 mM Glutamine (Gibco-BRL).
5.    Corning 96-well ELISA plates (Corning Cat. No.  
         25805-96).
- 15       6.    L4 or E38 monoclonal antibody specific for flk-1,  
         Purified by Protein-A agarose affinity chromatography (SUGEN,  
         Inc.).
7.    PBS (Dulbecco's Phosphate-Buffered Saline) Gibco  
         Cat. No. 450-1300EB).
- 20       8.    HNTG (see BIOCHEMICAL FGFR for preparation).
9.    Pierce BCA protein determination kit.
10.   Blocking buffer
11.   TBST (pH 7.0)
12.   Kinase Buffer
- 25       13.   Kinase Stop Solution: 200 mM EDTA.
14.   Biotinylated 4G10, specific for phosphotyrosine  
         (UBI, Cat. No. No. 16-103).
15.   AB kit (Vector Laboratories Cat. No. PK 4000).
16.   DMSO
- 30       17.   NUNC 96-well V bottom polypropylene plates (Applied  
         Scientific Cat. No. AS-72092).
18.   Turbo-TMB (Pierce).
19.   Turbo-TMB stop solution: 1 M H<sub>2</sub>SO<sub>4</sub>.
20.   ATP (Sigma Cat. No. A-7699).

WO 00/08202

309

PCT/US99/17845 ;

21. 20% DMSO in TBS (pH 7.0).

#### Procedure

#### Cell Growth and Lysate Preparation.

1. Seed cell into growth medium and grow for 2-3 days  
5 to 90-100% confluency at 37° C and 5% CO<sub>2</sub>. Do not exceed  
passage #20.

2. Remove the medium and wash the cells twice with PBS.  
Lyse with HNTG lysis buffer. Collect all lysates and vortex  
mix them for 20-30 seconds.

10 3. Remove insoluble material by centrifugation (5-10  
min at about 10,000 xg).

4. Determine the protein concentration using BCA kit.

5. Partition lysate into 1 mg aliquots, store at -80° C.

#### Assay Procedure

15 1. Coat Corning 96-well ELISA plates with 2 µg/well  
purified L4 (or E 38) in 100 µl of PBS. Store overnight at 4° C.

2. Remove unbound proteins from wells by inverting the  
plate to remove the liquid. Wash one time with dH<sub>2</sub>O, pat plate  
on paper towel to remove excess liquid.

20 3. Block plates with 150 µl blocking buffer per well.  
Incubate for 45-60 minutes with shaking at 4° C.

4. Remove the blocking buffer and wash the ELISA plate  
three times with dH<sub>2</sub>O and one time with TBST. Pat plate on  
paper towel to remove excess liquid.

25 5. Dilute lysate in PBS to give final concentration of  
50 µg/100 µl. Add 100 µl of diluted lysate to each well.  
Incubate with shaking at 4° C overnight.

6. Remove unbound proteins from wells by inverting the  
plate. Wash as in step 4.

30 7. Add 80 µl of kinase buffer to wells (90 µl to  
negative control wells).

8. Dilute test compounds (normally 10-fold) into wells  
of a polypropylene plate containing 20% DMSO in TBS.

WO 00/08202

310

PCT/US99/17845 ,

9. Add 10  $\mu$ l of the diluted compounds to the ELISA wells containing immobilized flk-1 and shake. Control wells receive no compounds.

10. From stock 1 mM ATP, prepare 0.3 mM ATP solution in 5 dH<sub>2</sub>O (alternatively, kinase buffer may be used).

11. Add 10  $\mu$ l of 0.3 mM ATP to all wells except the negative controls. Incubate for 60 min. at room temperature with shaking.

12. After 1 hr stop the kinase reaction by adding 11  $\mu$ l 10 200 mM EDTA. Shake for 1-2 min.

13. Wash the ELISA plate 4 times with dH<sub>2</sub>O and twice with TBST.

14. Add 100  $\mu$ l of 1:5000 biotinylated 4G10:TBST to all wells. Incubate 45 min with shaking at room temperature.

15 15. While the above is incubating, add 50  $\mu$ l of solutions A & B from the ABC kit to 10 ml of TBST. These solutions must be combined approximately 30 min prior to use.

16. Wash plates as in step 4.

17. Add 100  $\mu$ l of the preformed A & B complex to all 20 wells. Incubate 30 min with shaking at room temperature.

18. Wash plates as in step 4.

19. Add 100  $\mu$ l turbo-TMB. Shake at room temperature for 10- 15 min.

20. When the color in the positive control wells reaches 25 an absorbance of about 0.35 - 0.4, stop the reaction with 100  $\mu$ l of turbo-TMB stop solution.

21. Read plates on Dynatech MR7000 ELISA reader, test filter: 450 nM, reference filter: 410 nM.

#### HUV-EC-C Assay

30 The following protocol may also be used to measure a compound's activity against PDGF-R, FGF-R, VEGF, aFGF or Flk-1/KDR, all of which are naturally expressed by HUV-EC cells.

WO 00/08202

311

PCT/US99/17845 ;

DAY 0

1. Wash and trypsinize HUV-EC-C cells (human umbilical vein endothelial cells, (American Type Culture Collection, catalogue no. 1730 CRL). Wash with Dulbecco's phosphate-buffered saline (D-PBS, obtained from Gibco BRL, catalogue no. 14190-029) 2 times at about 1 ml/10 cm<sup>2</sup> of tissue culture flask. Trypsinize with 0.05% trypsin-EDTA in non-enzymatic cell dissociation solution (Sigma Chemical Company, catalogue no. C-1544). The 0.05% trypsin is made by diluting 0.25% trypsin/1 mM EDTA (Gibco, catalogue no. 25200-049) in the cell dissociation solution. Trypsinize with about 1 ml/25-30 cm<sup>2</sup> of tissue culture flask for about 5 minutes at 37°C. After cells have detached from the flask, add an equal volume of assay medium and transfer to a 50 ml sterile centrifuge tube (Fisher Scientific, catalogue no. 05-539-6).

2. Wash the cells with about 35 ml assay medium in the 50 ml sterile centrifuge tube by adding the assay medium, centrifuge for 10 minutes at approximately 200x g, aspirate the supernatant, and resuspend with 35 ml D-PBS. Repeat the wash two more times with D-PBS, resuspend the cells in about 1 ml assay medium/15 cm<sup>2</sup> of tissue culture flask. Assay medium consists of F12K medium (Gibco BRL, catalogue no. 21127-014) and 0.5% heat-inactivated fetal bovine serum. Count the cells with a Coulter Counter® (Coulter Electronics, Inc.) and add assay medium to the cells to obtain a concentration of 0.8-1.0 x 10<sup>5</sup> cells/ml.

3. Add cells to 96-well flat-bottom plates at 100 µl/well or 0.8-1.0 x 10<sup>4</sup> cells/well, incubate ~24h at 37°C, 5% CO<sub>2</sub>.

DAY 1

1. Make up two-fold test compound titrations in separate 96-well plates, generally 50 µM on down to 0 µM. Use the same assay medium as mentioned in day 0, step 2 above.

WO 00/08202

312

PCT/US99/17845 ,

5 Titrations are made by adding 90  $\mu$ l/well of test compound at 200  $\mu$ M (4X the final well concentration) to the top well of a particular plate column. Since the stock test compound is usually 20 mM in DMSO, the 200  $\mu$ M drug concentration contains 2% DMSO.

10 A diluent made up to 2% DMSO in assay medium (F12K + 0.5% fetal bovine serum) is used as diluent for the test compound titrations in order to dilute the test compound but keep the DMSO concentration constant. Add this diluent to the remaining wells in the column at 60  $\mu$ l/well. Take 60  $\mu$ l from the 120  $\mu$ l of 200  $\mu$ M test compound dilution in the top well of the column and mix with the 60  $\mu$ l in the second well of the column. Take 60  $\mu$ l from this well and mix with the 60  $\mu$ l in the third well of the column, and so on until two-fold titrations are completed. When the next-to-the-last well is mixed, take 60  $\mu$ l of the 120  $\mu$ l in this well and discard it. Leave the last well with 60  $\mu$ l of DMSO/media diluent as a non-test compound-containing control. Make 9 columns of titrated test compound, enough for triplicate wells each for: (1) VEGF (obtained from Pepro Tech Inc., catalogue no. 100-200, (2) endothelial cell growth factor (ECGF) (also known as acidic fibroblast growth factor, or aFGF) (obtained from Boehringer Mannheim Biochemica, catalogue no. 1439 600), or, (3) human PDGF B/B (1276-956, Boehringer Mannheim, Germany) and assay media control. ECGF comes as a preparation with sodium heparin.

2. Transfer 50  $\mu$ l/well of the test compound dilutions to the 96-well assay plates containing the  $0.8-1.0 \times 10^4$  cells/100  $\mu$ l/well of the HUV-EC-C cells from day 0 and incubate ~2 h at 37° C, 5% CO<sub>2</sub>.

3. In triplicate, add 50  $\mu$ l/well of 80  $\mu$ g/ml VEGF, 20 ng/ml ECGF, or media control to each test compound condition.

WO 00/08202

313

PCT/US99/17845

As with the test compounds, the growth factor concentrations are 4X the desired final concentration. Use the assay media from day 0 step 2 to make the concentrations of growth factors. Incubate approximately 24 hours at 37°C, 5% CO<sub>2</sub>.

- 5 Each well will have 50 µl test compound dilution, 50 µl growth factor or media, and 100 µl cells, which calculates to 200 µl/well total. Thus the 4X concentrations of test compound and growth factors become 1X once everything has been added to the wells.

10 DAY 2

1. Add <sup>3</sup>H-thymidine (Amersham, catalogue no. TRK-686) at 1 µCi/well (10 µl/well of 100 µCi/ml solution made up in RPMI media + 10% heat-inactivated fetal bovine serum) and incubate ~24 h at 37°C, 5% CO<sub>2</sub>. RPMI is obtained from Gibco BRL, catalogue no. 11875-051.

DAY 3

1. Freeze plates overnight at -20°C.

DAY 4

- Thaw plates and harvest with a 96-well plate harvester (Tomtec Harvester 96®) onto filter mats (Wallac, catalogue no. 1205-401), read counts on a Wallac Betaplate™ liquid scintillation counter.

In Vivo Animal Models

Xenograft Animal Models

- 25 The ability of human tumors to grow as xenografts in athymic mice (e.g., Balb/c, nu/nu) provides a useful in vivo model for studying the biological response to therapies for human tumors. Since the first successful xenotransplantation of human tumors into athymic mice, (Rygaard and Povlsen, 1969, 30 Acta Pathol. Microbial. Scand. 77:758-760), many different human tumor cell lines (e.g., mammary, lung, genitourinary, gastro-intestinal, head and neck, glioblastoma, bone, and malignant melanomas) have been transplanted and successfully



WO 00/08202

314

PCT/US99/17845 ,

grown in nude mice. The following assays may be used to determine the level of activity, specificity and effect of the different compounds of the present invention. Three general types of assays are useful for evaluating compounds:

- 5 cellular/catalytic, cellular/biological and in vivo. The object of the cellular/catalytic assays is to determine the effect of a compound on the ability of a TK to phosphorylate tyrosines on a known substrate in a cell. The object of the cellular/biological assays is to determine the effect of a
- 10 compound on the biological response stimulated by a TK in a cell. The object of the in vivo assays is to determine the effect of a compound in an animal model of a particular disorder such as cancer.

- Suitable cell lines for subcutaneous xenograft
- 15 experiments include C6 cells (glioma, ATCC # CCL 107), A375 cells (melanoma, ATCC # CRL 1619), A431 cells (epidermoid carcinoma, ATCC # CRL 1555), Calu 6 cells (lung, ATCC # HTB 56), PC3 cells (prostate, ATCC # CRL 1435), SKOV3TP5 cells and NIH 3T3 fibroblasts genetically engineered to overexpress
- 20 EGFR, PDGFR, IGF-1R or any other test kinase. The following protocol can be used to perform xenograft experiments:

- Female athymic mice (BALB/c, nu/nu) are obtained from Simonsen Laboratories (Gilroy, CA). All animals are maintained under clean-room conditions in Micro-isolator cages with
- 25 Alpha-dri bedding. They receive sterile rodent chow and water ad libitum.

- Cell lines are grown in appropriate medium (for example, MEM, DMEM, Ham's F10, or Ham's F12 plus 5% - 10% fetal bovine serum (FBS) and 2 mM glutamine (GLN)). All cell culture media,
- 30 glutamine, and fetal bovine serum are purchased from Gibco Life Technologies (Grand Island, NY) unless otherwise specified. All cells are grown in a humid atmosphere of 90-95% air and 5-10% CO<sub>2</sub> at 37°C. All cell lines are routinely subcultured twice a week and are negative for mycoplasma as

WO 00/08202

315

PCT/US99/17845 ,

determined by the Mycotect method (Gibco).

Cells are harvested at or near confluency with 0.05% Trypsin-EDTA and pelleted at 450 x g for 10 min. Pellets are resuspended in sterile PBS or media (without FBS) to a

5 particular concentration and the cells are implanted into the hindflank of the mice (8 - 10 mice per group,  $2 - 10 \times 10^6$  cells/animal). Tumor growth is measured over 3 to 6 weeks using venier calipers. Tumor volumes are calculated as a product of length x width x height unless otherwise indicated.

10 P values are calculated using the Students t-test. Test compounds in 50 - 100  $\mu$ L excipient (DMSO, or VPD:D5W) can be delivered by IP injection at different concentrations generally starting at day one after implantation.

#### Tumor Invasion Model

15 The following tumor invasion model has been developed and may be used for the evaluation of therapeutic value and efficacy of the compounds identified to selectively inhibit KDR/FLK-1 receptor.

#### **Procedure**

20 8 week old nude mice (female) (Simonsen Inc.) are used as experimental animals. Implantation of tumor cells can be performed in a laminar flow hood. For anesthesia, Xylazine/Ketamine Cocktail (100 mg/kg ketamine and 5 mg/kg Xylazine) are administered intraperitoneally. A midline  
25 incision is done to expose the abdominal cavity (approximately 1.5 cm in length) to inject  $10^7$  tumor cells in a volume of 100  $\mu$ l medium. The cells are injected either into the duodenal lobe of the pancreas or under the serosa of the colon. The peritoneum and muscles are closed with a  
30 6-0 silk continuous suture and the skin is closed by using wound clips. Animals are observed daily.

WO 00/08202

316

PCT/US99/17845

**Analysis**

After 2-6 weeks, depending on gross observations of the animals, the mice are sacrificed, and the local tumor metastases to various organs (lung, liver, brain, stomach, spleen, heart, muscle) are excised and analyzed (measurement of tumor size, grade of invasion, immunochemistry, in situ hybridization determination, etc.).

**Measurement Of Cell Toxicity**

Therapeutic compounds should be more potent in inhibiting receptor tyrosine kinase activity than in exerting a cytotoxic effect. A measure of the effectiveness and cell toxicity of a compound can be obtained by determining the therapeutic index, i.e.,  $IC_{50}/LD_{50}$ .  $IC_{50}$ , the dose required to achieve 50% inhibition, can be measured using standard techniques such as those described herein.  $LD_{50}$ , the dosage which results in 50% toxicity, can also be measured by standard techniques as well (Mossman, 1983, J. Immunol. Methods, 65:55-63), by measuring the amount of LDH released (Korzeniewski and Callewaert, 1983, J. Immunol. Methods, 64:313, Decker and Lohmann-Matthes, 1988, J. Immunol. Methods, 115:61), or by measuring the lethal dose in animal models. Compounds with a large therapeutic index are preferred. The therapeutic index should be greater than 2, preferably at least 10, more preferably at least 50.

**B. Examples - Biological Activity.**

Examples of the in vitro potency of compounds of this invention are shown in Table 2. The examples shown are not to be construed as limiting the scope of this invention in any manner whatsoever.

30

WO 00/08202

317

PCT/US99/17845

### CONCLUSION

It will be appreciated that the compounds, methods and pharmaceutical compositions of the present invention are effective in modulating PK activity and therefore are expected to be effective as therapeutic agents against RTK, CTK-, and STK-related disorders.

One skilled in the art would also readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent herein. The molecular complexes and the methods, procedures, treatments, molecules, specific compounds described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention are defined by the scope of the claims.

It will be readily apparent to one skilled in the art that varying substitutions and modifications may be made to the invention disclosed herein without departing from the scope and spirit of the invention.

All patents and publications mentioned in the specification are indicative of the levels of those skilled in the art to which the invention pertains. All patents and publications are herein incorporated by reference to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

The invention illustratively described herein suitably may be practiced in the absence of any element or elements, limitation or limitations which is not specifically disclosed herein. Thus, for example, in each instance herein any of the terms "comprising", "consisting essentially of" and "consisting of" may be replaced with either of the other two

WO 00/08202

318

PCT/US99/17845

terms. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions of excluding any equivalents of the features shown and  
5 described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification  
10 and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the appended claims.

In addition, where features or aspects of the invention  
15 are described in terms of Markush groups, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group. For example, if X is described as selected from the group consisting of bromine, chlorine,  
20 and iodine, claims for X being bromine and claims for X being bromine and chlorine are fully described.

Other embodiments are within the following claims.

WO 00/08202

319


PCT/US99/17845

## CLAIMS

What is claimed:

1. A compound having the chemical structure:


5



wherein:

A is selected from the group consisting of carbon and  
10 nitrogen;

Q is selected from the group consisting of



15 bonds a and b may be either single or double bonds as indicated by the dotted lines with the proviso that a and b are both single bonds or both double bonds in any one compound of this invention;

20 r is 1 when a and b are double bonds;

r is 2 when a and b are single bonds;

WO 00/08202

PCT/US99/17845

320

$R^0$  is selected from the group consisting of hydrogen, alkyl,  $-C(O)R^{19}$  and  $-C(O)OR^{19}$ ;

$R^1$  is selected from the group consisting of hydrogen, alkyl, alkoxy, halo, aryl,  $-(CH_2)_nOC(O)R^{19}$ ;  $-C(O)NR^{19}$ ;  $-C(O)OR^{19}$  and  $-(CH_2)_nR^{20}$ , wherein:

$R^{19}$  is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl and aryl;

10

$n$  is 1, 2, 3 or 4;

$R^{20}$  is selected from the group consisting of hydroxy, halo,  $-C(O)OR^{19}$ ,  $-OC(O)NR^{21}R^{22}$ ;  $-OC(S)NR^{21}R^{22}$ ,  $-OC(O)NHSO_2R^{21}R^{22}$ ;  $-NR^{21}R^{22}$  and a heteroalicyclic group containing at least one nitrogen atom in the ring, the ring being bonded to the adjacent  $CH_2$  group through the nitrogen atom;

$R^2$  is selected from the group consisting of hydrogen, alkyl, cycloalkyl, aryl, heteroaryl, heteroalicyclic, trihalomethyl, alkoxy, halo,  $-(CH_2)_nR^{20}$ ,  $-SO_2NR^{21}R^{22}$ ,  $-C(O)OR^{19}$ ,  $-C(O)R^{19}$ ,  $-NHC(O)OR^{19}$ ,  $-NR^{21}R^{22}$  and  $-N=CR^{23}$  wherein

$R^{21}$  and  $R^{22}$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heteroalicyclic;

$R^{23}$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, aryl and heteroaryl;

30

WO 00/08202

PCT/US99/17845

321

R<sup>3</sup> is selected from the group consisting of hydrogen, alkyl, trihalomethyl, alkoxy, aryl, aryloxy, heteraryl, heteroalicyclic, hydroxy, halo, -SO<sub>2</sub>NR<sup>21</sup>R<sup>22</sup>, -NHSO<sub>2</sub>R<sup>19</sup>, -C(O)OR<sup>19</sup>, -NR<sup>21</sup>R<sup>22</sup> and -NHC(O)R<sup>24</sup>, wherein

5

R<sup>24</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl and heteroalicyclic;


R<sup>4</sup> is selected from the group consisting of hydrogen, alkyl, alkoxy and halo;

10

R<sup>5</sup> is selected from the group consisting of hydrogen, alkyl, cycloalkyl, halo, aryl and heteroaryl;

R<sup>6</sup> is selected from the group consisting of hydrogen, alkyl, alkoxy, halo, cycloalkyl, aryl and, combined with R<sup>18</sup>, a heteroalicyclic group having the structure

15



wherein

20

y and y' are independently selected from the group consisting of hydrogen, alkyl and aryl;

R<sup>7</sup> is OR<sup>18</sup>, wherein R<sup>18</sup> is selected from the group consisting of alkyl, -(CH)<sub>n</sub>R<sup>20</sup> and, combined with R<sup>6</sup> or R<sup>8</sup>, a heteroalicyclic group having the structure


25




WO 00/08202

PCT/US99/17845

322



$R^8$  is selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkoxy, halo, aryl, heteroaryl and, combined with  
 5  $R^{18}$ , a heteroalicyclic ring having the structure



$R^9$  is selected from the group consisting of hydrogen, alkyl, alkoxy, halo and  $-NR^{21}R^{22}$ ;

10

$R^{10}$  is selected from the group consisting of alkyl and  $-C(O)OR^{19}$ ;

$R^{11}$  is selected from the group consisting of hydrogen, alkyl  
 15 and  $-C(O)OR^{19}$ ;

$R^{12}$  is  $-(CH_2)_nR^{20}$ ;

$R^{13}$  is either a bond through which Q is bonded to the rest of  
 20 the molecule or a group selected from the group consisting of hydrogen and alkyl;

WO 00/08202

PCT/US99/17845

323

R<sup>14</sup> is either a bond through which Q is bonded to the rest of the molecule or a group selected from the group consisting of hydrogen, alkyl and  $-(CH_2)_nR^{20}$ ;


- 5 R<sup>15</sup> is either a bond through which Q is bonded to the rest of the molecule or hydrogen;

with the proviso that only one of R<sup>13</sup>, R<sup>14</sup> or R<sup>15</sup> is the bond linking Q to the rest of the molecule in any one  
10 compound herein;

R<sup>16</sup> and R<sup>17</sup>, when a and b are single bonds, are independently selected from the group consisting of hydrogen, alkyl and aryl and, when a and b are double bonds, R<sup>16</sup> is hydrogen and R<sup>17</sup> does  
15 not exist; and,

physiologically acceptable salts thereof.

2. The compound or salt of claim 1, wherein Q is  
20



3. The compound or salt of claim 2, wherein:

- 25 R<sup>1</sup> is selected from the group consisting of hydrogen, lower alkyl and  $-C(O)OH$ ;

WO 00/08202

PCT/US99/17845

324

R<sup>2</sup> is selected from the group consisting of hydrogen, lower alkyl, halo, -C(O)R<sup>19</sup>, -C(O)OR<sup>19</sup>, lower alkoxy, -NR<sup>21</sup>R<sup>22</sup>, - (CH<sub>2</sub>)<sub>n</sub>R<sup>20</sup> and -NHC(O)OR<sup>19</sup>;

5 R<sup>3</sup> is selected from the group consisting of hydrogen, hydroxy, halo, -NHC(O)O(lower alkyl), -NHSO<sub>2</sub>R<sup>19</sup>, -NHC(O)R<sup>24</sup>, trihalomethyl, and aryl optionally substituted with one or more lower alkoxy groups;

10 R<sup>4</sup> is hydrogen;

R<sup>19</sup> is selected from the group consisting of hydrogen and lower alkyl;

15 n is 2 or 3;

R<sup>20</sup> is selected from the group consisting of hydroxy, -C(O)OH, morpholin-4-yl, piperidin-1-yl, piperazin-1-yl, pyrrolidin-1-yl and -NR<sup>21</sup>R<sup>22</sup>;

20

R<sup>21</sup> and R<sup>22</sup> are independently selected from the group consisting of hydrogen and lower alkyl; and,

R<sup>24</sup> is selected from the group consisting of hydrogen and  
25 lower alkyl.

4. The compound or salt of claim 3, wherein,

R<sup>5</sup> is selected from the group consisting of hydrogen, lower  
30 alkyl, halo, five-member ring heteroaryl and aryl optionally substituted with one or more groups selected from the group consisting of lower alkyl, halo, hydroxy, -NR<sup>21</sup>R<sup>22</sup> and lower


WO 00/08202

PCT/US99/17845

325

alkoxy;


R<sup>6</sup> is selected from the group consisting of hydrogen, lower alkyl, 3 to 7-member cycloalkyl, lower alkoxy, halo, aryl  
 5 optionally substituted with one or more groups independently selected from the group consisting of lower alkyl, halo, hydroxy, -NR<sup>21</sup>R<sup>22</sup> and lower alkoxy and, combined with R<sup>18</sup>, a group having the structure



10

wherein y and y' are either both hydrogen or both lower alkyl;

R<sup>7</sup> is -OR<sup>18</sup> wherein R<sup>18</sup> is selected from the group consisting of lower alkyl, -(CH<sub>2</sub>)<sub>n</sub>R<sup>20</sup> and, combined with R<sup>6</sup> or R<sup>8</sup>, a group  
 15 having the structure




R<sup>8</sup> is selected from the group consisting of hydrogen, lower alkyl, 3 to 6-member ring cycloalkyl, lower alkoxy, halo, aryl  
 20 optionally substituted with one or more groups selected from the group consisting of lower alkyl, lower alkoxy, halo, -NR<sup>21</sup>R<sup>22</sup> and -NHC(O)(lower alkyl), five-member heteroaryl having

WO 00/08202

PCT/US99/17845

326

from 1 to 3 heteroatoms in the ring and 6-member heteroaryl having from 1 to 3 heteroatoms in the ring and, combined with  $R^{18}$  a group having the structure




- 5  $R^9$  is selected from the group consisting of hydrogen, lower alkyl; hydroxy, lower alkoxy, halo and  $-NR^{21}R^{22}$ .

5. The compound or salt of claim 3, wherein:

- 10  $R^5$  is selected from the group consisting of hydrogen, lower alkyl, thien-2-yl, thien-3-yl and aryl optionally substituted with one or more lower alkoxy groups;

- $R^6$  is selected from the group consisting of hydrogen, lower  
15 alkyl, 5 or 6-member cycloalkyl, aryl optionally substituted with one or more lower alkoxy groups and, combined with  $R^{18}$ , a group having the structure



wherein


- 20 y and y' are either both hydrogen or both lower alkyl;

WO 00/08202

PCT/US99/17845

327

$R^7$  is  $-OR^{18}$ , wherein  $R^{18}$  is selected from the group consisting of lower alkyl,  $-(CH_2)_nR^{20}$  and, combined with  $R^6$  or  $R^8$ , a group having the structure




5  $n$  is 2 or 3;

$R^{20}$  is selected from the group consisting of hydroxy,  $-C(O)OH$ , morpholin-4-yl, piperidin-1-yl, piperazin-1-yl, pyrrolidin-1-yl and  $-NR^{21}R^{22}$ ;

10

$R^8$  is selected from the group consisting of hydrogen, lower alkyl, 5 or 6-member ring cycloalkyl, lower alkoxy, aryl optionally substituted with a  $-NHC(O)$  (lower alkyl) group, thien-2-yl, thien-3-yl, pyridin-2-yl, pyridin-3-yl and,

15 combined with  $R^{18}$ , a group having the structure



$R^9$  is selected from the group consisting of hydrogen and lower alkyl.


20

WO 00/08202

PCT/US99/17845 ;

328

6. The compound or salt of claim 1 wherein, Q is:



7. The compound or salt of claim 6, wherein:

5

R<sup>1</sup> is selected from the group consisting of hydrogen, lower alkyl and  $-(CH_2)_nR^{20}$ ;

n is 2 or 3;

10

R<sup>20</sup> is selected from the group consisting of hydroxy and  $-C(O)OH$ ;

R<sup>2</sup> is selected from the group consisting of hydrogen, halo, lower alkyl, lower alkoxy,  $-SO_2NR^{21}R^{22}$  and  $-C(O)OH$ ;

15

R<sup>3</sup> is selected from the group consisting of hydrogen, halo, lower alkyl, lower alkoxy and aryl optionally substituted with one or more lower alkoxy groups; and,

20

R<sup>4</sup> is hydrogen.

8. The compound or salt of claim 7, wherein:

R<sup>10</sup> is selected from the group consisting of lower alkyl and  $-C(O)OR^{19}$ ;

25

WO 00/08202

PCT/US99/17845

329

R<sup>11</sup> is selected from the group consisting of hydrogen, lower alkyl and -C(O)OR<sup>19</sup>;

R<sup>12</sup> is -(CH<sub>2</sub>)<sub>n</sub>R<sup>20</sup>;

5

R<sup>19</sup> is selected from the group consisting of hydrogen and lower alkyl;

n is 2 or 3; and,

10

R<sup>20</sup> is selected from the group consisting of hydroxy, -C(O)OH, morpholin-4-yl, piperidin-1-yl, piperazin-1-yl, pyrrolidin-1-yl and -NR<sup>21</sup>R<sup>22</sup>.

15

9. The compound or salt of claim 7, wherein:

R<sup>10</sup> is selected from the group consisting of lower alkyl and -C(O)O(lower alkyl);

20

R<sup>11</sup> is selected from the group consisting of hydrogen, lower alkyl and -C(O)OR<sup>19</sup>;

R<sup>12</sup> is -(CH<sub>2</sub>)<sub>n</sub>R<sup>20</sup>;

25

R<sup>19</sup> is selected from the group consisting of hydrogen and lower alkyl;

n is 2; and,

30

R<sup>20</sup> is selected from the group consisting of -C(O)OH, morpholin-1-yl, piperidin-1-yl, piperazin-1-yl, pyrrolidin-1-yl and -N(lower alkyl)<sub>2</sub>.




WO 00/08202

PCT/US99/17845

330

10. The compound or salt of claim 1, wherein:



Q is

5 R<sup>1</sup> is selected from the group consisting of hydrogen and lower alkyl;

R<sup>2</sup> is selected from the group consisting of hydrogen, lower alkyl, halo, -C(O)OR<sup>19</sup>, -C(O)R<sup>19</sup>, -NR<sup>21</sup>R<sup>22</sup>, -SO<sub>2</sub>NR<sup>21</sup>R<sup>22</sup> and

10 -N=C-R<sup>23</sup>;

R<sup>3</sup> is selected from the group consisting of hydrogen, halo, lower alkyl, lower alkoxy, morpholin-4-yl, piperidin-1-yl, piperazin-1-yl, pyrrolidin-1-yl and aryl optionally

15 substituted with one or more lower alkoxy groups,; and,

R<sup>4</sup> is hydrogen.

11. The compound or salt of claim 10, wherein,

20

a and b are both single bonds;

r is 2;

25 R<sup>13</sup> is a covalent bond through which Q is bonded to the rest of the molecule;

WO 00/08202

PCT/US99/17845

331

R<sup>14</sup> is selected from the group consisting of hydrogen, lower alkyl and  $-(CH_2)_nR^{20}$ ;

R<sup>15</sup> is hydrogen;

5

R<sup>16</sup> and R<sup>17</sup> are independently selected from the group consisting of hydrogen and lower alkyl.

10        12. The compound or salt of claim 11 wherein R<sup>20</sup> is  $-NR^{21}R^{22}$  wherein R<sup>21</sup> and R<sup>22</sup> are independently selected from the group consisting of hydrogen and lower alkyl.

15        13. The compound or salt of claim 10, wherein,

a and b are both double bonds;

r is 1;

20        R<sup>13</sup> is a covalent bond through which Q is bonded to the rest of the molecule;

R<sup>14</sup> is selected from the group consisting of hydrogen, lower alkyl and  $-(CH_2)_nR^{20}$ ;

25

R<sup>15</sup> and R<sup>16</sup> are hydrogen; and,

R<sup>17</sup> does not exist.

30        14. The compound or salt of claim 10, wherein:

a and b are both double bonds;

WO 00/08202

PCT/US99/17845

332

r is 1;

R<sup>13</sup> is selected from the group consisting of hydrogen and lower  
5 alkyl;

R<sup>14</sup> is a covalent bond through which Q is bonded to the rest of  
the molecule;

10 R<sup>15</sup> and R<sup>16</sup> are hydrogen; and,

R<sup>17</sup> does not exist.

15 15. The compound or salt of claim 10, wherein:

a and b are double bonds;

r is 1;

20 R<sup>13</sup> is selected from the group consisting of hydrogen and lower  
alkyl;

R<sup>14</sup> is selected from the group consisting of hydrogen, lower  
alkyl and  $-(CH_2)_nR^{20}$ ;

25 R<sup>15</sup> is a covalent bond through which Q is bonded to the rest of  
the molecule;

R<sup>16</sup> is hydrogen; and,

30 R<sup>17</sup> does not exist.

WO 00/08202

PCT/US99/17845 ,

333

16. A pharmaceutical composition, comprising
- a compound or salt of claim 1; and,
- 5 a pharmaceutically acceptable carrier or excipient.

17. A method for the modulation of the catalytic activity of a protein kinase comprising contacting said protein kinase with a compound or salt of claim 1.

10

18. The method of claim 17 wherein said protein kinase is selected from the group consisting of a receptor tyrosine kinase, a non-receptor tyrosine kinase and a serine-threonine kinase.

15

19. A method for treating or preventing a protein kinase related disorder in an organism comprising administering a therapeutically effective amount of a compound or salt of claim 1 to said organism.

20

20. The method of claim 19 wherein said protein kinase related disorder is selected from the group consisting of a receptor tyrosine kinase related disorder, a non-receptor tyrosine kinase related disorder and a serine-threonine
- 25 kinase related disorder.

21. The method of claim 19 wherein said protein kinase related disorder is selected from the group consisting of an EGFR related disorder, a PDGFR related disorder, an IGFR
- 30 related disorder and a flk related disorder.

WO 00/08202

PCT/US99/17845

334

22. The method of claim 19 wherein said protein kinase related disorder is a cancer selected from the group consisting of squamous cell carcinoma, astrocytoma, Kaposi's sarcoma, glioblastoma, lung cancer, bladder cancer, head and neck cancer, melanoma, ovarian cancer, prostate cancer, breast cancer, small-cell lung cancer, glioma, colorectal cancer, genitourinary cancer and gastrointestinal cancer.

23. The method of claim 19 wherein said protein kinase related disorder is selected from the group consisting of diabetes, an autoimmune disorder, a hyperproliferation disorder, restenosis, fibrosis, psoriasis, von Heppel-Lindau disease, osteoarthritis, rheumatoid arthritis, angiogenesis, an inflammatory disorder, an immunological disorder and a cardiovascular disorder.

24. The method of claim 19 wherein said organism is a human.

25. A compound of claim 1, or salt thereof, selected from the group consisting of:  
3-(3,5-Diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one  
5-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one  
N-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide  
3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-hydroxy-1,3-dihydroindol-2-one  
5-Acetyl-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one  
3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid methyl ester

WO 00/08202

PCT/US99/17845 ,

335

3- (3-Isopropyl-4-methoxybenzylidene) -1,3-dihydroindol-2-one  
3- (5-Isopropyl-4-methoxy-2-methylbenzylidene) -1,3-dihydroindol-2-one  
5-Chloro-3- (5-isopropyl-4-methoxy-2-methylbenzylidene) -1,3-dihydroindol-2-one  
3- (3-Cyclopentyl-4-methoxybenzylidene) -1,3-dihydroindol-2-one  
3- (3-Cyclopentyl-4-methoxybenzylidene) -5-fluoro-1,3-dihydroindol-2-one  
3- (3-Cyclohexyl-4-methoxybenzylidene) -1,3-dihydroindol-2-one  
5-Bromo-3- (6-methoxybiphenyl-3-ylmethylene) -1,3-dihydroindol-2-one  
5-Chloro-3- (2,3-dihydrobenzofuran-5-ylmethylene) -1,3-dihydroindol-2-one  
5-Chloro-3- (2,2-dimethylchroman-6-ylmethylene) -1,3-dihydroindol-2-one  
N- {3- [3-Cyclohexyl-4- (2-morpholin-4-ylethoxy) -benzylidene] -2-oxo-2,3-dihydro-1H-indol-6-yl} -acetamide  
3- (3,5-Diisopropyl-4-methoxybenzylidene) -5-methoxy-1,3-dihydroindol-2-one  
N- [3- (4-Methoxy-3-thiophen-3-ylbenzylidene) -2-oxo-2,3-dihydro-1H-indol-6-yl] -acetamide  
3- (3,5-Diisopropyl-4-methoxybenzylidene) -5-methyl-1,3-dihydroindol-2-one  
5-Amino-3- (3,5-diisopropyl-4-methoxybenzylidene) -1,3-dihydroindol-2-one  
5-Chloro-3- (4-methoxy-3,5-dimethylbenzylidene) -1,3-dihydroindol-2-one  
3- (3,5-Diisopropyl-4-methoxybenzylidene) -6-fluoro-1,3-dihydroindol-2-one  
3- (2,2-Dimethylchroman-6-ylmethylene) -5-fluoro-1,3-dihydroindol-2-one  
5-Chloro-3- [3,5-diisopropyl-4- (2-morpholin-4-ylethoxy) -

WO 00/08202

PCT/US99/17845 ;

336

benzylidene]-1,3-dihydroindol-2-one  
3-(3,5-Diisopropyl-4-methoxybenzylidene)-7-fluoro-1,3-dihydroindol-2-one  
3-(4-Methoxy-3-thiophen-3-ylbenzylidene)-5-(2-morpholin-4-ylethyl)-1,3-dihydroindol-2-one  
N-[3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide  
3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-ethyl-1,3-dihydroindol-2-one  
N-[2'-Methoxy-5'-(2-oxo-1,2-dihydroindol-3-ylidenemethyl)-biphenyl-3-yl]-acetamide  
5-Fluoro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one  
N-[3-(4-Methoxy-3-thiophen-2-ylbenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide  
6-Amino-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one  
N-[3-(2,2-Dimethylchroman-6-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide  
5-Bromo-3-(2,2-dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one  
3-(4-Methoxy-3-thiophen-3-ylbenzylidene)-1,3-dihydroindol-2-one  
5-Bromo-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydroindol-2-one  
5-Fluoro-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one  
3-(3-Isopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one  
3-(4,5-Dimethoxy-2-thiophen-2-ylbenzylidene)-1,3-dihydroindol-2-one  
N-{3-[4-(2-Morpholin-4-ylethoxy)-3-thiophen-2-ylbenzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide

WO 00/08202

PCT/US99/17845 ,

337

3-(2,2-Dimethylchroman-6-ylmethylene)-4-methyl-1,3-dihydroindol-2-one

3-(2,3-Dihydrobenzofuran-5-ylmethylene)-5-fluoro-1,3-dihydroindol-2-one

3-(3-Cyclohexyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one

5-Fluoro-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one

3-(3'-Ethoxy-6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one

3-(3-Cyclopentyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one

3-(3-Cyclopentyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

3-(4,5,2'-Trimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one

N-{3-[4-(2-Morpholin-4-ylethoxy)-3-thiophen-3-ylbenzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide

5-Chloro-3-(3-cyclohexyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-carbamic acid tert-butyl ester

3-(3,5-Diisopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

5-Bromo-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

N-{3-[3-tert-Butyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-2-oxo-2,3-dihydro-1H-indol-6-yl}-acetamide

3-(4-Methoxy-3,5-dimethylbenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one



WO 00/08202

PCT/US99/17845

338

5-Bromo-3-[3,5-diisopropyl-4-(2-morpholin-4-ylethoxy)-benzylidene]-1,3-dihydroindol-2-one

3-(3'-Ethoxy-4,5-dimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one

5-Chloro-3-(4-methoxy-3-thiophen-2-ylbenzylidene)-1,3-dihydroindol-2-one

5-Chloro-3-(4-methoxy-3-pyridin-3-ylbenzylidene)-1,3-dihydroindol-2-one

5-Chloro-3-(4,5,3'-trimethoxybiphenyl-2-ylmethylene)-1,3-dihydroindol-2-one

3-(4,5-Dimethoxy-2-naphthalen-2-ylbenzylidene)-1,3-dihydroindol-2-one

N-[3-(3'-Acetylamino-6-methoxybiphenyl-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-acetamide

6-Methoxy-3-(4-methoxy-3-thiophen-3-ylbenzylidene)-1,3-dihydroindol-2-one

3-(6-Methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one

3-(2,3-Dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one

5-Chloro-3-(6-methoxybiphenyl-3-ylmethylene)-1,3-dihydroindol-2-one

3-(3-Cyclohexyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

3-(2,3-dihydrobenzofuran-5-ylmethylene)-4-methyl-1,3-dihydroindol-2-one

3-(3-Isopropyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one

3-(6-Methoxybiphenyl-3-ylmethylene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one

3-(3-Cyclohexyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one

3-(2,3-Dihydrobenzofuran-5-ylmethylene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one

WO 00/08202

PCT/US99/17845 ,

339

3-(3,5-Diisopropyl-4-methoxybenzylidene)-1,3-dihydropyrrolo[2,3-b]pyridin-2-one

5-Bromo-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

5-Bromo-3-(3-cyclopentyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

5-Chloro-3-(3-cyclopentyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

5-Chloro-3-(6-methoxybiphenyl-3-ylmethylene)-4-methyl-1,3-dihydroindol-2-one

5-Chloro-3-(5-isopropyl-4-methoxy-2-methylbenzylidene)-4-methyl-1,3-dihydroindol-2-one

5-Chloro-3-(4-methoxy-3,5-dimethylbenzylidene)-4-methyl-1,3-dihydroindol-2-one

3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-trifluoromethyl-1,3-dihydroindol-2-one

6-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

3-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-5-yl]-propionic acid

3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-methoxy-1,3-dihydroindol-2-one

5-Butyl-3-(3,5-diisopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid

3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-(3-methoxyphenyl)-1,3-dihydroindol-2-one

7-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-5-methyl-1,3-dihydroindol-2-one

[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-5-yl]-carbamic acid tert-butyl ester

WO 00/08202

PCT/US99/17845

340

5-Chloro-3-(3-isopropyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

5-Chloro-3-(3-cyclopentyl-4-methoxybenzylidene)-1,3-dihydroindol-2-one

3-(6-Methoxybiphenyl-3-ylmethylene)-4-methyl-1,3-dihydroindol-2-one

3-(5-Isopropyl-4-methoxy-2-methylbenzylidene)-4-methyl-1,3-dihydroindol-2-one

5-Bromo-3-(2,3-dihydrobenzofuran-5-ylmethylene)-1,3-dihydroindol-2-one

5-Chloro-3-(3-isopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

5-Chloro-3-(3,5-diisopropyl-4-methoxybenzylidene)-4-methyl-1,3-dihydroindol-2-one

5-Chloro-3-(2,2-dimethylchroman-6-ylmethylene)-4-methyl-1,3-dihydroindol-2-one

3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid

3-(3,5-Diisopropyl-4-methoxybenzylidene)-5,6-dimethoxy-1,3-dihydroindol-2-one

N-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-methanesulfonamide

N-[3-(3,5-Diisopropyl-4-methoxybenzylidene)-2-oxo-2,3-dihydro-1H-indol-6-yl]-benzamide

3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-(3-ethoxyphenyl)-1,3-dihydroindol-2-one

3-(3,5-Diisopropyl-4-methoxybenzylidene)-6-phenyl-1,3-dihydroindol-2-one

3-(3,5-Diisopropyl-4-methoxybenzylidene)-5-fluoro-1,3-dihydroindol-2-one

5-Fluoro-3-(4-methoxy-3,5-dimethylbenzylidene)-1,3-dihydroindol-2-one

WO 00/08202

PCT/US99/17845 ,

341

3-(2,2-dimethylchroman-6-ylmethylene)-1,3-dihydroindol-2-one

26. A compound of claim 1, or a physiologically acceptable salt thereof, selected from the group consisting of:

- 5-methyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,
- 3-(3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-5-sulfonic acid amide,
- 3-(3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide,
- 3-(3-methyl-1H-indole-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide,
- 3-(3-methyl-1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid,
- 5-acetyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,
- 5-acetyl-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,
- 3-(1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indol-5-sulfonic acid amide,
- 5-amino-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,
- 3-(1H-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-carboxylic acid,
- 6-chloro-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,
- 3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,
- 5-chloro-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,
- 5-bromo-3-(1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,
- 3-(1H-indol-2-ylmethylene)-4-methyl-1,3-dihydroindol-2-one,
- 3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,
- 5-chloro-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-dihydroindol-2-one,
- 5-bromo-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-

WO 00/08202

PCT/US99/17845 ,

342

dihydroindol-2-one,  
4-methyl-3-(3-methyl-1H-indol-2-ylmethylene)-1,3-  
dihydroindol-2-one, and  
3-(1H-indol-2-ylmethylene)-5[(1H-indol-2-ylmethylene)amino]-  
5 1,3-dihydroindol-2-one.

27. A compound of claim 1, or a physiologically acceptable salt thereof, wherein said compound is selected from the group consisting of:

10 3-[2-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
3-[2-(5-chloro-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
3-2-(5-bromo-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-  
15 4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
3-[2-(4-methyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
3-[2-(5-methyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
20 3-[2-(6-chloro-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
3-[2-(6-methoxy-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
N,N-dimethyl-3-[2-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-  
25 4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide,  
3-[3-(3-dimethylamino-propyl)-4,5,6,7-tetrahydro-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one,  
3-[2-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide,  
30 3-[3-(3-morpholin-4-yl-3-oxo-propyl)-4,5,6,7-tetrahydro-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one,  
N-methyl-3-[2-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-

WO 00/08202

PCT/US99/17845

343

- 4,5,6,7-tetrahydro-1H-indol-3-yl]-propionamide,  
N-(2-morpholin-4-yl-ethyl)-3-[2-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-  
propionamide, 3-[2-(2-oxo-1,2-dihydro-pyrrolo[2,3-b]pyridin-  
5 3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic  
acid,  
3-{2-[6-(3-methoxy-phenyl)-2-oxo-1,2-dihydro-indol-3-ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl}-propionic  
acid,  
10 3-{2-[6-(4-methoxy-phenyl)-2-oxo-1,2-dihydro-indol-3-ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl}-propionic  
acid,  
3-[2-(2-oxo-6-phenyl-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic acid,  
15 3-{2-[6-(2-methoxy-phenyl)-2-oxo-1,2-dihydro-indol-3-ylidenemethyl]-4,5,6,7-tetrahydro-1H-indol-3-yl}-propionic  
acid,  
3-[2-(5-isopropylsulfamoyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic  
20 acid,  
3-[2-(6-morpholin-4-yl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic  
acid,  
3-[2-(5-chloro-4-methyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic  
25 acid,  
3-[2-(5-bromo-4-methyl-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-propionic  
acid,  
30 3-[2-(5-bromo-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-4,5,6,7-tetrahydro-1H-indol-3-yl]-N-(2-morpholin-4-yl-ethyl)-propionamide,

WO 00/08202

PCT/US99/17845

344

3-[2-(5-chloro-2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-  
4,5,6,7-tetrahydro-1H-indol-3-yl]-N-(2-morpholin-4-yl-  
ethyl)-propionamide, and  
3-[2-(2-oxo-1,2-dihydro-indol-3-ylidenemethyl)-phenyl]-  
5 propionic acid.

28. A process for the preparation of a 3-(3-( $R^{27}R^{28}N$ )propyl)-4,5,6,7-tetrahydro-1H-indole, comprising:  
reacting 5-aminolevulinic acid or its hydrochloride  
10 salt with 1,3-cyclohexadione, optionally substituted in the  
5-position with one or two groups independently selected  
from the group consisting of alkyl and aryl, in a first  
solvent in the presence of a base at a temperature of from  
about 60° to about 180° C to give an optionally substituted  
15 3-(4-oxo-4,5,6,7-tetrahydro-1H-indol-3-yl)propionic acid;  
reacting said 3-(4-oxo-4,5,6,7-tetrahydro-1H-indol-3-  
yl)propionic acid with an  $R^{27}R^{28}NH$  group in which  $R^{27}$  and  $R^{28}$   
are independently selected from the group consisting of  
hydrogen, alkyl, alkenyl, alkynyl and aryl or, combined,  $R^{27}$   
20 and  $R^{28}$  form a five-member heteroaryl group with the nitrogen  
atom or form a five-member or a six-member heteroalicyclic  
group with the nitrogen atom, to give an optionally  
substituted 3-(3-( $R^{27}R^{28}N$ )-3-oxopropyl)-4,5,6,7-tetrahydro-  
indol-4-one; and,  
25 reducing said 3-(3-( $R^{27}R^{28}N$ )-3-oxopropyl)-4,5,6,7-  
tetrahydroindol-4-one with a metal hydride reducing agent in  
a second solvent at a temperature of from about room  
temperature to about 80° C to give a 3-(3-( $R^{27}R^{28}N$ )propyl)-  
4,5,6,7-tetrahydro-1H-indole.

30

29. The process of claim 28 wherein reacting said 5-aminolevulinic acid or its hydrochloride salt with said

WO 00/08202

PCT/US99/17845 ,

345

optionally substituted 1,3-cyclohexadione comprises using water as said first solvent.

30. The process of claim 28 wherein said reacting said  
5 5-aminolevulinic acid or its hydrochloride salt with said optionally substituted 1,3-cyclohexadione comprises using sodium acetate as said base.

31. The process of claim 28, wherein reacting said 5-  
10 aminolevulinic acid or its hydrochloride salt with said optionally substituted 1,3-cyclohexadione comprises using a temperature of from about 100° C to about 120° C.

32. The process of claim 28 comprising using as R<sup>27</sup> and  
15 R<sup>28</sup> groups independently selected from the group consisting of lower alkyl and, combined, -(CH<sub>2</sub>)<sub>4</sub>-, -(CH<sub>2</sub>)<sub>5</sub>-, -(CH<sub>2</sub>)<sub>2</sub>O(CH<sub>2</sub>)<sub>2</sub>- and -(CH<sub>2</sub>)<sub>2</sub>N(CH<sub>2</sub>)<sub>2</sub>-.

33. The process of claim 28 in which said reaction of  
20 said R<sup>27</sup>R<sup>28</sup>NH with said 3-(4-oxo-4,5,6,7-tetrahydro-1H-indol-3-yl)propionic acid comprises using 1,1-carbonyldiimidazole.

34. The process of claim 28 in which reducing said 3-  
(3-(R<sup>27</sup>R<sup>28</sup>N)-3-oxopropyl)-4,5,6,7-tetrahydroindol-4-one  
25 comprises using lithium aluminum hydride.

35. The process of claim 28 in which reducing said 3-  
(3-(R<sup>27</sup>R<sup>28</sup>N)-3-oxopropyl)-4,5,6,7-tetrahydroindol-4-one  
comprises using tetrahydrofuran as said second solvent at a  
30 temperature of from about room temperature to the reflux temperature of said tetrahydrofuran.